

Ratified by HoC/WHO at the HoC meeting in Brisbane, October 2002



WORLD HEALTH ORGANIZATION

Update Reference Committee

30 November 2002

OFFICIAL UPDATES TO ICD-10

The following pages include the official changes to the tabular list, instruction manual and alphabetical index of ICD-10. These changes were approved in principle at the HoC meeting in Brisbane, Australia in October 2002. Each change has been uniquely identified and the source for each change has been indicated. The implementation date has also been documented.

(Note: Every effort has been made in the following pages to reproduce the content of the ICD-10 in the same format as the published volumes. Page references have not been used in all instances since these do not apply to electronic versions of the Classification. Additions/changes have been indicated through the use of instructions, underline and ~~strikeout~~).

This document is not issued to the general public, and all rights are reserved by the World Health Organization (WHO). The document may not be reviewed, abstracted, quoted, reproduced or translated, in part or in whole, without the prior written permission of WHO. No part of this document may be stored in a retrieval system or transmitted in any form or by any means – electronic, mechanical or other – without the prior written permission of WHO.

Volume 1

TABULAR LIST

Instruction	Tabular list entries	Source	Major/Minor update	Implementation date
Page 50 Modify category descriptor	<p>List of three character categories</p> <p>Cerebral palsy and other paralytic syndromes (G80-G83) G80 Infantile <u>Cerebral palsy</u></p>	Australia (URC:0045)	Major	January 2006
Page 71 Revise code Add category code and title	<p>Three character categories</p> <p>Other obstetric conditions, not elsewhere classified (O94-O99) O94 <u>Sequelae of complication of pregnancy, childbirth and the puerperium</u></p>	Australia (URC:0112)	Major	January 2006
Page 107 Add instruction	<p>Certain infectious and parasitic diseases (A00-B99)</p> <p><u>Use additional code (U80.- – U89.-), if desired, to identify the antibiotic to which a bacterial agent is resistant</u></p>	Australia/ NORDIC (URC:0026)	Major	January 2006
Add inclusion term Add excludes notes	<p>A98.5 Haemorrhagic fever with renal syndrome <u>Hantavirus disease with renal manifestations</u></p> <p><i>Excludes:</i> hantavirus (cardio)-pulmonary syndrome (B33.4† J17.1*)</p>	Brazil (URC:0042)	Major	January 2006
Add new code and description Add inclusion terms Add instructional note Add excludes note	<p>B33 Other viral diseases, not elsewhere classified</p> <p><u>B33.4† Hantavirus (cardio)-pulmonary syndrome [HPS] [HCPS] (J17.1*)</u></p> <p><u>Hantavirus disease with pulmonary manifestations</u> <u>Sin Nombre virus disease</u></p> <p><u>Use additional code (N17.9) , if desired, to identify any renal failure associated with HPS caused by the Andes, Bayou and Black Creek Canal hantavirus aetiologies.</u></p> <p><i>Excludes:</i> haemorrhagic fever with renal manifestations (A98.5† N08.0*)</p>	Brazil (URC:0042)	Major	January 2006

Delete inclusion term	C71.0 Cerebrum, except lobes and ventricles Corpus callosum	UK (URC: 0015)	Minor	January 2004
Page 321 Add excludes note	.0 Acute intoxication <i>Excludes:</i> intoxication meaning poisoning (T36-T50)	MRG (URC:0116)	Major	January 2006
Add exclusion note Delete exclusion note	G11 Hereditary ataxia <i>Excludes:</i> cerebral palsy (G80.-) infantile cerebral palsy (G80.-)	Australia (URC:0045)	Major	January 2006
Modify title Delete includes note Modify title Add inclusion term Modify title Add inclusion term Modify title Add inclusion term Modify title Modify title	G80 Infantile Cerebral palsy <i>Includes:</i> Little's disease <i>Excludes:</i> hereditary spastic paraplegia (G11.4) G80.0 Spastic quadriplegic cerebral palsy <u>Spastic tetraplegic cerebral palsy</u> G80.1 Spastic diplegic cerebral palsy <u>Spastic cerebral palsy NOS</u> G80.2 Infantile hemiplegia Spastic hemiplegic cerebral palsy G80.3 Dyskinetic cerebral palsy <u>Dystonic cerebral palsy</u> G80.4 Ataxic cerebral palsy G80.8 Other infantile cerebral palsy G80.9 Infantile Cerebral palsy, unspecified	Australia (URC:0045)	Major	January 2006
Modify excludes note	G81 Hemiplegia <i>Excludes:</i> congenital and infantile cerebral palsy (G80.-)	Australia (URC:0045)	Major	January 2006

Modify excludes note	G82 Paraplegia and tetraplegia <i>Excludes:</i> congenital and infantile cerebral palsy (G80.-)	Australia (URC:0045)	Major	January 2006
Delete exclusion term	K56 Paralytic ileus and intestinal obstruction without hernia <i>Excludes:</i> neonatal intestinal obstructions classifiable to P76.-	MRG (URC:0115)	Minor	January 2004
Add excludes note	K56.6 Other and unspecified intestinal obstruction <i>Excludes:</i> <u>other and unspecified neonatal intestinal obstruction classifiable to P76.8, P76.9</u>			
Delete inclusion term	K62.8 Other specified diseases of anus and rectum Perforation (nontraumatic) of rectum	Germany (URC:0140)	Minor	January 2004
Add excludes note	Diseases of the musculoskeletal system and connective tissue (M00-M99) <i>Excludes:</i> <u>certain disorders of the temporomandibular joint (K07.6)</u>	Germany (URC:0029)	Minor	January 2004
Add note	N39.3 Stress incontinence <u>Use additional code (N32.8), if desired, to identify overactive bladder or destrusor muscle hyperactivity</u>	North America (URC:0040)	Major	January 2006
Add note	N39.4 Other specified urinary incontinence <u>Use additional code (N32.8), if desired, to identify overactive bladder or destrusor muscle hyperactivity</u>	North America (URC:0040)	Major	January 2006
Page 721 Revise code	Chapter XV Pregnancy, childbirth and the puerperium (O00-O99) This chapter contains the following blocks: <u>O94-O99 Other obstetric conditions, not elsewhere classified</u>	Australia (URC:0112)	Major	January 2006

<p>Page 760, Revise code Add subcategory</p> <p>Add instructional note</p> <p>Add excludes note</p>	<p>Other obstetric conditions, not elsewhere classified (O94-O99)</p> <p><u>O94 Sequelae of complication of pregnancy, childbirth and the puerperium</u></p> <p><i>Note:</i> This category is to be used for morbidity coding only to indicate conditions in categories O00 – O75 and O85 – O92 as the cause of sequelae, which are themselves classified elsewhere. The ‘sequelae’ include conditions specified as such or as late effects, or those present one year or more after the onset of the causal condition.</p> <p><i>Excludes:</i> that resulting in death (O96, O97)</p>	<p>Australia (URC:0112)</p>	<p>Major</p>	<p>January 2006</p>
<p>Delete exclusion note</p> <p>Add excludes note</p>	<p>P76 Other intestinal obstruction of newborn <i>Excludes:</i> intestinal obstruction classifiable to K56.</p> <p>P76.8 Other specified intestinal obstruction of newborn <i>Excludes:</i> intestinal obstruction classifiable to K56.0-K56.5</p>	<p>MRG (URC:0115)</p>	<p>Minor</p>	<p>January 2004</p>
<p>Modify subcategory title</p>	<p>S05.9 Injury of eye and orbit, part unspecified</p>	<p>Dutch Committee on Translation of ICD-10 (URC: 0082)</p>	<p>Minor</p>	<p>January 2004</p>
<p>Delete exclusion term</p> <p>Add exclusion term</p>	<p>Poisoning by drugs, medicaments and biological substances (T36-T50)</p> <p><i>Excludes:</i> drug dependence and related mental and behavioural disorders due to psychoactive substance use (F10-F19) intoxication meaning inebriation (F10-F19)</p>	<p>MRG (URC:0116)</p>	<p>Major</p>	<p>January 2006</p>
<p>Delete exclusion term</p> <p>Add exclusion term</p>	<p>T40 Poisoning by narcotics and psychodysleptics</p> <p><i>Excludes:</i> drug dependence and related mental and behavioural disorders due to psychoactive substance use (F10-F19) intoxication meaning inebriation (F10-F19)</p>	<p>MRG (URC:0116)</p>	<p>Major</p>	<p>January 2006</p>

<p>Delete exclusion term</p> <p>Add exclusion term</p>	<p>T42 Poisoning by antiepileptic, sedative-hypnotic and antiparkinsonism drugs</p> <p><i>Excludes:</i> drug dependence and related mental and behavioural disorders due to psychoactive substance use (F10-F19) <u>intoxication meaning inebriation (F10-F19)</u></p>	<p>MRG (URC:0116)</p>	<p>Major</p>	<p>January 2006</p>
<p>Delete exclusion term</p> <p>Add exclusion term</p>	<p>T43 Poisoning by psychotropic drugs, not elsewhere classified</p> <p><i>Excludes:</i> drug dependence and related mental and behavioural disorders due to psychoactive substance use (F10-F19) <u>intoxication meaning inebriation (F10-F19)</u></p>	<p>MRG (URC:0116)</p>	<p>Major</p>	<p>January 2006</p>
<p>Modify excludes note</p>	<p>T44.5 Predominantly β-adrenoreceptor agonists, not elsewhere classified</p> <p><i>Excludes:</i> salbutamol <u>β-adrenoreceptor agonists used in asthma therapy (T48.6)</u></p>	<p>Australia (URC:0076)</p>	<p>Major</p>	<p>January 2006</p>
<p>Add inclusion term</p> <p>Modify excludes note</p>	<p>T48.6 Antiasthmatics, not elsewhere classified</p> <p><u>β-adrenoreceptor agonists used in asthma therapy</u></p> <p><i>Excludes:</i> <u>β-adrenoreceptor agonists not used in asthma therapy (T44.5)</u></p>	<p>Australia (URC:0076)</p>	<p>Major</p>	<p>January 2006</p>

Add block	<u>Bacterial agents resistant to antibiotics (U80-U89)</u>	Australia/ NORDIC (URC:0026)	Major	January 2006
Add note	<u>Note: These categories should never be used in primary coding. They are provided for use as supplementary or additional codes when it is desired to identify the antibiotic to which a bacterial agent is resistant, in bacterial infection classified elsewhere</u>			
Add category, codes and code descriptions	<u>U80 Agent resistant to penicillin and related antibiotics</u> <u>U80.0 Penicillin resistant agent</u> <u>U80.1 Methicillin resistant agent</u> <u>U80.8 Agent resistant to other penicillin-related antibiotic</u>			
Add category, codes and code descriptions	<u>U81 Agent resistant to vancomycin and related antibiotics</u> <u>U81.0 Vancomycin resistant agent</u> <u>U81.8 Agent resistant to other vancomycin-related antibiotic</u>			
Add category	<u>U88 Agent resistant to multiple antibiotics</u>			
Add note	<u>Note: This category is provided for use when a bacterial agent is resistant to two or more antibiotics but there is insufficient detail to determine which antibiotic is contributing most to the “main condition”. It should also be used for primary tabulation purposes when it is more convenient to record a single code; otherwise each specific antibiotic-resistant agent should be coded separately.</u>			
Add category, codes and code descriptions	<u>U89 Agent resistant to other and unspecified antibiotics</u> <u>U89.8 Agent resistant to other single specified antibiotic</u> <u>U89.9 Agent resistant to unspecified antibiotic</u>			
Modify code title	X25 Contact with other specified venomous arthropods	Dutch Committee on Translation of ICD-10 (URC: 0082)	Major	January 2006

Instruction	Instruction manual entries	Source	Major/Minor update	Implementation date
Page vi Add contents reference to new section in Volume 2	5a. Recommendations 138a	MRG (URC:0113)	Major	January 2006
Page 42 Add text Add text	<p>4.1.9 The modification rules</p> <p>Rule B. Trivial conditions Where the selected cause is a trivial condition unlikely to cause death and a more serious condition (<u>any condition except an ill-defined or another trivial condition</u>) is reported, reselect the underlying cause... ..of the trivial condition, select the adverse reaction.</p> <p><u>When a trivial condition is reported as causing any other condition, the trivial condition is not discarded, i.e. Rule B is not applicable.</u></p>	MRG (URC:0114)	Major	January 2006
Page 45 Change existing text as indicated. Change existing text as indicated Change existing text as indicated	<p>Rule B. Trivial conditions (A) Where the selected cause is a trivial condition unlikely to cause death and a more serious condition (<u>any condition except an ill-defined or another trivial condition</u>) is reported, reselect the underlying cause as if the trivial condition has not been reported. If the death was the result of an adverse reaction to treatment of the trivial condition, select the adverse reaction.</p> <p><i>Example 38:</i> I (a) Dental caries II Cardiac arrest Diabetes</p> <p>Code to cardiac arrest (I46.9) diabetes (E14.9). Dental caries, selected by the General Principle, is ignored.</p> <p><i>Example 39:</i> (no change to existing example)</p> <p><u>(B) If the death was the result of an adverse reaction to treatment of the trivial condition, select the adverse reaction.</u></p> <p><i>Example 40:</i> I (a) Intraoperative haemorrhage</p>	MRG (URC:0114)	Major	January 2006

<p>Change existing text as indicated</p>	<p>(b) Tonsillectomy (c) Hypertrophy of tonsils</p> <p>Code to haemorrhage during surgical operation (Y60.0). <u>Code to the adverse reaction to treatment of the hypertrophy of tonsils, selected by the General Principle.</u></p>			
<p>Change existing text as indicated</p>	<p><u>(C) When a trivial condition is reported as causing any other condition, the trivial condition is not discarded (i.e. Rule B is not applicable).</u></p> <p><i>Example 41:</i> I (a) Bursitis and ulcerative colitis <u>Septicaemia</u> (b) <u>Impetigo</u></p>			
<p>Change existing text as indicated</p>	<p><u>Code to ulcerative colitis (K51.9). Bursitis, selected by Rule 2 (see Example 20), is ignored.</u> <u>Code to impetigo (L01.0) The trivial condition selected by the General Principle is not discarded since it is reported as the cause of</u> <u>another condition.</u></p> <p><i>Example 42:</i> I (a) Paronychia <u>Respiratory insufficiency</u> H (b) Tetanus <u>Upper respiratory infection</u></p> <p><u>Code to tetanus (A35). Paronychia, selected by the General Principle, is ignored.</u> <u>Code to upper respiratory infection (J06.9). The trivial condition selected by the General Principle is not discarded since it is reported as the cause of another condition.</u></p>			
<p>Page 52 Add text Delete text</p>	<p>F10-F19 Mental and behavioural disorders due to psychoactive substance use</p> <p><i>with mention of:</i> <u>X40-X49 Accidental poisoning by and exposure to noxious substances, code X40-X49</u> <u>X60-X69 Intentional self-poisoning by and exposure to noxious substances, code X60-X69</u> <u>X85-X90 Assault by noxious substances, code X85-X90</u> <u>Y10-Y19 Poisoning by and exposure to drugs, chemicals and noxious substances, code Y10-Y19</u></p> <p>Fourth characters .0 (Acute intoxication) and .5 (Psychotic disorder)</p>	<p>MRG (URC:0117)</p>	<p>Major</p>	<p>January 2006</p>

<p>Add text</p>	<p>with mention of Dependence syndrome (.2), code F10-F19 with fourth character .2</p> <p><u>Fourth character .0 (Acute intoxication), code X40-X49, X60-X69, X85-X90 or Y10-Y19</u> <u>Fourth character .5 (Psychotic disorder) with mention of Dependence syndrome (.2), code F10-F19 with fourth character .2</u></p>			
<p>Page 61 Add text Add text</p>	<p><u>P70.3 – P72.0 Transitory endocrine and metabolic disorders specific to fetus and newborn</u></p> <p><u>Not to be used for underlying cause mortality coding. If no other perinatal cause is reported, code to Condition originating in the perinatal period, unspecified (P96.9).</u></p> <p><u>P72.2 – P74 Transitory endocrine and metabolic disorders specific to fetus and newborn</u></p> <p><u>Not to be used for underlying cause mortality coding. If no other perinatal cause is reported, code to Condition originating in the perinatal period, unspecified (P96.9).</u></p>	<p>MRG (URC:0120)</p>	<p>Minor</p>	<p>January 2004</p>
<p>Page 61 Delete text</p>	<p>T36 T50 Poisoning by drugs, medicaments and biological substances ———— (accidental poisoning and poisoning of undetermined intent by ———— alcohol or dependence producing drugs)</p> <p><i>with mention of:</i></p> <p>F10 F19 with fourth character .2 (alcohol dependence or drug dependence), code F10 F19 with fourth character .2</p>	<p>MRG (URC:0117)</p>	<p>Major</p>	<p>January 2006</p>
<p>Page 62 Delete text</p>	<p>X40 X49 Accidental poisoning by and exposure to noxious substances Y10 Y15 Poisoning by and exposure to noxious substances, undetermined intent ———— (poisoning by alcohol or dependence producing drugs)</p> <p><i>with mention of:</i></p> <p>F10 F19 with fourth character .2 (alcohol dependence or drug dependence), code F10 F19 with fourth character .2</p>	<p>MRG (URC:0117)</p>	<p>Major</p>	<p>January 2006</p>

<p>Page 62</p> <p>Add codes</p> <p>Page 65</p> <p>Delete codes</p>	<p>Table 1. Summary of linkages by code number</p> <table border="1"> <thead> <tr> <th>Selected cause:</th> <th>With mention of:</th> <th>Resulting linked code</th> </tr> </thead> <tbody> <tr> <td>E86</td> <td>A00-A09</td> <td>A00-A09</td> </tr> <tr> <td><u>F10-F19</u></td> <td><u>X40-X49</u></td> <td><u>X40-X49</u></td> </tr> <tr> <td><u>F10-F19</u></td> <td><u>X60-X69</u></td> <td><u>X60-X69</u></td> </tr> <tr> <td><u>F10-F19</u></td> <td><u>X85-X90</u></td> <td><u>X85-X90</u></td> </tr> <tr> <td><u>F10-F19</u></td> <td><u>Y10-Y19</u></td> <td><u>Y10-Y19</u></td> </tr> </tbody> </table> <table border="1"> <tbody> <tr> <td>T36 T50</td> <td>F10 F19 (F1x.2)</td> <td>F10 F19 (F1x.2)</td> </tr> <tr> <td>X40 X49 }</td> <td>F10 F19 (F1x.2)</td> <td>F10 F19 (F1x.2)</td> </tr> <tr> <td>Y10 Y15 }</td> <td>F10 F19 (F1x.2)</td> <td>F10 F19 (F1x.2)</td> </tr> </tbody> </table>	Selected cause:	With mention of:	Resulting linked code	E86	A00-A09	A00-A09	<u>F10-F19</u>	<u>X40-X49</u>	<u>X40-X49</u>	<u>F10-F19</u>	<u>X60-X69</u>	<u>X60-X69</u>	<u>F10-F19</u>	<u>X85-X90</u>	<u>X85-X90</u>	<u>F10-F19</u>	<u>Y10-Y19</u>	<u>Y10-Y19</u>	T36 T50	F10 F19 (F1x.2)	F10 F19 (F1x.2)	X40 X49 }	F10 F19 (F1x.2)	F10 F19 (F1x.2)	Y10 Y15 }	F10 F19 (F1x.2)	F10 F19 (F1x.2)	<p>MRG (URC:0117)</p>	<p>Major</p>	<p>January 2006</p>
Selected cause:	With mention of:	Resulting linked code																													
E86	A00-A09	A00-A09																													
<u>F10-F19</u>	<u>X40-X49</u>	<u>X40-X49</u>																													
<u>F10-F19</u>	<u>X60-X69</u>	<u>X60-X69</u>																													
<u>F10-F19</u>	<u>X85-X90</u>	<u>X85-X90</u>																													
<u>F10-F19</u>	<u>Y10-Y19</u>	<u>Y10-Y19</u>																													
T36 T50	F10 F19 (F1x.2)	F10 F19 (F1x.2)																													
X40 X49 }	F10 F19 (F1x.2)	F10 F19 (F1x.2)																													
Y10 Y15 }	F10 F19 (F1x.2)	F10 F19 (F1x.2)																													
<p>Page 66</p> <p>Add codes to existing table</p>	<p>Table 2. Summary of codes not to be used in underlying cause mortality coding</p> <p>Codes not to be used for underlying cause mortality coding (code to item in parentheses; if no code is indicated, code to R99)</p> <p><u>P70.3 – P72.0 (code to P96.9)</u></p> <p><u>P72.2 – P74 (code to P96.9)</u></p>	<p>MRG (URC:0120)</p>	<p>Minor</p>	<p>January 2004</p>																											
<p>Page 66</p> <p>Add list of codes to existing table</p>	<p>Table 2. Summary of codes not to be used in underlying cause mortality coding</p> <p>Codes not to be used for underlying cause mortality coding (code to item in parentheses; if no code is indicated, code to R99)</p> <p><u>F10.0 (code to X45, X65, X85, or Y15)</u></p> <p><u>F11.0 (code to X42, X62, X85, or Y12)</u></p> <p><u>F12.0 (code to X42, X62, X85, or Y12)</u></p> <p><u>F13.0 (code to X41, X61, X85, or Y11)</u></p> <p><u>F14.0 (code to X42, X62, X85, or Y12)</u></p> <p><u>F15.0 (code to X41, X61, X85, or Y11)</u></p> <p><u>F16.0 (code to X42, X62, X85, or Y12)</u></p> <p><u>F17.0 (code to X49, X69, X89, or Y19)</u></p> <p><u>F18.0 (code to X46, X66, X89, or Y16)</u></p> <p><u>F19.0 (code to X40-X49, X60-X69, X85-X90, or Y10-Y19)</u></p>	<p>MRG (URC:0116)</p>	<p>Major</p>	<p>January 2006</p>																											

<p>Page 67</p> <p>Delete text as indicated Add text as indicated Add text as indicated</p>	<p>4.2.2 Interpretation of “highly improbable” The expression “highly improbable” has been used since ... the following relationships should be regarded as “highly improbable”:</p> <p>(a) any infectious disease may be accepted as “due to” disorders of the immune mechanism such as human immunodeficiency virus (HIV) disease or AIDS; (b) an infectious or parasitic disease (A00-B99) reported as “due to” any disease outside this chapter, except that:</p> <ul style="list-style-type: none"> • diarrhoea and gastroenteritis of presumed infectious origin (A09)) • septicaemia (A40-A41)) • erysipelas (A46)) may be accepted • gas gangrene (A48.0)) as “due to” • Vincent’s angina (A69.1)) any other • mycoses (B35-B49)) disease • any infectious disease may be accepted as “due to” immunosuppression by chemicals (chemotherapy) and radiation. Any infectious disease classified to A00-B19 or B25-B64 reported as “due to” a malignant neoplasm will also be an acceptable sequence; • <u>any infectious disease classified to A00-B19 or B25-B64 reported as “due to” a malignant neoplasm will also be an acceptable sequence</u> • varicella and zoster infections (B01-B02) may be accepted as “due to” diabetes, tuberculosis and lymphoproliferative neoplasms; <p>(c) a malignant neoplasm reported as “due to” any other disease, except human immunodeficiency virus (HIV) disease;</p>	<p>MRG (URC: 0122)</p>	<p>Minor</p>	<p>January 2004</p>
<p>Page 68</p> <p>Revise text as shown</p>	<p>4.2.2 Interpretation of “highly improbable” ... the following relationships should be regarded as “highly improbable”:</p> <p>(l) a congenital anomaly (Q00-Q99) reported as “due to” any other disease of the individual, including immaturity; except for:</p> <ul style="list-style-type: none"> • <u>a congenital anomaly reported as “due to” a chromosome abnormality or a congenital malformation syndrome</u> • <u>pulmonary hypoplasia reported as “due to” a congenital anomaly</u> 	<p>MRG (URC:0118)</p>	<p>Major</p>	<p>January 2006</p>
<p>Page 68</p> <p>Delete existing text and</p>	<p>4.2.2 Interpretation of “highly improbable” ... the following relationships should be regarded as “highly improbable”:</p> <p>(i) any cerebrovascular disease (I60-I69) reported as “due to” a disease of the digestive system (K00-K92) or endocarditis (I05-I08, I09.1, I33-I38), except for</p>	<p>MRG (URC:0119)</p>	<p>Minor</p>	<p>January 2004</p>

<p>replace with the following text</p>	<p>cerebral embolism in I65 I66 or intracranial haemorrhage (I60-I62);</p> <p>(i)</p> <p><u>(1) cerebrovascular diseases (I60-I69) reported as “due to” a disease of the digestive system (K00-K92);</u></p> <p><u>(2) cerebral infarction due to thrombosis of precerebral arteries (I63.0)</u> <u>cerebral infarction due to unspecified occlusion of precerebral arteries (I63.2)</u> <u>cerebral infarction due to thrombosis of cerebral arteries (I63.3)</u> <u>cerebral infarction due to unspecified occlusion of cerebral arteries (I63.5)</u> <u>cerebral infarction due to cerebral venous thrombosis, nonpyogenic (I63.6)</u> <u>other cerebral infarction (I63.8)</u> <u>cerebral infarction, unspecified (I63.9)</u> <u>stroke, not specified as haemorrhage or infarction (I64)</u> <u>other cerebrovascular diseases (I67)</u> <u>sequelae of stroke, not specified as haemorrhage or infarction (I69.4)</u> <u>sequelae of other and unspecified cerebrovascular diseases (I69.8)</u></p> <p><u>reported as “due to” endocarditis (I05-I08, I09.1, I33-I38);</u></p> <p><u>(3) occlusion and stenosis of precerebral arteries, not resulting in cerebral infarction (I65), except embolism</u> <u>occlusion and stenosis of cerebral arteries, not resulting in cerebral infarction (I66), except embolism</u> <u>sequelae of cerebral infarction (I69.3), except embolism</u></p> <p><u>reported as “due to” endocarditis (I05-I08, I09.1, I33-I38);</u></p>			
<p>Page 86</p> <p>Add text</p>	<p>4.2.9 Congenital malformations, deformations and chromosomal abnormalities</p> <p>If the interval between onset...on the medical certificate.</p> <p><u>On neonatal or infant death certificates, where lung or pulmonary hypoplasia is given with any mention of immaturity, prematurity, short gestation or low birth weight, code to pulmonary immaturity (P28.0) and not to Q33.6.</u></p>	<p>MRG (URC:0118)</p>	<p>Major</p>	<p>January 2006</p>
<p>Paste these recommendations into Volume 2 after page 138 and before page 139.</p>	<p>Recommendations</p> <p>1. Responsibility for medical certification of cause of death (see section 5.2)</p> <p>The medical certification of the cause of death is normally the responsibility of the attending physician. In the case of deaths certified by coroners or other legal authorities, the medical evidence supplied to the certifier should be stated on the</p>	<p>MRG (URC:0113)</p>	<p>Major</p>	<p>January 2006</p>

	<p>certificate in addition to any legal findings.</p> <p>2. Form of medical certificate of cause of death (see sections 5.2, 4.1.3, and 4.3.1)</p> <p>The medical certificate of cause of death should be in line with the international recommendation (see section 4.1.3). Collection of perinatal mortality statistics should be consistent with the recommendations presented in section 4.3.1.</p> <p>3. Confidentiality of medical information (see section 5.2)</p> <p>Administrative procedures should ensure the confidentiality of data from the death certificate or other medical records.</p> <p>4. Selection of the cause for mortality tabulation (see section 4.1.1)</p> <p>The causes of death to be entered on the medical certificate of cause of death are all diseases, morbid conditions or injuries resulting in or contributing to death and the circumstances of the accident or violence resulting in injuries. When only one cause of death is recorded, this cause is selected for tabulation. When more than one cause of death is recorded, selection should be made in accordance with the rules and guidelines given in the ICD.</p> <p>5. Use of the International Classification of Diseases (see sections 2.1, 2.2, and 3.3)</p> <p>The purpose of the ICD is to permit the systematic recording, analysis, interpretation and comparison of mortality and morbidity data collected in different countries or areas and at different times. The “core” classification of ICD-10 is the three-character code, which is the mandatory level of coding for international reporting to the WHO mortality database and for general international comparisons. The four-character subcategories, while not mandatory for reporting at the international level, are recommended for many purposes and form an integral part of the ICD, as do the special tabulation lists.</p> <p>Mortality and morbidity statistics should be coded according to the tabular list of inclusions and the alphabetical index. Fourth-character subcategories, when published, should be those of the ICD. Any additions or variations should be indicated in published statistical tables.</p> <p>6. Perinatal mortality statistics (see sections 5.7.2 and 5.7.3)</p> <p>It is recommended that all fetuses and infants weighing at least 500 g at birth, whether alive or dead, should be included in <i>national</i> statistics. When information on birth weight is unavailable, the corresponding criteria for gestational age (22 completed weeks) or body length (25 cm crown-heel) should be used. The criteria for deciding whether an event has taken place within the perinatal period should be applied in the order: (1) birth weight, (2) gestational age, (3) crown-heel length. The inclusion of</p>			
--	--	--	--	--

	<p>fetuses and infants weighing between 500 g and 1000 g in national statistics is recommended both because of its inherent value and because it improves the coverage of reporting at 1000 g and over.</p> <p>In statistics for international comparison, inclusion of the extremely low-birth-weight group disrupts the validity of comparisons and is not recommended. Countries should also present statistics in which both the numerator and the denominator of all ratios and rates are restricted to fetuses and infants weighing 1000 g or more (weight-specific ratios and rates); where information on birth weight is not available, the corresponding gestational age (28 completed weeks) or body length (35 cm crown-heel) should be used.</p> <p>7. Maternal mortality statistics (see sections 5.8.2 and 5.8.3)</p> <p>Published maternal mortality rates should always specify the numerator, which can be given as: the number of recorded direct obstetric deaths, or the number of recorded obstetric deaths (direct plus indirect). For the purpose of international reporting of maternal mortality, only those maternal deaths occurring before the end of the 42-day reference period should be included in the calculation of the various ratios and rates, although the recording of later deaths is useful for national analytical purposes.</p> <p>8. Statistical tables (see sections 5.6.1 and 5.7.4)</p> <p>The degree of detail in cross-classification by cause, sex, age, and geographical area will depend both on the purpose and range of the statistics and on the practical limits to their tabulation. Standard ways of presenting statistics are described in sections 5.6.1 and 5.7.4 to promote international compatibility.</p> <p>9. Tabulation of causes of death (see sections 5.6.2 and 5.6.4)</p> <p>Statistics of causes of death for a defined area should be in accordance with recommendations in section 5.6.1. Deaths should preferably be classified by sex and age group as in recommendations in section 5.6.1. For statistics of perinatal mortality, full-scale multiple-cause analysis of all conditions reported will be of greatest benefit. Where such analysis is impracticable, analysis of the main disease or condition in the fetus or infant and of the main maternal condition affecting the fetus or infant with cross-tabulation of groups of these two conditions should be regarded as the minimum. Where it is necessary to select only one condition, the main disease or condition in the fetus or infant should be selected.</p>			
	<p>Appendix 7.1</p> <p>List of conditions unlikely to cause death</p>	<p>MRG (URC:0122)</p>	<p>Minor</p>	<p>January 2004</p>

Ratified by HoC/WHO at the HoC meeting in Brisbane, October 2002

	Code	Category or subcategory			
Revise code	H54.0 – <u>H54.7</u>	Blindness and low vision			
Delete codes and text	H59.0 – H59.9	Postprocedural disorders of eye and adnexa, not elsewhere classified			
Delete code And text	M72.5	Fasciitis, not elsewhere classified, except necrotizing fasciitis			
Revise code	N92.0 – <u>N92.6</u>	Excessive, frequent, and irregular menstruation			

Volume 3**ALPHABETIC INDEX**

Instruction	Alphabetic index entries	Source	Major/Minor update	Implementation date
Revise code Add modifier Add subterm and code	Anomaly, anomalous (congenital) (unspecified type) Q89.9 - hydatid of Morgagni - - female <u>Q50.5</u> - - male (<u>epididymal</u>) Q55.4 - - - testicular <u>Q55.2</u>	UK (URC:0017)	Major	January 2006
Add modifier and revise code	Aphasia (amnesic)...R47.0 - auditory (<u>developmental</u>) F80.2	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Add subterm and code Revise code	Appendage - epididymal (organ of Morgagni) <u>Q55.4</u> - testicular (organ of Morgagni) <u>Q55.2</u>	UK (URC:0017)	Major	January 2006
Revise code Add modifier Add subterm and code Revise code	Appendix, appendicular – see also condition - Morgagni - - female <u>Q50.5</u> - - male (<u>epididymal</u>) Q55.4 - - - testicular <u>Q55.2</u> - testis <u>Q55.2</u>	UK (URC:0017)	Major	January 2006
Revise code	Atresia, atretic - hymen <u>Q52.3</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

Revise code	<p>Calcification - kidney N28.8 - - tuberculous <u>B90.1† N29.1*</u></p>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Revise code	<p>Cerebrospasticity (congenital) <u>G80.1</u></p>	Australia (URC:0045)	Major	January 2006
Add text as indicated	<p>Condition - see Disease <u>Conditions arising in the perinatal period</u></p> <p><u>Note - Conditions arising in the perinatal period, even though death or morbidity occurs later, should, as far as possible, be coded to chapter XVI, which takes precedence over chapters containing codes for diseases by their anatomical site.</u></p> <p><u>These exclude:</u> <u>Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)</u> <u>Endocrine, nutritional and metabolic diseases (E00-E99)</u> <u>Injury, poisoning and certain other consequences of external causes (S00-T99)</u> <u>Neoplasms (C00-D48)</u> <u>Tetanus neonatorum (A33)</u></p> <p>- <u>ablatio, ablation</u> - - <u>placentae, affecting fetus or newborn P02.1</u> - <u>abnormal, abnormality, abnormalities</u> - - <u>amnion, amniotic fluid, affecting fetus or newborn P02.9</u> - - <u>anticoagulation, newborn (transient) P61.6</u> - - <u>cervix, maternal (acquired) (congenital), in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - - <u>chorion, affecting fetus or newborn P02.9</u> - - <u>coagulation, newborn, transient P61.6</u> - - <u>fetus, fetal</u> - - - <u>causing disproportion, affecting fetus or newborn P03.1</u> - - <u>forces of labor affecting fetus or newborn P03.6</u> - - <u>labor NEC, affecting fetus or newborn P03.6</u> - - <u>membranes (fetal)</u> - - - <u>affecting fetus or newborn P02.9</u> - - - <u>specified type NEC, affecting fetus or newborn P02.8</u> - - <u>organs or tissues of maternal pelvis, in pregnancy or childbirth</u></p>	MRG (URC: 0105)	Minor	January 2004

	<p> <u>- - - - affecting fetus or newborn P03.8</u> <u>- - - - causing obstructed labor</u> <u>- - - - affecting fetus or newborn P03.1</u> <u>- - parturition, affecting fetus or newborn P03.9</u> <u>- - presentation (fetus)</u> <u>- - - before labor, affecting fetus or newborn P01.7</u> <u>- - - causing obstructed labour, affecting fetus or newborn (any, except breech) P03.1</u> <u>- - - - breech P03.0</u> <u>- - pulmonary</u> <u>- - - function, newborn P28.8</u> <u>- - - ventilation, newborn P28.8</u> <u>- - umbilical cord, affecting fetus or newborn P02.6</u> <u>- - uterus, maternal, in pregnancy or childbirth</u> <u>- - - - affecting fetus or newborn P03.8</u> <u>- - - - causing obstructed labor</u> <u>- - - - - affecting fetus or newborn P03.1</u> <u>- - vagina, maternal (acquired) (congenital), in pregnancy or childbirth</u> <u>- - - causing obstructed labor</u> <u>- - - - affecting fetus or newborn P03.1</u> <u>- - vulva and perineum, maternal (acquired) (congenital), in pregnancy or childbirth</u> <u>- - - causing obstructed labor</u> <u>- - - - affecting fetus or newborn P03.1</u> <u>- ABO hemolytic disease (fetus or newborn) P55.1</u> <u>- aborter, habitual or recurrent NEC</u> <u>- - current abortion, affecting fetus or newborn P01.8</u> <u>- abortion (complete) (incomplete)</u> <u>- - fetus or newborn P96.4</u> <u>- - habitual or recurrent, with current abortion, fetus P01.8</u> <u>- - legal (induced), fetus P96.4</u> <u>- - medical, fetus P96.4</u> <u>- - spontaneous, fetus P01.8</u> <u>- - - threatened, affecting fetus or newborn P01.8</u> <u>- - therapeutic, fetus P96.4</u> <u>- - threatened (spontaneous), affecting fetus or newborn P01.8</u> <u>- abruptio placentae, affecting fetus or newborn P02.1</u> <u>- abscess (embolic) (infective) (metastatic) (multiple) (perforated) (pyogenic) (septic)</u> <u>- - breast (acute) (chronic) (nonpuerperal), newborn P39.0</u> <u>- - kidney, maternal, complicating pregnancy</u> <u>- - - affecting fetus or newborn P00.1</u> </p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>navel, newborn P38</u> - - <u>umbilicus, newborn P38</u> - <u>absorption</u> - - <u>chemical</u> - - - <u>through placenta (fetus or newborn) P04.8</u> - - - - <u>environmental substance P04.6</u> - - - - <u>nutritional substance P04.5</u> - - - - <u>obstetric anesthetic or analgesic drug P04.0</u> - - <u>drug NEC (fetus or newborn) – see also Conditions arising in the perinatal period, reaction, drug</u> - - - <u>through placenta P04.1</u> - - - - <u>addictive P04.4</u> - - - - <u>obstetric anesthetic or analgesic medication P04.0</u> - - <u>maternal medication NEC through placenta (fetus or newborn) P04.1</u> - <u>accident</u> - - <u>birth - see Conditions originating in the perinatal period, birth, injury</u> - - <u>during pregnancy, to mother, affecting fetus or newborn P00.5</u> - <u>acidosis (lactic) (respiratory)</u> - - <u>fetal - see Conditions originating in the perinatal period, distress, fetal</u> - - <u>intrauterine – see Conditions originating in the perinatal period, distress, fetal</u> - - <u>metabolic NEC</u> - - - <u>late, of newborn P74.0</u> - - - <u>newborn – see Conditions originating in the perinatal period, distress, fetal</u> - <u>acrocyanosis, newborn P28.2</u> - <u>addiction, maternal</u> - - <u>alcohol, alcoholic (ethyl) (methyl) (wood), complicating pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P04.3</u> - - <u>drug NEC, maternal, complicating pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P04.4</u> - - - <u>withdrawal symptoms in newborn P96.1</u> - <u>adhesions, adhesive (postinfective)</u> - - <u>amnion to fetus, affecting fetus or newborn P02.8</u> - <u>adiponecrosis neonatorum P83.8</u> - <u>aeration lung imperfect, newborn P28.1</u> - <u>albuminuria, albuminuric (acute) (chronic) (subacute)</u> - - <u>pre-eclamptic, affecting fetus or newborn P00.0</u> - <u>alcoholism (acute) (chronic), complicating pregnancy or childbirth</u> - - <u>affecting fetus or newborn P04.3</u> - <u>amnionitis, affecting fetus or newborn P02.7</u> - <u>amputation, any part of fetus, to facilitate delivery P03.8</u> - <u>anaeriosis of newborn P28.8</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - <u>anasarca, fetus or newborn P83.2</u> - <u>android pelvis, maternal</u> - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>anemia</u> - - <u>congenital P61.4</u> - - - <u>due to isoimmunization NEC P55.9</u> - - - <u>following fetal blood loss P61.3</u> - - due to - - - <u>fetal blood loss P61.3</u> - - - <u>prematurity P61.2</u> - - <u>erythroblastic, fetus or newborn (see also Conditions originating in the perinatal period, disease, hemolytic) P55.9</u> - - <u>fetus or newborn P61.4</u> - - - due to - - - - <u>ABO (antibodies) (isoimmunization) (maternal/fetal incompatibility) P55.1</u> - - - - <u>Rh (antibodies) (isoimmunization) (maternal/fetal incompatibility) P55.0</u> - - - <u>following fetal blood loss P61.3</u> - - - <u>posthemorrhagic P61.3</u> - - <u>hemolytic, acute, fetus or newborn (see also Conditions originating in the perinatal period, disease, hemolytic) P55.9</u> - - <u>maternal, of or complicating pregnancy</u> - - - <u>affecting fetus or newborn P00.8</u> - - <u>of prematurity P61.2</u> - - <u>posthemorrhagic (chronic), newborn P61.3</u> - <u>anomaly, anomalous (congenital) (unspecified type)</u> - - <u>cervix, maternal, in pregnancy or childbirth NEC</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - - <u>uterus, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - <u>anoxemia – see Conditions originating in the perinatal period, anoxia</u> - <u>anoxia (see also Conditions, originating in the perinatal period, hypoxia)</u> - - <u>cerebral, newborn (see also Conditions originating in the perinatal period, asphyxia, newborn) P21.9</u> - - <u>newborn (see also Conditions, originating in the perinatal period, asphyxia, newborn) P21.9</u> - <u>anteversion</u> - - <u>uterus, uterine, maternal (cervix) (postinfectional) (postpartal, old), in pregnancy</u> 			
--	--	--	--	--

	<p><u>or childbirth</u></p> <ul style="list-style-type: none"> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - <u>anthropoid pelvis, maternal</u> - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>antibodies (blood group) (see also Conditions originating in the perinatal period, incompatibility)</u> - - <u>anti-D, fetus or newborn P55.0</u> - <u>anuria, newborn P96.0</u> - <u>apgar (score)</u> - - <u>low NEC, with asphyxia P21.9</u> - - <u>0-3 at 1 minute, with asphyxia P21.0</u> - - <u>4-7 at 1 minute, with asphyxia P21.1</u> - <u>apnea, apneic (spells), newborn NEC P28.4</u> - - <u>sleep (primary) P28.3</u> - <u>arrest, arrested</u> - - <u>active phase of labor, affecting fetus or newborn P03.6</u> - - <u>cardiac, newborn P29.1</u> - - <u>coronary, infant P29.1</u> - - <u>deep transverse, affecting fetus or newborn P03.1</u> - - <u>development or growth, fetus P05.9</u> - - <u>respiratory, newborn P28.5</u> - <u>arrhythmia (cardiac) (ventricular), newborn P29.1</u> - <u>asphyxia, asphyxiation (by)</u> - - <u>antenatal (see also Conditions originating in the perinatal period, distress, fetal) P20.9</u> - - <u>birth (see also Conditions originating in the perinatal period, asphyxia, newborn) P21.9</u> - - <u>fetal (see also Conditions originating in the perinatal period, distress, fetal) P20.9</u> - - <u>food or foreign body (in), newborn P24.3</u> - - <u>intrauterine (see also Conditions originating in the perinatal period, distress, fetal) P20.9</u> - - <u>mucus (in), newborn P24.1</u> - - <u>newborn P21.9</u> - - - <u>with 1-minute Apgar score</u> - - - - <u>low NEC P21.9</u> - - - - <u>0-3 P21.0</u> - - - - <u>4-7 P21.1</u> - - - <u>blue P21.1</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - - <u>livida P21.1</u> - - - <u>mild or moderate P21.1</u> - - - <u>pallida P21.0</u> - - - <u>severe P21.0</u> - - - <u>white P21.0</u> - - <u>perinatal – see Conditions originating in the perinatal period, asphyxia, newborn</u> - - <u>postnatal – see Conditions originating in the perinatal period, asphyxia, newborn</u> - - <u>perinatal (see also Conditions originating in the perinatal period, distress, fetal)</u> <u>P20.9</u> - <u>aspiration</u> - - <u>amniotic fluid (newborn) P24.1</u> - - <u>blood, newborn P24.2</u> - - <u>liquor (amni) (newborn) P24.1</u> - - <u>meconium (newborn) P24.0</u> - - <u>milk (newborn) P24.3</u> - - <u>mucus, newborn P24.1</u> - - <u>newborn (massive) (syndrome) P24.9</u> - - - <u>meconium P24.0</u> - - <u>vernix caseosa (newborn) P24.8</u> - <u>asymmetry</u> - - <u>lumbar spine with disproportion, affecting fetus or newborn P03.1</u> - - <u>pelvis with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>atelectasis (massive) (partial) (pressure) (pulmonary)</u> - - <u>fetus or newborn (secondary) P28.1</u> - - - <u>due to resorption P28.1</u> - - - <u>partial P28.1</u> - - - <u>primary P28.0</u> - - - <u>subtotal P28.1</u> - <u>atonia, atony, atonic</u> - - <u>congenital P94.2</u> - - <u>uterus, maternal, during labor</u> - - - <u>affecting fetus or newborn P03.6</u> - <u>atresia, atretic</u> - - <u>cervix, maternal (acquired), in pregnancy or childbirth</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - <u>attack</u> - - <u>cyanotic, newborn P28.2</u> - - <u>respiration, respiratory, newborn P28.8</u> - <u>awareness of heart beat</u> - - <u>fetal P20.9</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>newborn P29.1</u> - <u>baby, floppy (syndrome) P94.2</u> - <u>bacteremia</u> - - <u>due to bacterial organisms NEC – see also Conditions originating in the perinatal period, infection, by specified organism</u> - - - <u>newborn P36.9</u> - <u>Bandl's ring (contraction), complicating delivery</u> - - <u>affecting fetus or newborn P03.6</u> - <u>bicornate or bicornis maternal uterus, in pregnancy or childbirth</u> - - <u>affecting fetus or newborn P03.8</u> - <u>bigeminal pulse</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>birth</u> - - <u>abnormal NEC, affecting fetus or newborn P03.9</u> - - <u>delayed, fetus P03.8</u> - - <u>difficult NEC, affecting fetus or newborn P03.9</u> - - <u>forced NEC, affecting fetus or newborn P03.8</u> - - <u>forceps, affecting fetus or newborn P03.2</u> - - <u>immature (between 28 and 37 completed weeks) P07.3</u> - - - <u>extremely (less than 28 completed weeks) P07.2</u> - - <u>induced, affecting fetus or newborn P03.8</u> - - <u>injury P15.9</u> - - - <u>basal ganglia P11.1</u> - - - <u>brachial plexus NEC P14.3</u> - - - <u>brain (compression) (pressure) P11.2</u> - - - <u>central nervous system NEC P11.9</u> - - - <u>cerebellum P11.1</u> - - - <u>cerebral hemorrhage P10.1</u> - - - <u>external genitalia P15.5</u> - - - <u>eye P15.3</u> - - - <u>face P15.4</u> - - - <u>fracture</u> - - - - <u>bone P13.9</u> - - - - <u>specified NEC P13.8</u> - - - - <u>clavicle P13.4</u> - - - - <u>femur P13.2</u> - - - - <u>humerus P13.3</u> - - - - <u>long bone, except femur P13.3</u> - - - - <u>radius and ulna P13.3</u> - - - - <u>skull P13.0</u> 			
--	--	--	--	--

	<p> <u>---- spine P11.5</u> <u>---- tibia and fibula P13.3</u> <u>--- intracranial P11.2</u> <u>---- laceration or hemorrhage P10.9</u> <u>----- specified NEC P10.8</u> <u>--- intraventricular hemorrhage P10.2</u> <u>--- laceration</u> <u>---- brain P10.1</u> <u>---- by scalpel P15.8</u> <u>---- peripheral nerve P14.9</u> <u>--- liver P15.0</u> <u>--- meninges</u> <u>---- brain P11.1</u> <u>---- spinal cord P11.5</u> <u>--- nerve</u> <u>---- brachial plexus P14.3</u> <u>---- cranial NEC (except facial) P11.4</u> <u>---- facial P11.3</u> <u>---- peripheral P14.9</u> <u>---- phrenic (paralysis) P14.2</u> <u>--- penis P15.5</u> <u>--- scalp P12.9</u> <u>--- scalpel wound P15.8</u> <u>--- scrotum P15.5</u> <u>--- skull NEC P13.1</u> <u>---- fracture P13.0</u> <u>--- specified site or type NEC P15.8</u> <u>--- spinal cord P11.5</u> <u>--- spine P11.5</u> <u>--- spleen P15.1</u> <u>--- sternomastoid (hematoma) P15.2</u> <u>--- subarachnoid hemorrhage P10.3</u> <u>--- subcutaneous fat necrosis P15.6</u> <u>--- subdural hemorrhage P10.0</u> <u>--- tentorial tear P10.4</u> <u>--- testes P15.5</u> <u>--- vulva P15.5</u> <u>-- instrumental NEC, affecting fetus or newborn P03.8</u> <u>-- multiple, affecting fetus or newborn P01.5</u> <u>-- palsy or paralysis, newborn, NEC (birth injury) P14.9</u> <u>-- post-term (42 weeks or more) P08.2</u> </p>			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>precipitate, affecting fetus or newborn P03.5</u> - - <u>premature (infant) P07.3</u> - - <u>prolonged, affecting fetus or newborn P03.8</u> - - <u>retarded, affecting fetus or newborn P03.8</u> - - <u>shock, newborn P96.8</u> - - <u>trauma - see Conditions originating in perinatal period, birth, injury</u> - - <u>twin, affecting fetus or newborn P01.5</u> - - <u>vacuum extractor, affecting fetus or newborn P03.3</u> - - <u>ventouse, affecting fetus or newborn P03.3</u> - - <u>weight</u> - - - <u>low (between 1000 and 2499 grams at birth) P07.1</u> - - - - <u>extremely (999 grams or less at birth) P07.0</u> - - - <u>4500 grams or more P08.0</u> - <u>bleb(s) lung (ruptured), fetus or newborn P25.8</u> - <u>bleeding (see also Conditions originating in perinatal period, hemorrhage)</u> - - <u>rectum, rectal, newborn P54.2</u> - - <u>umbilical stump P51.9</u> - - <u>vagina, vaginal (abnormal), newborn P54.6</u> - <u>blood dyscrasia, fetus or newborn P61.9</u> - <u>born in toilet (see also Birth, precipitate fetus or newborn) P03.5</u> - <u>brachycardia</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>bradycardia (any type) (sinoatrial) (sinus) (vagal)</u> - - <u>fetal - see Conditions originating in the perinatal period, distress, fetal</u> - - <u>newborn P29.1</u> - <u>breech</u> - - <u>delivery NEC, affecting fetus or newborn P03.0</u> - - <u>extraction, affecting fetus or newborn P03.0</u> - - <u>presentation</u> - - - <u>before labor, affecting fetus or newborn P01.7</u> - - - <u>during labor, affecting fetus or newborn P03.0</u> - <u>bronze baby syndrome P83.8</u> - <u>bruise (skin surface intact)</u> - - <u>fetus or newborn P54.5</u> - - <u>scalp, due to birth injury, newborn P12.3</u> - - <u>umbilical cord, affecting fetus or newborn P02.6</u> - <u>bubbly lung syndrome P27.0</u> - <u>bullae, lung, fetus or newborn P25.8</u> - <u>candidiasis, candidal</u> - - <u>congenital P37.5</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>neonatal P37.5</u> - <u>caput succedaneum P12.8</u> - <u>catastrophe, catastrophe, cardiorespiratory, newborn P28.8</u> - <u>caul over face (causing asphyxia) P21.9</u> - <u>cellulitis (diffuse) (with lymphangitis)</u> - - <u>navel, newborn P38</u> - - <u>umbilicus, newborn P38</u> - <u>cephalohematocele, cephal(o)hematocele</u> - - <u>fetus or newborn P52.8</u> - - - <u>birth injury P10.8</u> - <u>cephalohematoma, cephalohematoma (calcified)</u> - - <u>fetus or newborn (birth injury) P12.0</u> - <u>cervicitis, maternal (acute) (chronic) (nonvenereal) (subacute) (with ulceration)</u> - - <u>complicating pregnancy, affecting fetus or newborn P00.8</u> - <u>cesarean operation or section</u> - - - <u>affecting fetus or newborn P03.4</u> - - - <u>post mortem, affecting fetus or newborn P01.6</u> - <u>cessation</u> - - <u>cardiac, newborn P29.0</u> - - <u>cardiorespiratory, newborn P29.0</u> - - <u>respiratory, newborn P28.5</u> - <u>chemotherapy (session) (for)</u> - - <u>cancer, maternal, affecting fetus or newborn P04.1</u> - <u>chickenpox, congenital P35.8</u> - <u>chignon, fetus or newborn (birth injury) (from vacuum extraction) P12.1</u> - <u>chorioamnionitis, fetus or newborn P02.7</u> - <u>chorioretinitis, in toxoplasmosis, congenital (active) P37.1† H32.0*</u> - <u>circulation</u> - - <u>failure (peripheral)</u> - - - <u>fetus or newborn P29.8</u> - - <u>fetal, persistent P29.3</u> - <u>cirrhosis, cirrhotic (hepatic)</u> - - <u>liver (chronic) (hepatolienal) (hypertrophic) (nodular) (splenomegalic)</u> - - - <u>congenital P78.8</u> - <u>cleidotomy, fetus or newborn P03.8</u> - <u>clotting, disseminated, intravascular, newborn P60</u> - <u>coagulation, intravascular (diffuse) (disseminated)</u> - - <u>antepartum, affecting fetus or newborn P02.1</u> - - <u>fetus or newborn P60</u> - <u>coagulopathy (see also Conditions arising in the perinatal period, defect, coagulation)</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>consumption, newborn P60</u> - <u>cold injury syndrome (newborn) P80.0</u> - <u>collapse</u> - - <u>cardiocirculatory, newborn P29.8</u> - - <u>cardiopulmonary, newborn P29.8</u> - - <u>cardiovascular, newborn P29.8</u> - - <u>circulatory (peripheral), fetus or newborn P29.8</u> - - <u>respiratory, newborn P28.8</u> - - <u>vascular (peripheral)</u> - - - <u>during labor and delivery, affecting fetus or newborn P03.8</u> - - - <u>fetus or newborn P29.8</u> - <u>coma, newborn P91.5</u> - <u>complications (from) (of)</u> - - <u>intrauterine procedure NEC, affecting fetus or newborn P96.5</u> - - <u>maternal sedation during labor and delivery, affecting fetus or newborn P04.0</u> - - <u>umbilical cord, affecting fetus or newborn P02.6</u> - <u>compression</u> - - <u>during birth (fetus or newborn) P15.9</u> - - <u>umbilical cord, affecting fetus or newborn P02.5</u> - - - <u>with cord prolapse P02.4</u> - <u>compromise, respiratory, newborn P28.5</u> - <u>congestion, congestive (chronic) (passive)</u> - - <u>facial, due to birth injury P15.4</u> - <u>conjunctivitis (in) (due to)</u> - - <u>chlamydial, neonatal P39.1</u> - - <u>neonatal P39.1</u> - <u>constriction</u> - - <u>cervix, cervical (canal), in pregnancy or childbirth</u> - - - <u>causing obstructed labour, affecting fetus or newborn P03.1</u> - - <u>ring dystocia (uterus), affecting fetus or newborn P03.6</u> - <u>contraction, contracture, contracted</u> - - <u>hourglass uterus (complicating delivery), affecting fetus or newborn P03.6</u> - - <u>pelvis, maternal (acquired) (general)</u> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - - <u>ring (Bandl's) (complicating delivery), affecting fetus or newborn P03.6</u> - - <u>uterus, maternal, abnormal NEC, affecting fetus or newborn P03.6</u> - <u>contusion (skin surface intact)</u> - - <u>fetus or newborn P54.5</u> - - <u>scalp, due to birth injury P12.3</u> - <u>convulsions (idiopathic), newborn P90</u> - <u>cord around neck (tightly) (with compression), affecting fetus or newborn P02.5</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - <u>coupled rhythm</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>cranioclasia, fetus P03.8</u> - <u>craniotabes (cause unknown), neonatal P96.3</u> - <u>craniotomy, fetus P03.8</u> - <u>cretin, cretinism (congenital) (endemic) (nongoitrous) (sporadic)</u> - - <u>pelvis, maternal, with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>cyst (colloid) (mucous) (retention) (simple)</u> - - <u>periventricular, acquired, newborn P91.1</u> - <u>cystitis, maternal (exudative) (hemorrhagic) (septic) (suppurative)</u> - - <u>complicating pregnancy, affecting fetus or newborn P00.1</u> - <u>cystocele(-rectocele), maternal, in pregnancy or childbirth</u> - - <u>affecting fetus or newborn P03.8</u> - - <u>causing obstructed labor</u> - - - <u>affecting fetus or newborn P03.1</u> - <u>dacryocystitis (acute) (phlegmonous), neonatal P39.1</u> - <u>damage</u> - - <u>brain (nontraumatic)</u> - - - <u>anoxic, hypoxic</u> - - - - <u>at birth P21.9</u> - - - - <u>fetal P20.9</u> - - - - <u>intrauterine P20.9</u> - - - - <u>newborn P21.9</u> - - - <u>due to birth injury P11.2</u> - - - <u>ischemic, newborn P91.0</u> - - - <u>newborn P11.2</u> - - <u>eye, birth injury P15.3</u> - <u>deadborn fetus P95</u> - <u>death</u> - - <u>fetus, fetal (cause not stated) (intrauterine) P95</u> - - <u>infant, from intrauterine coil P00.7</u> - - <u>neonatal NEC P96.8</u> - - <u>obstetric, maternal (cause unknown), affecting fetus or newborn P01.6</u> - <u>debility (chronic) (general), congenital or neonatal NEC P96.9</u> - <u>decapitation, fetal (to facilitate delivery) P03.8</u> - <u>deciduitis (acute), affecting fetus or newborn P00.8</u> - <u>decompensation, lung (pulmonary), newborn P28.8</u> - <u>defect, defective coagulation (factor)</u> - - <u>antepartum with hemorrhage, maternal, affecting fetus or newborn P02.1</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>newborn, transient P61.6</u> - <u>defibrination (syndrome)</u> - - <u>antepartum, maternal, affecting fetus or newborn P02.1</u> - - <u>fetus or newborn P60</u> - <u>deficiency, deficient</u> - - <u>coagulation</u> - - - <u>antepartum, maternal, affecting fetus or newborn P02.1</u> - - - <u>newborn, transient P61.6</u> - - <u>surfactant P28.0</u> - - <u>vitamin K, of newborn P53</u> - <u>deformity</u> - - <u>fetal, causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>pelvis, pelvic, maternal (acquired) (bony)</u> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - - <u>soft parts, maternal organs or tissues (of pelvis), in pregnancy or childbirth NEC</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor</u> - - - - <u>affecting fetus or newborn P03.1</u> - <u>dehydration, newborn P74.1</u> - <u>delay, delayed,</u> - - <u>birth or delivery NEC, affecting fetus or newborn P03.8</u> - - <u>closure, ductus arteriosus (Botalli) P29.3</u> - - <u>delivery, second twin, triplet, etc.</u> - - - <u>affecting fetus or newborn P03.8</u> - - <u>primary respiration (<i>see also</i> Conditions originating in the perinatal period, asphyxia, newborn) P21.9</u> - <u>delivery (single)</u> - - <u>breech NEC, affecting fetus or newborn P03.0</u> - - <u>cesarean (for), affecting fetus or newborn P03.4</u> - - <u>extremely rapid, newborn P03.5</u> - - <u>forceps, affecting fetus or newborn P03.2</u> - - <u>precipitate, affecting fetus or newborn P03.5</u> - - <u>premature or preterm NEC, affecting fetus or newborn P07.3</u> - - <u>vacuum extractor NEC, affecting fetus or newborn P03.3</u> - - <u>ventouse NEC, affecting fetus or newborn P03.3</u> - <u>demise, fetal P95</u> - <u>dependence</u> - - <u>due to drug NEC</u> - - - <u>maternal, complicating pregnancy or childbirth</u> - - - - <u>affecting fetus or newborn P04.4</u> - - - - <u>withdrawal symptoms in newborn P96.1</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - <u>depression</u> - - <u>central nervous system (CNS) NEC, newborn P91.4</u> - - <u>cerebral, newborn P91.4</u> - - <u>respiration, respiratory, newborn P28.8</u> - - <u>vital centers, newborn P91.4</u> - <u>destruction, live fetus to facilitate delivery (fetus) P03.8</u> - <u>development</u> - - <u>arrested, fetus P05.9</u> - - <u>incomplete P05.9</u> - <u>device, contraceptive, intrauterine, affecting fetus or newborn P00.8</u> - <u>diabetes, diabetic (controlled) (familial) (mellitus) (on insulin) (severe) (uncontrolled)</u> - - <u>arising in pregnancy, maternal, affecting fetus or newborn P70.0</u> - - <u>complicating pregnancy or childbirth, maternal, affecting fetus or newborn P70.1</u> - - - <u>arising in pregnancy, affecting fetus or newborn P70.0</u> - - - <u>gestational, affecting fetus or newborn P70.0</u> - - <u>neonatal (transient) P70.2</u> - <u>diarrhea, diarrheal (disease) (endemic) (infantile) (summer) (see also Conditions originating in the perinatal period, enteritis)</u> - - <u>neonatal (noninfective) P78.3</u> - <u>difficult, difficulty (in)</u> - - <u>birth, affecting fetus or newborn P03.9</u> - - <u>feeding, newborn P92.9</u> - - - <u>breast P92.5</u> - - - <u>specified NEC P92.8</u> - - <u>respiratory, newborn P28.8</u> - <u>dilatation cervix (uteri), maternal - see also Conditions originating in the perinatal period, incompetency, cervix</u> - - <u>incomplete, poor, slow, affecting fetus or newborn P03.6</u> - <u>disease, diseased - see also Conditions originating in the perinatal period, syndrome</u> - - <u>breast, inflammatory, fetus or newborn P83.4</u> - - <u>cardiorespiratory, newborn P96.8</u> - - <u>cardiovascular, fetus or newborn P29.9</u> - - - <u>specified NEC P29.8</u> - - <u>circulatory (system) NEC, fetus or newborn P29.9</u> - - - <u>maternal, affecting fetus or newborn P00.3</u> - - <u>facial nerve (seventh), newborn (birth injury) P11.3</u> - - <u>heart (organic)</u> - - - <u>congenital</u> 			
--	---	--	--	--

	<p> <u>--- maternal, affecting fetus or newborn P00.3</u> <u>--- rheumatic (chronic) (inactive) (old) (quiescent) (with chorea)</u> <u>--- maternal, affecting fetus or newborn P00.3</u> <u>-- hemolytic (fetus) (newborn) P55.9</u> <u>--- due to</u> <u>--- incompatibility</u> <u>----- ABO (blood group) P55.1</u> <u>----- blood (group) (Duffy) (K(ell)) (Kidd) (Lewis) (M) (S) NEC P55.8</u> <u>----- Rh (blood group) (factor) P55.0</u> <u>----- Rh-negative mother P55.0</u> <u>--- specified type NEC P55.8</u> <u>-- hemorrhagic, fetus or newborn P53</u> <u>-- hyaline (diffuse) (generalized), membrane (lung) (newborn) P22.0</u> <u>-- infectious, infective</u> <u>--- congenital P37.9</u> <u>----- specified NEC P37.8</u> <u>--- maternal, complicating pregnancy or childbirth</u> <u>---- affecting fetus or newborn P00.2</u> <u>-- maternal, unrelated to pregnancy NEC, affecting fetus or newborn P00.9</u> <u>-- pelvis, pelvic, maternal</u> <u>-- inflammatory (female), complicating pregnancy</u> <u>--- affecting fetus or newborn P00.8</u> <u>-- placenta, affecting fetus or newborn P02.2</u> <u>-- renal, maternal (functional) (pelvis)</u> <u>--- complicating pregnancy, affecting fetus or newborn P00.1</u> <u>-- respiratory (tract)</u> <u>--- chronic NEC, fetus or newborn P27.9</u> <u>---- specified NEC P27.8</u> <u>--- newborn P28.9</u> <u>---- specified type NEC P28.8</u> <u>-- viral, virus NEC</u> <u>--- congenital P35.9</u> <u>---- specified NEC P35.8</u> <u>- disorder (of) – see also Conditions originating in the perinatal period, disease</u> <u>-- amino-acid, neonatal, transitory P74.8</u> <u>-- coagulation (factor)</u> <u>--- antepartum with hemorrhage, maternal, affecting fetus or newborn P02.1</u> <u>--- newborn, transient P61.6</u> <u>-- digestive (system), fetus or newborn P78.9</u> <u>--- specified NEC P78.8</u> <u>-- feeding, newborn P92.9</u> </p>			
--	---	--	--	--

	<ul style="list-style-type: none"> - - fetus or newborn P96.9 - - - specified NEC P96.8 - - hematological, fetus or newborn P61.9 - - - specified NEC P61.8 - - hemorrhagic NEC, newborn P53 - - integument, fetus or newborn P83.9 - - - specified NEC P83.8 - - membranes or fluid, amniotic, affecting fetus or newborn P02.9 - - muscle tone, newborn P94.9 - - - specified NEC P94.8 - - seizure, newborn P90 - - skin, fetus or newborn P83.9 - - - specified NEC P83.8 - - temperature regulation, fetus or newborn P81.9 - - - specified NEC P81.8 - - thyroid (gland) function NEC, neonatal, transitory P72.2 - disproportion (fetopelvic), affecting fetus or newborn P03.1 - distortion (congenital) lumbar spine, maternal - - with disproportion, affecting fetus or newborn P03.1 - distress - - cardiac - - - congenital P20.9 - - - newborn P29.8 - - cardiopulmonary, newborn P96.8 - - cardiorespiratory, newborn P96.8 - - circulatory, newborn P96.8 - - fetal (syndrome) P20.- - - - first noted - - - - before onset of labor P20.0 - - - - during labor and delivery P20.1 - - intrauterine - <i>see</i> Conditions originating in the perinatal period, distress, fetal - - respiratory, newborn P22.9 - - - specified NEC P22.8 - disturbance - <i>see also</i> Conditions originating in the perinatal period, disease - - cerebral status, newborn P91.9 - - - specified NEC P91.8 - - electrolyte - - - newborn, transitory P74.4 - - - - potassium balance P74.3 - - - - sodium balance P74.2 - - - - specified type NEC P74.4 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>endocrine (gland), neonatal, transitory P72.9</u> - - - <u>specified NEC P72.8</u> - - <u>feeding, newborn P92.9</u> - - <u>metabolism, neonatal, transitory P74.9</u> - - - <u>calcium and magnesium P71.9</u> - - - - <u>specified type NEC P71.8</u> - - - <u>carbohydrate metabolism P70.9</u> - - - - <u>specified type NEC P70.8</u> - - - <u>specified NEC P74.8</u> - - <u>potassium balance, newborn P74.3</u> - - <u>sodium balance, newborn P74.2</u> - - <u>temperature regulation, newborn P81.9</u> - - - <u>specified NEC P81.8</u> - <u>double uterus, maternal, in pregnancy or childbirth</u> - - <u>affecting fetus or newborn P03.8</u> - <u>Duchenne's paralysis, birth injury P14.0</u> - <u>dyscrasia, blood, fetus or newborn P61.9</u> - - <u>specified type NEC P61.8</u> - <u>dysfunction uterus, complicating delivery</u> - - <u>affecting fetus or newborn P03.6</u> - <u>dysmaturity (see also Conditions originating in the perinatal period, immaturity) P05.0</u> - - <u>pulmonary (newborn) (Wilson-Mikity) P27.0</u> - <u>dysplasia - see also Conditions originating in the perinatal period, anomaly</u> - - <u>bronchopulmonary (perinatal) P27.1</u> - - <u>lung (congenital), associated with short gestation P28.0</u> - <u>dyspnea (nocturnal) (paroxysmal), newborn P22.8</u> - <u>dysrhythmia (cardiac), newborn P29.1</u> - <u>dystocia</u> - - <u>affecting fetus or newborn P03.1</u> - - <u>cervical, maternal (hypotonic), affecting fetus or newborn P03.6</u> - - <u>contraction ring, maternal, affecting fetus or newborn P03.6</u> - - <u>fetal, fetus, affecting fetus or newborn P03.1</u> - - <u>maternal, affecting fetus or newborn P03.1</u> - - <u>positional, affecting fetus or newborn P03.1</u> - - <u>shoulder (girdle), affecting fetus or newborn P03.1</u> - - <u>uterine NEC, maternal, affecting fetus or newborn P03.6</u> - <u>ecchymosis (see also Conditions originating in the perinatal period, hemorrhage)</u> - - <u>fetus or newborn P54.5</u> - <u>eclampsia, eclamptic (coma) (convulsions) (delirium) (with pre-existing or pregnancy-related hypertension) NEC</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>pregnancy, affecting fetus or newborn P00.0</u> - <u>edema, edematous brain</u> - - <u>due to birth injury P11.0</u> - - <u>fetus or newborn (anoxia or hypoxia) P52.4</u> - - - <u>birth injury P11.0</u> - - <u>fetus or newborn NEC, classifiable to R60.- P83.3</u> - - <u>newborn, classifiable to R60.- P83.3</u> - <u>effect, adverse NEC</u> - - <u>anesthesia in labor and delivery, affecting fetus or newborn P04.0</u> - <u>effusion, pleura, pleurisy, pleuritic, pleuropericardial</u> - - <u>fetus or newborn P28.8</u> - <u>embarrassment, respiratory, newborn (see Conditions originating in the perinatal period, distress, respiratory)</u> - <u>embolism (septic), air (any site) (traumatic), newborn NEC P25.8</u> - <u>embryotomy (to facilitate delivery), fetus P03.8</u> - <u>emphysema (atrophic) (chronic) (interlobular) (lung) (obstructive) (pulmonary) (senile)</u> - - <u>congenital (interstitial) P25.0</u> - - <u>fetus or newborn (interstitial) P25.0</u> - - <u>interstitial, congenital P25.0</u> - - - <u>perinatal period P25.0</u> - - <u>mediastinal, fetus or newborn P25.2</u> - <u>encephalitis in toxoplasmosis, congenital P37.1† G05.2*</u> - <u>encephalopathy hyperbilirubinemia, newborn P57.9</u> - <u>due to isoimmunization (conditions in P55.-) P57.0</u> - <u>encephalopathy (acute) (cerebral)</u> - - <u>hyperbilirubinemic, newborn P57.9</u> - - - <u>due to isoimmunization (conditions in P55.-) P57.0</u> - - <u>in birth injury P11.1</u> - - <u>ischemic</u> - - - <u>anoxic (hypoxic), newborn P21.9</u> - - - <u>newborn P91.0</u> - <u>endometritis (nonspecific) (purulent) (septic) (suppurative)</u> - - <u>maternal, complicating pregnancy, affecting fetus or newborn P00.8</u> - <u>engorgement, breast, newborn P83.4</u> - <u>entanglement umbilical cord(s), affecting fetus or newborn P02.5</u> - <u>enteritis (acute) (diarrheal) (hemorrhagic) (see also Conditions originating in the perinatal period, diarrhea)</u> - - <u>necrotizing of fetus or newborn P77</u> - <u>enterocolitis (see also Conditions originating in the perinatal period, diarrhea)</u> - - <u>necrotizing (chronic), fetus or newborn P77</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - <u>Erb(-Duchenne) paralysis (birth injury) (newborn) P14.0</u> - <u>Erb's palsy, paralysis (brachial) (birth) (newborn) P14.0</u> - <u>erythema, erythematous</u> - - <u>neonatorum P83.8</u> - - - <u>toxic P83.1</u> - - <u>rash, newborn P83.8</u> - - <u>toxic, toxicum NEC, newborn P83.1</u> - <u>erythroblastosis (fetalis) (newborn) P55.9</u> - - <u>due to</u> - - - <u>ABO (antibodies) (incompatibility) (isoimmunization) P55.1</u> - - - <u>Rh (antibodies) (incompatibility) (isoimmunization) P55.0</u> - <u>erythroderma neonatorum P83.8</u> - - <u>toxic P83.1</u> - <u>evisceration, birth injury P15.8</u> - <u>excess, excessive, excessively</u> - - <u>large, fetus or infant P08.0</u> - - <u>long umbilical cord (entangled), affecting fetus or newborn P02.5</u> - - <u>short umbilical cord, affecting fetus or newborn P02.6</u> - <u>exhaustion (physical NEC)</u> - - <u>fetus or newborn P96.8</u> - - <u>maternal, complicating delivery, affecting fetus or newborn P03.8</u> - <u>exsanguination – see Conditions originating in the perinatal period, hemorrhage</u> - <u>extraction</u> - - <u>with hook (fetus) P03.8</u> - - <u>breech NEC, affecting fetus or newborn P03.0</u> - <u>face, facial presentation, affecting fetus or newborn P01.7</u> - <u>failure, failed</u> - - <u>cardiac, newborn P29.0</u> - - <u>cardiopulmonary (acute) (chronic), newborn P29.0</u> - - <u>cervical dilatation in labor, affecting fetus or newborn P03.6</u> - - <u>descent of head (at term), affecting fetus or newborn P03.1</u> - - <u>expansion, terminal respiratory units (newborn) (primary) P28.0</u> - - <u>fetal head to enter pelvic brim, affecting fetus or newborn P03.1</u> - - <u>forceps NEC (with subsequent delivery by cesarean section)</u> - - - <u>affecting fetus or newborn P03.1</u> - - <u>heart (acute) (sudden), newborn P29.0</u> - - - <u>congestive P29.0</u> - - - <u>left (ventricular) P29.0</u> - - <u>pulmonary, newborn P28.5</u> - - <u>renal, congenital P96.0</u> - - <u>respiration, respiratory, newborn P28.5</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>trial of labor (with subsequent cesarean section), affecting fetus or newborn P03.1</u> - - <u>vacuum extraction NEC (with subsequent cesarean section), affecting fetus or newborn P03.1</u> - - <u>ventilatory, newborn P28.5</u> - - <u>ventouse NEC (with subsequent cesarean section), affecting fetus or newborn P03.1</u> - - <u>ventricular, left, newborn P29.0</u> - - <u>vital centers, fetus or newborn P91.8</u> - <u>fecalith (impaction), congenital P76.8</u> - <u>feeding problem, newborn P92.9</u> - - <u>specified NEC P92.8</u> - <u>fever, newborn P81.9</u> - - <u>environmental P81.0</u> - <u>fibroid (tumor), maternal</u> - - <u>in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - <u>fibromyoma, maternal</u> - - <u>uterus (corpus), in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - <u>fibrosis, fibrotic</u> - - <u>lung (atrophic) (capillary) (chronic) (confluent) (massive) (perialveolar) (peribronchial), congenital P27.8</u> - - <u>perineum, maternal, in pregnancy or childbirth</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>pulmonary, congenital P27.8</u> - <u>fit, newborn P90</u> - <u>flail chest, newborn (birth injury) P13.8</u> - <u>flat - see also Conditions arising in the perinatal period, anomaly, by site</u> - - <u>pelvis, maternal</u> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>floppy baby syndrome (nonspecific) P94.2</u> - <u>forced birth or delivery NEC, affecting fetus or newborn P03.8</u> - <u>forceps delivery NEC, affecting fetus or newborn P03.2</u> - <u>fracture (abduction) (adduction) (avulsion) (comminuted) (compression) (dislocation) (oblique) (separation)</u> - - <u>bone NEC, birth injury P13.9</u> - - <u>clavicle (acromial end) (interligamentous) (shaft), birth injury P13.4</u> - - <u>femur, femoral, birth injury P13.2</u> - - <u>skull, birth injury P13.0</u> - - <u>vertebra, vertebral (back) (body) (column) (neural arch) (pedicle) (spinous process) (transverse process), birth injury P11.5</u> - <u>funnel pelvis, maternal (acquired)</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>gallop rhythm</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>ganglionitis geniculate, newborn (birth injury) P11.3</u> - <u>gestation (period) - see also Conditions originating in the perinatal period, pregnancy</u> - - <u>less than 28 weeks P07.2</u> - - <u>28 weeks but less than 37 weeks P07.3</u> - - <u>42 or more completed weeks P08.2</u> - <u>glaucoma, traumatic, newborn (birth injury) P15.3</u> - <u>goiter (plunging) (substernal)</u> - - <u>congenital (nontoxic), transitory, with normal functioning P72.0</u> - - <u>neonatal NEC P72.0</u> - <u>gonococcus, gonococcal (disease) (infection) (see also condition)</u> - - <u>maternal, complicating pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P00.2</u> - <u>gonorrhea (acute) (chronic)</u> - - <u>maternal, complicating pregnancy or childbirth</u> - - <u>affecting fetus or newborn P00.2</u> - <u>granuloma, umbilicus, newborn P38</u> - <u>Gray syndrome (newborn) P93</u> - <u>Grey syndrome (newborn) P93</u> - <u>heart beat</u> - - <u>abnormality</u> - - - <u>fetal P20.9</u> - - - <u>newborn P29.1</u> - - <u>awareness</u> - - - <u>fetal P20.9</u> - - - <u>newborn P29.1</u> - <u>heavy-for-dates NEC (fetus or infant) P08.1</u> - - <u>exceptionally (4500 g or more) P08.0</u> - <u>hematemesis, newborn, neonatal P54.0</u> - - <u>due to swallowed maternal blood P78.2</u> - <u>hematoma (traumatic) (skin surface intact)</u> - - <u>birth injury NEC P15.8</u> - - <u>brain (traumatic), fetus or newborn NEC P52.4</u> - - - <u>birth injury P10.1</u> - - <u>face, birth injury P15.4</u> - - <u>liver (subcapsular), birth injury P15.0</u> - - <u>penis, birth injury P15.5</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>scrotum, superficial, birth injury P15.5</u> - - <u>spinal (cord) (meninges)</u> - - - <u>fetus or newborn (birth injury) P11.5</u> - - <u>sternocleidomastoid, birth injury P15.2</u> - - <u>sternomastoid, birth injury P15.2</u> - - <u>subarachnoid, fetus or newborn (nontraumatic) P52.5</u> - - - <u>birth injury P10.3</u> - - <u>subdural, fetus or newborn (localized) P52.8</u> - - - <u>birth injury P10.0</u> - - <u>superficial, fetus or newborn P54.5</u> - - <u>testis, birth injury P15.5</u> - - <u>umbilical cord, complicating delivery, affecting fetus or newborn P02.6</u> - - <u>vulva, fetus or newborn (birth injury) P15.5</u> - <u>hematomyelia (central), fetus or newborn (birth injury) P11.5</u> - <u>hematorachis, hematorrhachis, fetus or newborn (birth injury) P11.5</u> - <u>hemiplegia, newborn NEC P91.8</u> - - <u>birth injury P11.9</u> - <u>hemolysis</u> - - <u>autoimmune, newborn P55.9</u> - - <u>intravascular NEC, newborn P60</u> - - <u>neonatal (excessive) P58.8</u> - <u>hemopericardium, newborn P54.8</u> - <u>hemoptysis, newborn P26.9</u> - <u>hemorrhage, hemorrhagic</u> - - <u>accidental antepartum, affecting fetus or newborn P02.1</u> - - <u>adrenal (capsule) (gland), newborn P54.4</u> - - <u>alveolar, lung, newborn P26.8</u> - - <u>antepartum (see also Conditions originating in the perinatal period, hemorrhage, pregnancy)</u> - - - <u>affecting fetus or newborn P02.1</u> - - <u>bowel, newborn P54.3</u> - - <u>brain (miliary) (nontraumatic) (petechial)</u> - - - <u>due to birth injury P10.1</u> - - - <u>fetus or newborn P52.4</u> - - - - <u>birth injury P10.1</u> - - - <u>stem, newborn P52.4</u> - - <u>cerebellar, cerebellum (nontraumatic), fetus or newborn P52.6</u> - - <u>cerebral, cerebrum, fetus or newborn (anoxic) P52.4</u> - - - <u>birth injury P10.1</u> - - <u>complicating delivery</u> - - - <u>affecting fetus or newborn P02.1</u> 			
--	---	--	--	--

	<p><u>- - - associated with coagulation defect, maternal, affecting fetus or newborn P03.8</u> <u>- - - - afibrinogenemia P03.8</u> <u>- - - due to</u> <u>- - - - low-lying placenta, affecting fetus or newborn P02.0</u> <u>- - - - placenta previa, affecting fetus or newborn P02.0</u> <u>- - - - premature separation of placenta, affecting fetus or newborn P02.1</u> <u>- - - - trauma, affecting fetus or newborn P03.8</u> <u>- - - - uterine leiomyoma, affecting fetus or newborn P03.8</u> <u>- - conjunctiva, newborn P54.8</u> <u>- - cord, newborn (stump) P51.9</u> <u>- - cutaneous, fetus or newborn P54.5</u> <u>- - disease, fetus or newborn P53</u> <u>- - epicranial subaponeurotic (massive), birth injury P12.2</u> <u>- - extradural, fetus or newborn (anoxic) (nontraumatic) P52.8</u> <u>- - - birth injury P10.8</u> <u>- - fetal, fetus (<i>see also</i> Conditions arising in the perinatal period, hemorrhage, by specified sites) P50.9</u> <u>- - - from</u> <u>- - - - cut end of co-twin's cord P50.5</u> <u>- - - - placenta P50.2</u> <u>- - - - ruptured cord P50.1</u> <u>- - - - vasa previa P50.0</u> <u>- - - into</u> <u>- - - - co-twin P50.3</u> <u>- - - - maternal circulation P50.4</u> <u>- - - specified NEC P50.8</u> <u>- - fetal-maternal P50.4</u> <u>- - gastroenteric, newborn P54.3</u> <u>- - gastrointestinal (tract), newborn P54.3</u> <u>- - internal (organs) NEC, newborn P54.8</u> <u>- - intestine, newborn P54.3</u> <u>- - intra-alveolar (lung), newborn P26.8</u> <u>- - intracerebral (nontraumatic), fetus or newborn P52.4</u> <u>- - - birth injury P10.1</u> <u>- - intracranial, fetus or newborn (nontraumatic) P52.9</u> <u>- - - birth injury P10.9</u> <u>- - - specified NEC P52.8</u> <u>- - intraventricular, fetus or newborn (nontraumatic) P52.3</u> <u>- - - - birth injury P10.2</u> <u>- - - - grade</u> <u>- - - - - 1 P52.0</u></p>			
--	--	--	--	--

	<p> <u>----- 2 P52.1</u> <u>----- 3 P52.2</u> <u>-- lung, newborn P26.9</u> <u>--- massive P26.1</u> <u>--- specified NEC P26.8</u> <u>-- massive umbilical, newborn P51.0</u> <u>-- maternal, gestational, affecting fetus or newborn P02.1</u> <u>-- mucous membrane NEC, newborn P54.8</u> <u>-- nasal turbinate, newborn P54.8</u> <u>-- navel, newborn P51.9</u> <u>-- newborn P54.9</u> <u>--- specified NEC P54.8</u> <u>-- nose, newborn P54.8</u> <u>-- placenta NEC, affecting fetus or newborn P02.1</u> <u>--- from surgical or instrumental damage, affecting fetus or newborn P02.1</u> <u>--- previa, affecting fetus or newborn P02.0</u> <u>-- posterior fossa (nontraumatic), fetus or newborn P52.6</u> <u>-- pregnancy - <i>see also</i> Conditions originating in the perinatal period,</u> <u>hemorrhage, antepartum</u> <u>--- due to</u> <u>---- abruptio placentae, affecting fetus or newborn P02.1</u> <u>---- afibrinogenemia, or other coagulation defect (conditions in category D65-</u> <u>D68), affecting fetus or newborn P02.1</u> <u>---- leiomyoma, uterus, affecting fetus or newborn P02.1</u> <u>---- placenta previa, affecting fetus or newborn P02.0</u> <u>---- premature separation of placenta (normally implanted), affecting fetus or</u> <u>newborn P02.1</u> <u>---- threatened abortion, affecting fetus or newborn P02.1</u> <u>---- trauma, affecting fetus or newborn P02.1</u> <u>--- early, affecting fetus or newborn P02.1</u> <u>-- pulmonary, newborn P26.9</u> <u>--- massive P26.1</u> <u>--- specified NEC P26.8</u> <u>-- rectum (sphincter), newborn P54.2</u> <u>-- skin, fetus or newborn P54.5</u> <u>-- slipped umbilical ligature P51.8</u> <u>-- spinal (cord), fetus or newborn (birth injury) P11.5</u> <u>-- stomach, newborn P54.3</u> <u>-- subarachnoid, fetus or newborn (nontraumatic) P52.5</u> <u>--- birth injury P10.3</u> <u>-- subconjunctival, birth injury P15.3</u> </p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>subdural (acute)</u> - - - <u>fetus or newborn (nontraumatic) (anoxic) (hypoxic) P52.8</u> - - - - <u>birth injury P10.0</u> - - <u>subependymal, fetus or newborn P52.0</u> - - - <u>with intraventricular extension P52.1</u> - - - - <u>and intracerebral extension P52.2</u> - - <u>suprarenal (capsule) (gland), newborn P54.4</u> - - <u>tentorium (traumatic) NEC, fetus or newborn (birth injury) P10.4</u> - - <u>tracheobronchial, newborn P26.0</u> - - <u>umbilicus, umbilical</u> - - - <u>cord,</u> - - - - <u>after birth, newborn P51.9</u> - - - - <u>fetus, from ruptured cord P50.1</u> - - - <u>newborn P51.9</u> - - - - <u>massive P51.0</u> - - - - <u>slipped ligature P51.8</u> - - - <u>stump P51.9</u> - - <u>unavoidable (anteartum) (due to placenta previa), affecting fetus or newborn P02.0</u> - - <u>vagina (abnormal), newborn P54.6</u> - - <u>vasa previa, affecting fetus or newborn P50.0</u> - - <u>viscera NEC, newborn P54.8</u> - <u>hemothorax (bacterial) (nontuberculous), newborn P54.8</u> - <u>hepatitis</u> - - <u>fetus or newborn P59.2</u> - - <u>in toxoplasmosis, congenital (active) P37.1† K77.0*</u> - - <u>neonatal (toxic) P59.2</u> - - <u>viral, virus (acute), congenital P35.3</u> - <u>herpes simplex, congenital P35.2</u> - <u>hyaline membrane (disease) (lung) (pulmonary) (newborn) P22.0</u> - <u>hydramnios, affecting fetus or newborn P01.3</u> - <u>hydrocele (spermatic cord) (testis) (tunica vaginalis), congenital P83.5</u> - - <u>fetus or newborn P83.5</u> - <u>hydrocephalus (acquired) (external) (internal) (malignant) (recurrent)</u> - - <u>causing disproportion, affecting fetus or newborn P03.1</u> - - <u>due to toxoplasmosis (congenital) P37.1</u> - <u>hydrops fetal(is) or newborn (idiopathic) P83.2</u> - - <u>due to</u> - - - <u>ABO isoimmunization P56.0</u> - - - <u>hemolytic disease NEC P56.9</u> - - - <u>isoimmunization (ABO) (Rh) P56.0</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - - <u>Rh incompatibility P56.0</u> - <u>hyperbilirubinemia NEC</u> - - <u>neonatal (transient) (see also Conditions originating in the perinatal period, jaundice, fetus or newborn) P59.9</u> - - - <u>of prematurity P59.0</u> - <u>hyperemesis (see also Conditions originating in the perinatal period, vomiting)</u> - - <u>gravidarum, maternal (mild), affecting fetus or newborn P01.8</u> - <u>hypermagnesemia, neonatal P71.8</u> - <u>hypermaturity (fetus or newborn) P08.2</u> - <u>hypertension, hypertensive (accelerated) (benign) (essential) (idiopathic) (malignant) (primary) (systemic)</u> - - <u>maternal, complicating pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P00.0</u> - - <u>newborn P29.2</u> - - - <u>pulmonary (persistent) P29.3</u> - - <u>pulmonary (artery), of newborn (persistent) P29.3</u> - <u>hyperthermia (of unknown origin), newborn, environmental P81.0</u> - <u>hyperthyroidism (apathetic) (latent) (pre-adult) (recurrent), neonatal, transitory P72.1</u> - <u>hypertony, hypertonia, hypertonicity</u> - - <u>congenital P94.1</u> - - <u>uterus, uterine (contractions) (complicating delivery), affecting fetus or newborn P03.6</u> - <u>hypertrophy, hypertrophic breast, fetus or newborn P83.4</u> - <u>hypocalcemia, neonatal P71.1</u> - - <u>due to cow's milk P71.0</u> - - <u>phosphate-loading (newborn) P71.1</u> - <u>hypoglycemia (spontaneous)</u> - - <u>in infant of diabetic mother P70.1</u> - - - <u>gestational diabetes P70.0</u> - - <u>neonatal (transitory) P70.4</u> - - - <u>iatrogenic P70.3</u> - - - <u>maternal diabetes P70.1</u> - - - - <u>gestational P70.0</u> - - <u>transitory neonatal P70.4</u> - <u>hypomagnesemia, neonatal P71.2</u> - <u>hypoparathyroidism, neonatal, transitory P71.4</u> - <u>hypoplasia, hypoplastic</u> - - <u>lung (congenital) (lobe), associated with short gestation P28.0</u> - - <u>pulmonary, associated with short gestation P28.0</u> - <u>hypoprothrombinemia (congenital) (hereditary) (idiopathic), newborn, transient</u> 			
--	---	--	--	--

	<p><u>P61.6</u> - <u>hypothermia (accidental) (due to)</u> - - <u>neonatal P80.9</u> - - - <u>environmental (mild) NEC P80.8</u> - - - <u>mild P80.8</u> - - - <u>severe (chronic) (cold injury syndrome) P80.0</u> - - - <u>specified NEC P80.8</u> - <u>hypothyroidism (acquired), neonatal, transitory P72.2</u> - <u>hypotonia, hypotonicity, hypotony NEC, congenital (benign) P94.2</u> - <u>hypoxia - see also Conditions originating in the perinatal period, anoxia</u> - - <u>fetal - see Conditions originating in the perinatal period, distress, fetal</u> - - <u>intrauterine P20.9</u> - - - <u>first noted</u> - - - - <u>before onset of labor P20.0</u> - - - - <u>during labor and delivery P20.1</u> - - <u>newborn (see also Conditions originating in the perinatal period, asphyxia, newborn) P21.9</u> - <u>hysterotomy, affecting fetus or newborn P03.8</u> - <u>icterus neonatorum (see also Conditions originating in the perinatal period, jaundice, fetus or newborn) P59.9</u> - <u>ileus (bowel) (colon) (inhibitory) (intestine) (neurogenic), newborn, transitory P76.1</u> - <u>imbalance, electrolyte</u> - - <u>neonatal, transitory NEC P74.4</u> - - - <u>potassium P74.3</u> - - - <u>sodium P74.2</u> - <u>immature, birth (28 completed weeks or more but less than 37 completed weeks) P07.3</u> - - - <u>extremely (less than 28 completed weeks) P07.2</u> - <u>immaturity (28 completed weeks or more but less than 37 completed weeks) P07.3</u> - - <u>extreme (less than 28 completed weeks) P07.2</u> - - <u>fetal, fetus P07.3</u> - - <u>fetus or infant light-for-dates – see Conditions originating in the perinatal period, light-for-dates</u> - - <u>gross P07.2</u> - - <u>infant P07.3</u> - - <u>newborn P07.3</u> - - <u>pulmonary, fetus or newborn P28.0</u> - - <u>respiratory P28.0</u> - <u>immunization</u></p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>ABO (see also Conditions originating in the perinatal period, isoimmunization, ABO)</u> - - - <u>affecting management of pregnancy, in fetus or newborn P55.1</u> - <u>impaction, impacted shoulder, affecting fetus or newborn P03.1</u> - <u>imperfect aeration, lung (newborn) NEC P28.1</u> - <u>improperly tied umbilical cord (causing hemorrhage) P51.8</u> - <u>inability to breathe properly, newborn P28.8</u> - <u>inadequate, inadequacy</u> - - <u>development</u> - - - <u>fetus P05.9</u> - - - <u>lungs, associated with short gestation P28.0</u> - - <u>pulmonary function, newborn P28.5</u> - - - <u>ventilation, newborn P28.5</u> - - <u>ventilation, newborn P28.5</u> - <u>incompatibility</u> - - <u>ABO, fetus or newborn P55.1</u> - - <u>blood (group) (Duffy) (K(ell)) (Kidd) (Lewis) (M) (S) NEC</u> - - - <u>fetus or newborn P55.8</u> - - <u>Rh (blood group) (factor), fetus or newborn P55.0</u> - <u>incompatible with life (newborn) (nonviable) P07.2</u> - <u>incompetency, incompetent cervix, cervical (os), maternal</u> - - <u>in pregnancy, affecting fetus or newborn P01.0</u> - <u>incomplete – see also condition</u> - - <u>expansion, lungs (newborn) NEC P28.1</u> - <u>incoordinate, incoordination</u> - - <u>uterus (action) (contractions) (complicating delivery)</u> - - - <u>affecting fetus or newborn P03.6</u> - <u>induction of labor, affecting fetus or newborn P03.8</u> - <u>inertia uterus, uterine during labor</u> - - <u>affecting fetus or newborn P03.6</u> - <u>infancy, infantile, infantilism</u> - - <u>genitalia, genitals, maternal (after puberty)</u> - - - <u>in pregnancy or childbirth NEC, affecting fetus or newborn P03.8</u> - - <u>pelvis, maternal</u> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>infant(s) - see also Conditions originating in the perinatal period, infancy</u> - - <u>of diabetic mother (syndrome of) P70.1</u> - - - <u>gestational diabetes P70.0</u> - <u>infarct, infarction (of), placenta (complicating pregnancy), affecting fetus or newborn P02.2</u> - <u>infection, infected (opportunistic) (see also Conditions originating in the perinatal</u> 			
--	---	--	--	--

	<p><u>period, inflammation)</u> <u>-- amniotic fluid, sac or cavity, affecting fetus or newborn P02.7</u> <u>-- <i>Candida (albicans) (tropicalis)</i>,</u> <u>--- neonatal P37.5</u> <u>--- congenital P37.5</u> <u>-- <i>Citrobacter</i>, newborn P37.8</u> <u>-- <i>Clostridium</i>, clostridium, congenital P39.8</u> <u>-- congenital NEC P39.9</u> <u>--- <i>Candida (albicans)</i> P37.5</u> <u>--- clostridium, other than <i>Clostridium tetani</i> P39.8</u> <u>--- cytomegalovirus P35.1</u> <u>--- <i>Escherichia coli</i> P39.8</u> <u>---- sepsis P36.4</u> <u>--- hepatitis, viral P35.3</u> <u>--- herpes simplex P35.2</u> <u>--- infectious or parasitic disease P37.9</u> <u>---- specified NEC P37.8</u> <u>--- listeriosis (disseminated) P37.2</u> <u>--- malaria NEC P37.4</u> <u>---- <i>falciparum</i> P37.3</u> <u>--- <i>Plasmodium falciparum</i> P37.3</u> <u>--- poliomyelitis P35.8</u> <u>--- rubella P35.0</u> <u>--- salmonella P39.8</u> <u>--- skin P39.4</u> <u>--- streptococcal NEC P39.8</u> <u>---- sepsis P36.1</u> <u>---- group B P36.0</u> <u>--- toxoplasmosis (acute) (chronic) (subacute) P37.1</u> <u>--- tuberculosis P37.0</u> <u>--- urinary (tract) P39.3</u> <u>--- vaccinia P35.8</u> <u>--- virus P35.9</u> <u>---- specified type NEC P35.8</u> <u>-- cytomegalovirus, cytomegaloviral, congenital P35.1</u> <u>-- <i>Enterobacter (cloacae)</i>, newborn P37.8</u> <u>-- <i>Escherichia (E.) coli</i> NEC, congenital P39.8</u> <u>--- sepsis P36.4</u> <u>-- fetus (<i>see also</i> Conditions originating in the perinatal period, infection, congenital) P39.9</u> <u>--- intra-amniotic NEC P39.2</u></p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>genital organ or tract, maternal</u> - - - <u>complicating pregnancy, affecting fetus or newborn P00.8</u> - - <u>herpes (simplex), congenital P35.2</u> - - <u>intra-amniotic, fetus P39.2</u> - - <u>intrauterine (complicating pregnancy)</u> - - - <u>fetus or newborn P00.8</u> - - - <u>specified infection NEC, fetus P39.2</u> - - <u>kidney, maternal (cortex) (hematogenous)</u> - - - <u>complicating pregnancy, affecting fetus or newborn P00.1</u> - - <u>Listeria monocytogenes, congenital P37.2</u> - - <u>Monilia, neonatal P37.5</u> - - <u>navel, newborn P38</u> - - <u>newborn P39.9</u> - - - <u>skin P39.4</u> - - - <u>specified type NEC (see also Conditions originating in the perinatal period, infection, congenital) P39.8</u> - - <u>perinatal period NEC P39.9</u> - - - <u>specified type NEC P39.8</u> - - <u>polymicrobial, newborn P37.9</u> - - <u>respiratory (tract) NEC</u> - - - <u>fetus P28.8</u> - - - <u>newborn, neonatal P28.8</u> - - <u>rubella, congenital P35.0</u> - - <u>Salmonella (arizonae) (cholerae-suis) (enteritidis) (typhimurium)</u> - - - <u>congenital P39.8</u> - - <u>skin (local) (staphylococcal) (streptococcal)</u> - - - <u>newborn P39.4</u> - - <u>staphylococcal NEC</u> - - - <u>hemolytic, newborn P36.3</u> - - - <u>newborn P37.8</u> - - <u>streptococcal NEC</u> - - - <u>congenital P39.8</u> - - - - <u>sepsis P36.1</u> - - - - <u>group B P36.0</u> - - <u>umbilicus, newborn P38</u> - - <u>urinary (tract) NEC</u> - - - <u>maternal, complicating pregnancy, affecting fetus or newborn P00.1</u> - - - <u>newborn P39.3</u> - <u>inflammation, inflamed, inflammatory (with exudation) (see also Conditions originating in the perinatal period, infection)</u> - - <u>navel, newborn P38</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>umbilicus, newborn P38</u> - <u>influenza (specific virus not identified)</u> - - <u>maternal, affecting fetus or newborn P00.2</u> - <u>inhalation, meconium (newborn) P24.0</u> - <u>injury, injuries (see also specified injury type)</u> - - <u>birth (see also Conditions originating in the perinatal period, birth, injury) P15.9</u> - - <u>brachial plexus, newborn P14.3</u> - - <u>brain, anoxic</u> - - - <u>at birth P21.9</u> - - - <u>newborn P21.9</u> - - <u>childbirth (fetus or newborn) (see also Conditions originating in the perinatal period, birth, injury) P15.9</u> - - <u>delivery (fetus or newborn) P15.9</u> - - <u>maternal, during pregnancy, affecting fetus or newborn P00.5</u> - - <u>nerve, facial, newborn P11.3</u> - - <u>scalp, fetus or newborn (birth injury) P12.9</u> - - - <u>due to monitoring (electrode) (sampling incision) P12.4</u> - - - <u>specified NEC P12.8</u> - - <u>skeleton, skeletal, birth injury P13.9</u> - - - <u>specified part NEC P13.8</u> - <u>inспissated bile syndrome (newborn) P59.1</u> - <u>insufficiency, insufficient</u> - - <u>circulatory NEC, fetus or newborn P29.8</u> - - <u>heart, newborn P29.0</u> - - <u>lung, newborn P28.5</u> - - <u>myocardial, myocardium (acute) (chronic), newborn P29.0</u> - - <u>placental, affecting fetus or newborn P02.2</u> - - <u>pulmonary, newborn P28.5</u> - - <u>respiratory (acute), newborn P28.5</u> - - <u>ventilation, ventilatory, newborn P28.8</u> - <u>insufficiently tied umbilical cord P51.8</u> - <u>interruption (of)</u> - - <u>oxygen cycle, newborn P28.8</u> - - <u>respiration, newborn P28.8</u> - <u>intoxication NEC, drug, newborn P93</u> - <u>intrauterine contraceptive device (IUD), affecting fetus or newborn P01.8</u> - <u>irritable, irritability, cerebral, in newborn P91.3</u> - <u>ischemia, ischemic</u> - - <u>cerebral (chronic) (generalized)</u> - - - <u>newborn P91.0</u> - - - <u>prenatal P91.0</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>myocardium, myocardial, transient, of newborn P29.4</u> - <u>isoimmunization NEC (see also Conditions originating in the perinatal period, incompatibility)</u> - - <u>fetus or newborn P55.9</u> - - - <u>with</u> - - - - <u>hydrops fetalis P56.0</u> - - - - <u>kernicterus P57.0</u> - - - - <u>ABO P55.1</u> - - - - <u>Rh P55.0</u> - - - <u>specified type NEC P55.8</u> - <u>jaundice (yellow)</u> - - <u>breast-milk (inhibitor) P59.3</u> - - <u>due to or associated with</u> - - - <u>delivery due to delayed conjugation P59.0</u> - - - <u>preterm delivery P59.0</u> - - <u>fetus or newborn (physiological) P59.9</u> - - - <u>due to or associated with</u> - - - - <u>ABO</u> - - - - - <u>antibodies P55.1</u> - - - - - <u>incompatibility, maternal/fetal P55.1</u> - - - - - <u>isoimmunization P55.1</u> - - - - <u>absence or deficiency of enzyme system for bilirubin conjugation (congenital) P59.8</u> - - - - <u>bleeding P58.1</u> - - - - <u>breast milk inhibitors to conjugation P59.3</u> - - - - <u>bruising P58.0</u> - - - - <u>delayed conjugation P59.8</u> - - - - - <u>associated with preterm delivery P59.0</u> - - - - <u>drugs or toxins</u> - - - - - <u>given to newborn P58.4</u> - - - - - <u>transmitted from mother P58.4</u> - - - - <u>excessive hemolysis NEC P58.9</u> - - - - - <u>specified type NEC P58.8</u> - - - - <u>hepatocellular damage P59.2</u> - - - - <u>hereditary hemolytic anemia P58.8</u> - - - - <u>incompatibility, maternal/fetal NEC P55.9</u> - - - - <u>infection P58.2</u> - - - - <u>inspissated bile syndrome P59.1</u> - - - - <u>isoimmunization NEC P55.9</u> - - - - <u>polycythemia P58.3</u> - - - - <u>preterm delivery P59.0</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - - - Rh - - - - antibodies P55.0 - - - - incompatibility, maternal/fetal P55.0 - - - - isoimmunization P55.0 - - - - swallowed maternal blood P58.5 - - - specified cause NEC P59.8 - - nuclear, newborn – <i>see</i> Conditions originating in the perinatal period, kernicterus of newborn - kernicterus of newborn P57.9 - - due to isoimmunization (conditions in P55.-) P57.0 - - specified type NEC P57.8 - Klumpke(-Déjerine) palsy, paralysis (birth) (newborn) P14.1 - knot (true), umbilical cord, affecting fetus or newborn P02.5 - labor (<i>see also</i> Conditions originating in the perinatal period, delivery) - - abnormal NEC, affecting fetus or newborn P03.6 - - arrested active phase, affecting fetus or newborn P03.6 - - desultory, affecting fetus or newborn P03.6 - - dyscoordinate, affecting fetus or newborn P03.6 - - forced or induced, affecting fetus or newborn P03.8 - - hypertonic, affecting fetus or newborn P03.6 - - hypotonic, affecting fetus or newborn P03.6 - - incoordinate, affecting fetus or newborn P03.6 - - irregular, affecting fetus or newborn P03.6 - - obstructed - - - affecting fetus or newborn P03.1 - - - by or due to - - - - abnormal - - - - pelvis (bony), affecting fetus or newborn P03.1 - - - - presentation or position, affecting fetus or newborn P03.1 - - - - size, fetus, affecting fetus or newborn P03.1 - - - cephalopelvic disproportion (normally formed fetus), affecting fetus or newborn P03.1 - - - disproportion, fetopelvic NEC, affecting fetus or newborn P03.1 - - - malpresentation, affecting fetus or newborn P03.1 - - precipitate, affecting fetus or newborn P03.5 - - prolonged or protracted, affecting fetus or newborn P03.8 - laceration (<i>see also</i> Conditions originating in the perinatal period, wound) - - brain (any part) (cortex) (diffuse) (membrane), during birth P10.8 - - - with hemorrhage P10.1 - - cerebral (diffuse), during birth P10.8 - - - with hemorrhage P10.1 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>intracranial NEC, birth injury P10.9</u> - - <u>spinal cord (meninges), fetus or newborn (birth injury) P11.5</u> - <u>lack of exchange of gases, newborn P28.5</u> - <u>large-for-dates NEC (fetus or infant) P08.1</u> - - <u>exceptionally (4500 g or more) P08.0</u> - <u>laryngismus (stridulus), congenital P28.8</u> - <u>lateroversion, uterus, uterine, maternal (cervix) (postinfectious) (postpartal, old)</u> - - <u>in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - <u>leak, leakage of air (bronchus) (intrathoracic) (lung) (pleural) (pulmonary) (thorax), newborn P25.1</u> - <u>lesion, vascular (nontraumatic)</u> - - <u>umbilical cord, complicating delivery</u> - - - <u>affecting fetus or newborn P02.6</u> - <u>leukomalacia, cerebral, newborn P91.2</u> - <u>lie, abnormal (maternal care) (see also Conditions originating in the perinatal period, presentation, fetal, abnormal)</u> - - <u>before labor, affecting fetus or newborn P01.7</u> - <u>light, fetus or newborn, for gestational age P05.0</u> - <u>light-for-dates (infant) P05.0</u> - - <u>and small-for-dates P05.1</u> - <u>listeriosis, listerellosis, congenital (disseminated) P37.2</u> - - <u>fetal P37.2</u> - - <u>neonatal (disseminated) P37.2</u> - <u>lithopedion P95</u> - <u>locked twins causing obstructed labor, affecting fetus or newborn P03.1</u> - <u>long labor, affecting fetus or newborn P03.8</u> - <u>loss of fluid (acute), fetus or newborn P74.1</u> - <u>low</u> - - <u>birthweight (2499 grams or less) (see also Conditions originating in the perinatal period, weight) P07.1</u> - - - <u>extreme (999 grams or less) P07.0</u> - - - <u>for gestational age P05.0</u> - - <u>cardiac, output, newborn P29.8</u> - <u>lupus erythematosus, systemic (discoid) (local)</u> - - <u>maternal, affecting fetus or newborn P00.8</u> - <u>luxation, eyeball, birth injury P15.3</u> - <u>lymphadenopathy (generalized), due to toxoplasmosis</u> - - <u>congenital (acute) (chronic) (subacute) P37.1</u> - <u>maceration of fetus or newborn (cause not stated) P95</u> - <u>malaria, malarial fever, congenital NEC P37.4</u> - - <u>falciparum P37.3</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - <u>male type pelvis, maternal</u> - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>malformation (congenital) - see also Conditions originating in the perinatal period, anomaly</u> - - <u>pelvic organs or tissues NEC, maternal</u> - - - <u>in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>umbilical cord NEC (complicating delivery)</u> - - - <u>affecting fetus or newborn P02.6</u> - <u>malnutrition</u> - - <u>intrauterine or fetal P05.2</u> - - - <u>light-for-dates P05.0</u> - - - <u>small-for-dates P05.1</u> - - <u>maternal, affecting fetus or newborn P00.4</u> - <u>mastitis (acute) (nonpuerperal) (subacute)</u> - - <u>infective, newborn or neonatal P39.0</u> - - <u>noninfective, newborn or neonatal P83.4</u> - <u>maternal condition, affecting fetus or newborn P00.9</u> - - <u>acute yellow atrophy of liver P00.8</u> - - <u>alcohol use P04.3</u> - - <u>anesthesia or analgesia P04.0</u> - - <u>blood loss (gestational) P02.1</u> - - <u>cancer chemotherapy P04.1</u> - - <u>chorioamnionitis P02.7</u> - - <u>circulatory disease (conditions in I00-I99, Q20-Q28) P00.3</u> - - <u>complication of pregnancy NEC P01.9</u> - - <u>congenital heart disease (conditions in Q20-Q24) P00.3</u> - - <u>cortical necrosis of kidney P00.1</u> - - <u>cytotoxic drug P04.1</u> - - <u>death P01.6</u> - - <u>diabetes mellitus (conditions in E10-E14) P70.1</u> - - <u>disease NEC P00.9</u> - - <u>drug abuse P04.4</u> - - <u>eclampsia P00.0</u> - - <u>exposure to environmental chemical substances P04.6</u> - - <u>genital tract infections NEC P00.8</u> - - <u>glomerular diseases (conditions in N00- N08) P00.1</u> - - <u>hemorrhage, gestational P02.1</u> - - <u>hepatitis, acute, malignant or subacute P00.8</u> - - <u>hyperemesis (gravidarum) P01.8</u> - - <u>hypertension (conditions in O10-O11, O13-O16) P00.0</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>infectious and parasitic diseases (conditions in A00-B99, J10-J11) P00.2</u> - - <u>influenza P00.2</u> - - - <u>manifest influenza in the infant P35.8</u> - - <u>injury (conditions in S00-T79) P00.5</u> - - <u>intrauterine coil P01.8</u> - - <u>malaria P00.2</u> - - - <u>manifest malaria NEC in infant or fetus P37.4</u> - - - - <u>falciparum P37.3</u> - - <u>malnutrition P00.4</u> - - <u>necrosis of liver P00.8</u> - - <u>nephritis, nephrotic syndrome and nephrosis (conditions in N00-N08) P00.1</u> - - <u>noxious influence transmitted via breast milk or placenta P04.9</u> - - - <u>specified NEC P04.8</u> - - <u>nutritional disorder (conditions in E40- E64) P00.4</u> - - <u>operation unrelated to current pregnancy P00.6</u> - - <u>pre-eclampsia P00.0</u> - - <u>previous surgery, uterus or pelvic organs P03.8</u> - - <u>proteinuria P00.1</u> - - <u>pyelitis or pyelonephritis P00.1</u> - - <u>renal disease or failure P00.1</u> - - <u>respiratory disease (conditions in J00-J99, Q30-Q34) P00.3</u> - - <u>rheumatic heart disease (chronic) (conditions in I05-I09) P00.3</u> - - <u>rubella (conditions in B06) P00.2</u> - - - <u>manifest rubella in the infant or fetus P35.0</u> - - <u>septate vagina P03.8</u> - - <u>stenosis or stricture of vagina P03.8</u> - - <u>surgery unrelated to current pregnancy P00.6</u> - - - <u>to uterus or pelvic organs P03.8</u> - - <u>syphilis (conditions in A50-A53) P00.2</u> - - <u>thrombophlebitis P00.3</u> - - <u>tobacco use P04.2</u> - - <u>toxemia (of pregnancy) P00.0</u> - - <u>toxoplasmosis (conditions in B58.-) P00.2</u> - - - <u>manifest toxoplasmosis (acute) (chronic) (subacute) in the infant or fetus P37.1</u> - - <u>transmission of chemical substance through the placenta (see also Conditions originating in the perinatal period, absorption, chemical, through placenta) P04.8</u> - - <u>uremia P00.1</u> - - <u>urinary tract conditions (conditions in N00- N39) P00.1</u> - - <u>vomiting (pernicious) (persistent) (vicious) P01.8</u> - <u>meconium</u> - - <u>obstruction, fetus or newborn P76.0</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>peritonitis P78.0</u> - - <u>plug syndrome (newborn) NEC P76.0</u> - - <u>stain, fetus or newborn P20.9</u> - <u>melena, newborn, neonatal P54.1</u> - - <u>due to swallowed maternal blood P78.2</u> - <u>membranitis, affecting fetus or newborn P02.7</u> - <u>meningoencephalitis, in toxoplasmosis, congenital P37.1† G05.2*</u> - <u>meningoencephalomyelitis, due to toxoplasma or toxoplasmosis, congenital P37.1† G05.2*</u> - <u>microcephalus, microcephalic, microcephaly, due to toxoplasmosis (congenital) P37.1</u> - <u>Mikity-Wilson disease or syndrome P27.0</u> - <u>molding, head (during birth) P13.1</u> - <u>mole (pigmented), hydatid, hydatidiform (benign) (complicating pregnancy) (delivered) (undelivered)</u> - - <u>newborn P02.2</u> - <u>moniliasis, neonatal P37.5</u> - <u>mucus</u> - - <u>asphyxia or suffocation, newborn P24.1</u> - - <u>plug</u> - - - <u>aspiration, newborn P24.1</u> - - - <u>tracheobronchial, newborn P24.1</u> - <u>multiple, multiplex</u> - - <u>birth, affecting fetus or newborn P01.5</u> - <u>myasthenia, myasthenic gravis, neonatal, transient P94.0</u> - <u>necrosis, necrotic (ischemic) (necrotizing)</u> - - <u>fat (generalized)</u> - - - <u>subcutaneous, due to birth injury P15.6</u> - - <u>kidney (bilateral)</u> - - - <u>tubular, complicating pregnancy , affecting fetus or newborn P00.1</u> - - <u>liver (cell), maternal</u> - - - <u>complicating pregnancy or childbirth, affecting fetus or newborn P00.8</u> - - <u>subcutaneous fat, fetus or newborn P83.8</u> - - <u>umbilical cord, affecting fetus or newborn P02.6</u> - <u>nephritis, nephritic, maternal, complicating pregnancy or childbirth</u> - - <u>with secondary hypertension, pre-existing, affecting fetus or newborn P00.0</u> - - <u>affecting fetus or newborn P00.1</u> - <u>neuritis, cranial nerve</u> - - <u>seventh or facial, newborn (birth injury) P11.3</u> - <u>neutropenia, neutropenic (congenital) (cyclic) (drug-induced) (periodic) (primary) (splenic) (toxic)</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>neonatal, transitory (isoimmune) (maternal transfer) P61.5</u> - <u>nonexpansion, lung (newborn) P28.0</u> - <u>nonviable P07.2</u> - <u>nuchal cord, newborn P02.5</u> - <u>obstetric trauma NEC (complicating delivery), affecting fetus or newborn P03.8</u> - <u>obstruction, obstructed, obstructive</u> - - <u>intestine (mechanical) (neurogenic) (paroxysmal) (postinfective) (reflex)</u> - - - <u>newborn P76.9</u> - - - - <u>due to</u> - - - - - <u>fecaliths P76.8</u> - - - - - <u>inspissated milk P76.2</u> - - - - - <u>meconium (plug) P76.0</u> - - - - <u>specified NEC P76.8</u> - - <u>labor (see also Conditions originating in the perinatal period, labor, obstructed)</u> - - - <u>affecting fetus or newborn P03.1</u> - - <u>meconium plug, newborn P76.0</u> - <u>oligohydramnios, affecting fetus or newborn P01.2</u> - <u>omphalitis (congenital) (newborn) (with mild hemorrhage) P38</u> - <u>omphalorrhagia, newborn P51.9</u> - <u>operation</u> - - <u>for delivery, affecting fetus or newborn (see also Conditions originating in the perinatal period, delivery, by type, affecting fetus) P03.8</u> - - <u>maternal, unrelated to current delivery, affecting fetus or newborn P00.6</u> - <u>ophthalmia neonatorum, newborn P39.1</u> - <u>overfeeding, newborn P92.4</u> - <u>oversize fetus P08.1</u> - - <u>causing disproportion, affecting fetus or newborn P03.1</u> - - <u>exceptionally large (more than 4500 g) P08.0</u> - <u>palpitations (heart)</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>palsy (see also Conditions originating in the perinatal period, paralysis)</u> - - <u>brachial plexus NEC</u> - - - <u>fetus or newborn (birth injury) P14.3</u> - - <u>Erb's P14.0</u> - - <u>facial, newborn (birth injury) P11.3</u> - - <u>Klumpke(-Déjerine) P14.1</u> - <u>papyraceous fetus P95</u> - <u>paralysis, paralytic (complete) (incomplete)</u> - - <u>birth injury P14.9</u> - - <u>brachial plexus NEC, birth injury P14.3</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - - <u>newborn P14.3</u> - - <u>Duchenne's, birth injury P14.0</u> - - <u>Erb(-Duchenne) (birth) (newborn) P14.0</u> - - <u>facial (nerve), birth injury P11.3</u> - - - <u>newborn P11.3</u> - - <u>Klumpke(-Déjerine) (birth injury) (newborn) P14.1</u> - - <u>nerve</u> - - - <u>birth injury P14.9</u> - - - <u>facial, birth injury (newborn) P11.3</u> - - - <u>newborn (birth injury) P14.9</u> - - - <u>phrenic (birth injury) (newborn) P14.2</u> - - - <u>radial, birth injury (newborn) P14.3</u> - - - <u>seventh or facial (birth injury) (newborn) P11.3</u> - - <u>newborn P91.8</u> - - - <u>birth injury P11.9</u> - - <u>radial nerve, birth injury (newborn) P14.3</u> - <u>paraplegia (lower), newborn P91.8</u> - - <u>birth injury P11.9</u> - <u>patent cervix, maternal, complicating pregnancy</u> - - <u>affecting fetus or newborn P01.0</u> - <u>pendulous abdomen, maternal, in pregnancy or childbirth</u> - - <u>affecting fetus or newborn P03.8</u> - <u>perforation, perforated (nontraumatic) (see also Conditions originating in the perinatal period, rupture)</u> - - <u>intestine (bowel) (colon) (ileum) (jejunum) (rectum) (sigmoid), fetus or newborn P78.0</u> - <u>peritonitis (adhesive) (fibrinous) (with effusion)</u> - - <u>congenital NEC P78.1</u> - - <u>meconium (newborn) P78.0</u> - - <u>neonatal P78.1</u> - - - <u>meconium P78.0</u> - <u>persistence, persistent (congenital) fetal circulation P29.3</u> - - <u>occipitoposterior or transverse (position), affecting fetus or newborn P03.1</u> - <u>petechia, petechiae, fetus or newborn P54.5</u> - <u>pithecoïd maternal pelvis, with disproportion (fetopelvic)</u> - - <u>affecting fetus or newborn P03.1</u> - <u>placenta, placental (see also condition)</u> - - <u>ablatio, affecting fetus or newborn P02.1</u> - - <u>abnormal, abnormality NEC</u> - - - <u>with hemorrhage, affecting fetus or newborn P02.1</u> - - - <u>affecting fetus or newborn P02.2</u> 			
--	--	--	--	--

	<p>- - <u>abruptio, affecting fetus or newborn P02.1</u> - - <u>disease NEC, affecting fetus or newborn P02.2</u> - - <u>infarction, affecting fetus or newborn P02.2</u> - - <u>insufficiency, affecting fetus or newborn P02.2</u> - - <u>marginal (hemorrhage) (rupture), affecting fetus or newborn P02.0</u> - - <u>previa (central) (complete) (marginal) (partial) (total) (with hemorrhage), affecting fetus or newborn P02.0</u> - <u>placental, affecting fetus or newborn P02.7</u> - <u>platypelloid maternal pelvis, with disproportion (fetopelvic)</u> - - <u>affecting fetus or newborn P03.1</u> - <u>plug, meconium (newborn), syndrome P76.0</u> - <u>pneumomediastinum, congenital or perinatal (newborn) P25.2</u> - <u>pneumonia (acute) (community acquired) (double) (hemorrhagic) (lobe) (migratory) (nosocomial) (primary) (purulent) (septic) (unresolved) (see also Conditions originating in the perinatal period, pneumonitis)</u> - - <u>aspiration</u> - - - <u>newborn P24.9</u> - - - - <u>meconium P24.0</u> - - <u>chlamydial, congenital P23.1</u> - - <u>congenital (infective) P23.9</u> - - - <u>due to</u> - - - - <u>bacterium NEC P23.6</u> - - - - <u>Chlamydia P23.1</u> - - - - <u>Escherichia coli P23.4</u> - - - - <u>Haemophilus influenzae P23.6</u> - - - - <u>infective organism NEC P23.8</u> - - - - <u>Klebsiella pneumoniae P23.6</u> - - - - <u>Mycoplasma P23.6</u> - - - - <u>Pseudomonas P23.5</u> - - - - <u>staphylococcus P23.2</u> - - - - <u>streptococcus (except group B) P23.6</u> - - - - - <u>group B P23.3</u> - - - - <u>viral agent P23.0</u> - - - <u>specified NEC P23.8</u> - - <u>in (due to), Chlamydia, neonatal P23.1</u> - - <u>meconium P24.0</u> - - <u>neonatal P23.9</u> - - - <u>aspiration P24.9</u> - - <u>viral, virus (broncho) (interstitial) (lobar), congenital P23.0</u> - <u>pneumonitis (acute) (primary) (see also Conditions originating in the perinatal period, pneumonia)</u></p>			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>due to toxoplasmosis, congenital P37.1† J17.3*</u> - - <u>meconium P24.0</u> - - <u>neonatal aspiration P24.9</u> - - <u>rubella, congenital P35.0</u> - <u>pneumopericardium</u> - - <u>congenital P25.3</u> - - <u>fetus or newborn P25.3</u> - <u>pneumothorax</u> - - <u>congenital P25.1</u> - - <u>newborn P25.1</u> - - <u>perinatal period P25.1</u> - - <u>spontaneous NEC</u> - - - <u>fetus or newborn P25.1</u> - <u>poliomyelitis (acute) (anterior) (epidemic), congenital P35.8</u> - <u>polycythemia (primary) (rubra) (vera), neonatorum P61.1</u> - <u>polyhydramnios, affecting fetus or newborn P01.3</u> - <u>polyp, polypus</u> - - <u>cervix (uteri), maternal, in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - - <u>umbilical, newborn P83.6</u> - - <u>uterus (body) (corpus) (mucous), maternal, in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - <u>poor</u> - - <u>contractions, labor, affecting fetus or newborn P03.6</u> - - <u>fetal growth NEC P05.9</u> - <u>postmaturity, postmature (fetus or newborn) P08.2</u> - <u>postterm (pregnancy), infant P08.2</u> - <u>precipitate labor or delivery, affecting fetus or newborn P03.5</u> - <u>pre-eclampsia, affecting fetus or newborn P00.0</u> - <u>pregnancy (single) (uterine)</u> - - <u>abdominal (ectopic), affecting fetus or newborn P01.4</u> - - <u>cornual, fetus or newborn P01.4</u> - - <u>ectopic (ruptured), affecting fetus or newborn P01.4</u> - - <u>multiple NEC, affecting fetus or newborn P01.5</u> - - <u>mural, fetus or newborn P01.4</u> - - <u>ovarian, fetus or newborn P01.4</u> - - <u>quadruplet, affecting fetus or newborn P01.5</u> - - <u>quintuplet, affecting fetus or newborn P01.5</u> - - <u>sextuplet, affecting fetus or newborn P01.5</u> - - <u>triplet, affecting fetus or newborn P01.5</u> - - <u>tubal (with abortion) (with rupture), affecting fetus or newborn P01.4</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - <u>twin, affecting fetus or newborn P01.5</u> - <u>premature</u> - - <u>birth NEC P07.3</u> - - <u>delivery, newborn NEC P07.3</u> - - <u>infant NEC P07.3</u> - - - <u>light-for-dates P05.0</u> - - <u>labor, newborn NEC P07.3</u> - - <u>lungs P28.0</u> - - <u>rupture, membranes or amnion, affecting fetus or newborn P01.1</u> - <u>prematurity NEC (less than 37 completed weeks) (see also Conditions originating in the perinatal period, immaturity) P07.3</u> - - <u>extreme (less than 28 completed weeks) P07.2</u> - - <u>gross P07.2</u> - - <u>marked P07.2</u> - - <u>severe P07.2</u> - <u>presentation, fetal</u> - - <u>abnormal</u> - - - <u>before labor, affecting fetus or newborn P01.7</u> - - - <u>causing obstructed labor, affecting fetus or newborn (any, except breech) P03.1</u> - - - - <u>breech P03.0</u> - - <u>breech</u> - - - <u>with external version before labor, affecting fetus or newborn P01.7</u> - - - <u>before labor, affecting fetus or newborn P01.7</u> - - - <u>causing obstructed labor, fetus or newborn P03.0</u> - <u>pressure</u> - - <u>birth, fetus or newborn, NEC P15.9</u> - - <u>brain injury at birth NEC P11.1</u> - - <u>increased, intracranial (benign), injury at birth P11.0</u> - <u>preterm infant, newborn NEC P07.3</u> - <u>previa</u> - - <u>placenta (with hemorrhage), affecting fetus or newborn P02.0</u> - - <u>vasa, affecting fetus or newborn P02.6</u> - <u>previable P07.2</u> - <u>problem (related to) (with) (see also Conditions originating in the perinatal period, disease)</u> - - <u>feeding newborn P92.9</u> - - - <u>breast P92.5</u> - - - <u>overfeeding P92.4</u> - - - <u>slow P92.2</u> - - - <u>specified NEC P92.8</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - - - <u>underfeeding P92.3</u> - <u>procedure (surgical), maternal (unrelated to current delivery), affecting fetus or newborn P00.6</u> - - <u>nonsurgical (medical) P00.7</u> - <u>prolapse, prolapsed</u> - - <u>arm or hand, in fetus or newborn P03.1</u> - - <u>fetal limb NEC, in fetus or newborn P03.1</u> - - <u>leg, in fetus or newborn P03.1</u> - - <u>umbilical cord, affecting fetus or newborn P02.4</u> - - <u>uterus (with prolapse of vagina), pregnant, affecting fetus or newborn P03.8</u> - <u>prolonged</u> - - <u>labor, affecting fetus or newborn P03.8</u> - - <u>uterine contractions in labor, affecting fetus or newborn P03.6</u> - <u>prominent ischial spine or sacral promontory, with disproportion (fetopelvic)</u> - - <u>affecting fetus or newborn P03.1</u> - <u>proteinuria, pre-eclamptic, affecting fetus or newborn P00.0</u> - <u>pseudomenses (newborn) P54.6</u> - <u>pseudomenstruation (newborn) P54.6</u> - <u>pseudoparalysis, atonic, congenital P94.2</u> - <u>pseudosclerema, newborn P83.8</u> - <u>pulse</u> - - <u>alternating</u> - - - <u>fetal P20.9</u> - - - <u>newborn P29.1</u> - - <u>bigeminal</u> - - - <u>fetal P20.9</u> - - - <u>newborn P29.1</u> - <u>pulsus alternans or trigeminus</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>purpura, thrombocytopenic (congenital) (hereditary), neonatal, transitory P61.0</u> - <u>pyelitis, maternal (congenital) (uremic), complicating pregnancy, affecting fetus or newborn P00.1</u> - <u>pyelonephritis, maternal, complicating pregnancy, affecting fetus or newborn P00.1</u> - <u>pyoderma, pyodermia NEC, newborn P39.4</u> - <u>pyrexia (of unknown origin), newborn, environmentally-induced P81.0</u> - <u>quadriplegia, newborn NEC P11.9</u> - <u>quadruplet, affecting fetus or newborn P01.5</u> - <u>quintuplet, affecting fetus or newborn P01.5</u> - <u>rachitic pelvis, maternal (late effect), with disproportion (fetopelvic), affecting</u> 			
--	--	--	--	--

	<p><u>fetus or newborn P03.1</u> <u>- radiology, maternal, affecting fetus or newborn P00.7</u> <u>- rapid</u> <u>- - heart (beat)</u> <u>- - - fetal P20.9</u> <u>- - - newborn P29.1</u> <u>- - second stage (delivery), affecting fetus or newborn P03.5</u> <u>- R.D.S. (newborn) P22.0</u> <u>- reaction - see also Conditions originating in the perinatal period, disorder</u> <u>- - drug NEC</u> <u>- - - newborn P93</u> <u>- - - withdrawal</u> <u>- - - - infant of dependent mother P96.1</u> <u>- - - - newborn P96.1</u> <u>- rectocele, maternal, in pregnancy or childbirth</u> <u>- - causing obstructed labor, affecting fetus or newborn P03.1</u> <u>- regurgitation, food, newborn P92.1</u> <u>- respiration, insufficient, or poor, newborn NEC P28.5</u> <u>- retardation</u> <u>- - growth, fetus P05.9</u> <u>- - - intrauterine P05.9</u> <u>- - physical, fetus P05.9</u> <u>- retraction, ring, uterus (Bandl's) (pathological), affecting fetus or newborn P03.6</u> <u>- retroversion, retroverted uterus, uterine, maternal, in pregnancy or childbirth</u> <u>- - causing obstructed labor, affecting fetus or newborn P03.1</u> <u>- Rh (factor)</u> <u>- - hemolytic disease (fetus or newborn) P55.0</u> <u>- - incompatibility, immunization or sensitization, fetus or newborn P55.0</u> <u>- - negative mother affecting fetus or newborn P55.0</u> <u>- rigid, rigidity</u> <u>- - cervix, maternal (uteri), in pregnancy or childbirth</u> <u>- - - affecting fetus or newborn P03.8</u> <u>- - - causing obstructed labor, affecting fetus or newborn P03.1</u> <u>- - pelvic floor, maternal, in pregnancy or childbirth</u> <u>- - - affecting fetus or newborn P03.8</u> <u>- - - causing obstructed labor, affecting fetus or newborn P03.1</u> <u>- - perineum or vulva, maternal, in pregnancy or childbirth</u> <u>- - - causing obstructed labor, affecting fetus or newborn P03.1</u> <u>- - vagina, maternal, in pregnancy or childbirth</u> <u>- - - causing obstructed labor, affecting fetus or newborn P03.8</u> <u>- ring(s)</u></p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>Bandl's, fetus or newborn P03.6</u> - - <u>contraction, complicating delivery, affecting fetus or newborn P03.6</u> - - <u>retraction, uterus, pathological, affecting fetus or newborn P03.6</u> - <u>rotation, manual, affecting fetus or newborn P03.8</u> - <u>rubella</u> - - <u>congenital P35.0</u> - - <u>maternal, affecting fetus or newborn P00.2</u> - - - <u>manifest rubella in infant P35.0</u> - <u>ruminant, newborn P92.1</u> - <u>rupture, ruptured - see also Conditions originating in the perinatal period, perforation</u> - - <u>fontanelle P13.1</u> - - <u>intestine (bowel) (colon) (ileum) (jejunum) (rectum) (sigmoid), fetus or newborn P78.0</u> - - <u>kidney, birth injury P15.8</u> - - <u>liver, birth injury P15.0</u> - - <u>marginal sinus (placental) (with hemorrhage), affecting fetus or newborn P02.1</u> - - - <u>with placenta previa, affecting fetus or newborn P02.0</u> - - <u>membranes</u> - - - <u>artificial, delayed delivery following</u> - - - - <u>affecting fetus or newborn P01.1</u> - - - <u>premature, affecting fetus or newborn P01.1</u> - - - <u>spontaneous, delayed delivery following</u> - - - - <u>affecting fetus or newborn P01.1</u> - - <u>spinal cord, fetus or newborn (birth injury) P11.5</u> - - <u>spleen (traumatic)</u> - - - <u>birth injury P15.1</u> - - - <u>congenital P15.1</u> - - <u>umbilical cord, complicating delivery, affecting fetus or newborn P50.1</u> - - <u>uterus, maternal</u> - - - <u>during or after labor, affecting fetus or newborn P03.8</u> - <u>salpingo-oophoritis, maternal (purulent) (ruptured) (septic) (suppurative)</u> - - <u>complicating pregnancy, affecting fetus or newborn P00.8</u> - <u>scar, scarring,</u> - - <u>cervix, maternal</u> - - - <u>in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - - <u>due to previous cesarean section, complicating pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - <u>uterus, maternal</u> - - - <u>in pregnancy or childbirth, affecting fetus or newborn P03.8</u> - <u>scleredema, newborn P83.0</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - <u>sclerema (newborn) P83.0</u> - - <u>adiposum P83.0</u> - - <u>edematosum P83.0</u> - <u>scleroderma, sclerodermia (diffuse) (generalized), newborn P83.8</u> - <u>scoliotic maternal pelvis with disproportion (fetopelvic)</u> - - <u>affecting fetus or newborn P03.1</u> - <u>section, cesarean</u> - - <u>affecting fetus or newborn P03.4</u> - - - <u>postmortem P01.6</u> - - - <u>previous, in pregnancy or childbirth P03.8</u> - <u>seizure(s), newborn P90</u> - <u>sepsis (generalized) (see also Conditions originating in the perinatal period, septicemia)</u> - - <u>bacterial, newborn P36.9</u> - - - <u>due to</u> - - - - <u>anaerobes NEC P36.5</u> - - - - <u>Escherichia coli P36.4</u> - - - - <u>Staphylococcus NEC P36.3</u> - - - - - <u>aureus P36.2</u> - - - - <u>streptococcus NEC P36.1</u> - - - - - <u>group B P36.0</u> - - - <u>specified type NEC P36.8</u> - - <u>newborn NEC P36.9</u> - - - <u>due to</u> - - - - <u>anaerobes NEC P36.5</u> - - - - <u>Escherichia coli P36.4</u> - - - - <u>Staphylococcus NEC P36.3</u> - - - - - <u>aureus P36.2</u> - - - - <u>streptococcus NEC P36.1</u> - - - - - <u>group B P36.0</u> - - - <u>specified NEC P36.8</u> - - <u>umbilical (newborn) (organism unspecified) P38</u> - <u>septic umbilical cord P38</u> - <u>septicemia, septicemic (generalized) (suppurative) (see also Conditions originating in the perinatal period, sepsis)</u> - - <u>Actinobacter, newborn P36.8</u> - - <u>Bacillus coli, newborn P36.8</u> - - <u>Citobacter, newborn P36.8</u> - - <u>Enterobacter (aerogenes) (clocae), newborn P36.8</u> - - <u>Friedlanders', newborn P36.8</u> - - <u>newborn NEC (see also Conditions originating in the perinatal period, sepsis,</u> 			
--	--	--	--	--

	<p><u>newborn) P36.9</u> - - <u>Streptococcus, streptococcal, neonatal P36.1</u> - <u>sextuplet, affecting fetus or newborn P01.5</u> - <u>shock (acute)</u> - - <u>birth, fetus or newborn NEC P96.8</u> - - <u>lung, newborn P22.0</u> - - <u>septic (see also Conditions originating in the perinatal period, septicemia)</u> - - - <u>newborn P36.9</u> - - <u>septicemic, newborn P36.9</u> - <u>short, shortening, shortness</u> - - <u>cord (umbilical), complicating delivery, affecting fetus or newborn P02.6</u> - <u>sinus, marginal, ruptured or bleeding, maternal, affecting fetus or newborn P02.1</u> - - <u>with placenta previa, affecting fetus or newborn P02.0</u> - <u>sleep apnea, newborn P28.3</u> - <u>slipped, slipping ligature, umbilical P51.8</u> - <u>slow</u> - - <u>feeding, newborn P92.2</u> - - <u>fetal growth NEC P05.9</u> - - <u>heart(beat)</u> - - - <u>fetal P20.9</u> - - - <u>newborn P29.1</u> - <u>small(ness)</u> - - <u>fetus or newborn for gestational age P05.1</u> - - <u>pelvis, maternal, with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>small-and-light-for-dates (infant) P05.1</u> - <u>small-for-dates (infant) P05.1</u> - <u>snuffles (non-syphilitic), newborn P28.8</u> - <u>spasm(s), spastic, spasticity (see also condition)</u> - - <u>cervix, maternal, complicating delivery, affecting fetus or newborn P03.6</u> - - <u>uterus, maternal, complicating labor, affecting fetus or newborn P03.6</u> - <u>spondylolisthesis, maternal (acquired)</u> - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>spondylolysis, maternal, lumbosacral region</u> - - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>spondylosis, maternal</u> - - <u>with disproportion (fetopelvic), affecting fetus or newborn P03.1</u> - <u>standstill respiration, respiratory, newborn P28.5</u> - <u>stenosis (cicatricial)</u> - - <u>cervix, cervical (canal), maternal, in pregnancy or childbirth</u></p>			
--	--	--	--	--

	<ul style="list-style-type: none"> - - - <u> affecting fetus or newborn P03.8</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u> vagina, maternal, in pregnancy or childbirth</u> - - - <u> affecting fetus or newborn P03.8</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - <u> stillbirth NEC P95</u> - <u> strangulation, strangled, umbilical cord, fetus or newborn P02.5</u> - - <u> with cord prolapse P02.4</u> - <u> stricture (see also Conditions originating in the perinatal period, stenosis)</u> - - <u> cervix, cervical (canal), maternal, in pregnancy or childbirth</u> - - - <u> affecting fetus or newborn P03.8</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - <u> stridor, congenital (larynx) NEC P28.8</u> - <u> stroke (apoplectic) (brain) (paralytic), hemorrhagic, newborn P52.4</u> - <u> surgery, previous, in pregnancy or childbirth</u> - - <u> cervix, maternal</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u> pelvic soft tissues NEC, maternal</u> - - - <u> affecting fetus or newborn P03.8</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u> uterus, maternal</u> - - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - <u> suspended maternal uterus, in pregnancy or childbirth</u> - - <u> causing obstructed labor, affecting fetus or newborn P03.1</u> - <u> syncope, bradycardia</u> - - <u> fetal P20.9</u> - - <u> newborn P29.1</u> - <u> syndrome - see also Conditions originating in the perinatal period, disease</u> - - <u> aspiration, of newborn (massive) P24.9</u> - - - <u> meconium P24.0</u> - - <u> bronze baby P83.8</u> - - <u> bubbly lung P27.0</u> - - <u> cardiorespiratory distress (idiopathic), newborn P22.0</u> - - <u> cold injury (newborn) P80.0</u> - - <u> congenital rubella (manifest) P35.0</u> - - <u> defibrination, fetus or newborn P60</u> - - <u> drug withdrawal, infant of dependent mother P96.1</u> - - <u> fetal transfusion P02.3</u> - - <u> fetomaternal dysfunctional P02.2</u> - - <u> floppy, baby P94.2</u> - - <u> gray (newborn) P93</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>idiopathic cardiorespiratory distress, newborn P22.0</u> - - <u>infant of diabetic mother P70.1</u> - - - <u>gestational diabetes P70.0</u> - - <u>inspissated bile (newborn) P59.1</u> - - <u>low cardiac output, newborn P29.8</u> - - <u>lower radicular, newborn (birth injury), P14.8</u> - - <u>meconium plug (newborn) P76.0</u> - - <u>perfusion, newborn P28.8</u> - - <u>placental</u> - - - <u>dysfunction, affecting fetus or newborn P02.2</u> - - - <u>insufficiency, affecting fetus or newborn P02.2</u> - - - <u>transfusion, in fetus or newborn P02.3</u> - - <u>pulmonary dysmaturity (Wilson-Mikity) P27.0</u> - - <u>radicular NEC, upper limbs, newborn (birth injury) P14.3</u> - - <u>respiratory distress (idiopathic) (newborn) P22.0</u> - - <u>transfusion, fetomaternal P50.4</u> - - <u>twin (to twin) transfusion, in fetus or newborn P02.3</u> - - <u>wet lung, newborn P22.1</u> - - <u>withdrawal, drug</u> - - - <u>infant of dependent mother P96.1</u> - - - <u>therapeutic use, newborn P96.2</u> - <u>syphilis, syphilitic, maternal (acquired), affecting fetus or newborn P00.2</u> - <u>complicating pregnancy or childbirth P00.2</u> - <u>tachycardia</u> - - <u>fetal - see Conditions originating in the perinatal period, distress, fetal</u> - - <u>newborn P29.1</u> - <u>tachypnea, transitory, of newborn P22.1</u> - <u>tear, torn (traumatic) - see also Conditions originating in the perinatal period,</u> <u>wound</u> - - <u>tentorial, at birth P10.4</u> - - <u>umbilical cord, affecting fetus or newborn P50.1</u> - <u>temperature, cold, trauma from, newborn P80.0</u> - <u>tetany (due to), neonatal (without calcium or magnesium deficiency) P71.3</u> - <u>threatened</u> - - <u>abortion, affecting fetus P01.8</u> - - <u>labor (see also Complications originating in the perinatal period, labor, false)</u> - - - <u>affecting fetus P01.8</u> - - <u>miscarriage, affecting fetus P01.8</u> - - <u>premature delivery, affecting fetus P01.8</u> - <u>thrombocytopenia, thrombocytopenic</u> - - <u>due to</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - - <u>exchange transfusion P61.0</u> - - - <u>idiopathic maternal thrombocytopenia P61.0</u> - - - <u>isoimmunization P61.0</u> - - <u>neonatal, transitory P61.0</u> - <u>thrombophlebitis, maternal</u> - - <u>antepartum (superficial), affecting fetus or newborn P00.3</u> - - <u>pregnancy (superficial), affecting fetus or newborn P00.3</u> - <u>thrombosis, thrombotic (multiple) (progressive) (septic) (vein) (vessel)</u> - - <u>umbilical cord (vessels), complicating delivery, affecting fetus or newborn P02.6</u> - <u>thyrotoxicosis, neonatal P72.1</u> - <u>tobacco (nicotine), maternal use, affecting fetus or newborn P04.2</u> - <u>torsion umbilical cord in fetus or newborn P02.5</u> - <u>torticollis (intermittent) (spastic), due to birth injury P15.2</u> - <u>toxemia, maternal (of pregnancy), affecting fetus or newborn P00.0</u> - <u>toxoplasma, toxoplasmosis</u> - - <u>congenital (acute) (chronic) (subacute) P37.1</u> - - <u>maternal, affecting fetus or newborn P00.2</u> - - - <u>manifest toxoplasmosis in infant or fetus (acute) (chronic) (subacute) P37.1</u> - <u>transfusion</u> - - <u>placental (syndrome) (mother), in fetus or newborn P02.3</u> - - <u>twin-to-twin, fetus or newborn P02.3</u> - <u>transverse</u> - - <u>arrest (deep), in labor, affecting fetus or newborn P03.1</u> - - <u>lie</u> - - - <u>before labor, affecting fetus or newborn P01.7</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - <u>trauma, traumatism (see also Conditions originating in the perinatal period, injury)</u> - - <u>birth - see Conditions originating in the perinatal period, birth, injury</u> - - <u>maternal, during pregnancy, affecting fetus or newborn P00.5</u> - <u>trigeminy</u> - - <u>fetal P20.9</u> - - <u>newborn P29.1</u> - <u>triplet, affecting fetus or newborn P01.5</u> - <u>tuberculosis, tubercular, tuberculous (caseous) (degeneration) (gangrene) (necrosis)</u> - - <u>congenital P37.0</u> - - <u>maternal, complicating pregnancy or childbirth, affecting fetus or newborn P00.2</u> - <u>tumor</u> 			
--	--	--	--	--

	<ul style="list-style-type: none"> - - <u>cervix, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>ovary, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - <u>pelvic, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>uterus (body), maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>vagina, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - - <u>vulva or perineum, maternal, in pregnancy or childbirth</u> - - - <u>affecting fetus or newborn P03.8</u> - - - <u>causing obstructed labor, affecting fetus or newborn P03.1</u> - <u>twin (pregnancy) (fetus or newborn) P01.5</u> - <u>twist, twisted umbilical cord in fetus or newborn P02.5</u> - <u>tyrosinemia, newborn, transitory P74.5</u> - <u>ulcer, ulcerated, ulcerating, ulceration, ulcerative</u> - - <u>intestine, intestinal, perforating, fetus or newborn P78.0</u> - - <u>peptic (site unspecified), newborn P78.8</u> - <u>underfeeding, newborn P92.3</u> - <u>underweight for gestational age P05.0</u> - <u>Underwood's disease P83.0</u> - <u>unstable lie, fetus or newborn, before labor P01.7</u> - <u>uremia, uremic (coma)</u> - - <u>congenital P96.0</u> - - <u>maternal NEC, affecting fetus or newborn P00.1</u> - - <u>newborn P96.0</u> - <u>urticaria neonatorum P83.8</u> - <u>use (of) harmful patent medicines, maternal, affecting fetus or newborn P04.1</u> - <u>uveitis (anterior), due to toxoplasmosis, congenital P37.1† H22.0*</u> - <u>vaccinia (generalized) (localized)</u> - - <u>congenital P35.8</u> - <u>vaginitis, maternal (acute), complicating pregnancy, affecting fetus or newborn P00.8</u> - <u>varicella, congenital P35.8</u> - <u>varicose vein (ruptured), umbilical cord, affecting fetus or newborn P02.6</u> - <u>varix (ruptured), umbilical cord, affecting fetus or newborn P02.6</u> 			
--	---	--	--	--

	<ul style="list-style-type: none"> - <u>vasa previa</u> - - <u>affecting fetus or newborn P02.6</u> - - <u>hemorrhage from, affecting fetus or newborn P50.0</u> - <u>ventilator lung, newborn P27.8</u> - <u>ventouse delivery NEC, affecting fetus or newborn P03.3</u> - <u>version, with extraction, affecting fetus or newborn P01.7</u> - <u>viremia, newborn P35.9</u> - <u>vitality, lack of, newborn P96.8</u> - <u>volvulus (bowel) (colon) (intestine), newborn K56.2</u> - <u>vomiting (see also Conditions originating in the perinatal period, hyperemesis)</u> - - <u>newborn P92.0</u> - <u>weak, weakness, newborn P96.8</u> - <u>weight</u> - - <u>999 grams or less at birth (extremely low) P07.0</u> - - <u>1000-2499 grams at birth (low) P07.1</u> - <u>wet lung, newborn P22.1</u> - <u>wide cranial sutures, newborn P96.3</u> - <u>Wilson-Mikity syndrome P27.0</u> - <u>withdrawal state, symptoms, syndrome, newborn</u> - - <u>correct therapeutic substance properly administered P96.2</u> - - <u>infant of dependent mother P96.1</u> - - <u>therapeutic substance, neonatal P96.2</u> - <u>wound (cut) (laceration) (open) (penetrating (puncture wound) (with penetrating foreign body)</u> - - <u>scalpel, fetus or newborn (birth injury) P15.8</u> 			
<p>Delete subterm and code</p> <p>Add subterm and code</p> <p>Add subterm and code</p>	<p>Cubitus</p> <p>- congenital Q68.1</p> <ul style="list-style-type: none"> - valgus (acquired) M21.0 - - <u>congenital Q68.8</u> - varus (acquired) M21.1 - - <u>congenital Q68.8</u> 	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
<p>Revise code</p> <p>Add modifier</p> <p>Add subterm and code</p>	<p>Cyst (colloid) (mucous) (retention) (simple)</p> <ul style="list-style-type: none"> - hydatid (<i>see also</i> Echinococcus) B67.9 - - Morgagni - - - female Q50.5 - - - male (<u>epididymal</u>) Q55.4 - - - - <u>testicular Q55.2</u> - mesonephric duct 	UK (URC:0017)	Major	January 2006

Revise code	- - female <u>Q50.5</u> - Morgagni (hydatid)			
Revise code	- - female <u>Q50.5</u>			
Add modifier	- - male (<u>epididymal</u>) Q55.4			
Add subterm and code	- - - <u>testicular Q55.2</u>			
Delete code	- paramesonephric duct <u>Q50.4</u>			
Add subterms and codes	- - female <u>Q50.4</u> - - male <u>Q55.2</u>			
Revise code	- wolffian - - female <u>Q50.5</u>			
Revise subterm & code	Deformity Q89.9 - finger (acquired) M20.0 - - congenital <u>NEC Q68.1</u> - mitral (leaflets) (valve) I05.8	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Delete subterm & code	- Ebstein's Q22.5			
Add subterm & code	- tricuspid (leaflets) (valve) I07.8 - - <u>Ebstein's Q22.5</u>			
Revise subterm and code	Diplegia (upper limbs) G83.0 - infantile or congenital (cerebral) (spastic) (spinal) <u>G80.8</u>	Australia (URC:0045)	Major	January 2006
Add subterm and code	- - <u>spastic G80.1</u>			
Add subterms and codes	Disease, diseased – <i>see also</i> Syndrome - viral, virus (<i>see also</i> Disease, by type of virus) B34.9 - - <u>Hanta (with renal manifestations) (Dobrava) (Puumala) (Seoul) A98.5 † N08.0*</u> - - - <u>with pulmonary manifestations (Andes) (Bayou) (Bermejo) (Black Creek Canal) (Choclo) (Jujuitiba) (Laguna negra) Lechiguanas) (New York) (Oran) (Sin Nombre) B33.4† J17.1*</u>	Brazil (URC:0042)	Major	January 2006
Add modifier	- - Hantaan (<u>Korean hemorrhagic fever</u>) A98.5† N08.0*			
Add subterm and codes	- - <u>Sin Nombre (Hantavirus (cardio)-pulmonary syndrome) B33.4† J17.1*</u>			
Add subterm and cross reference	Drug - <u>resistant bacterial agent in bacterial infection – see Resistance (to), antibiotic by bacterial agent</u>	Australia/NORDIC (URC:0026)	Major	January 2006

Revise code	Exposure (to) (<i>see also</i> Contact, with) T75.8 - pollution NEC Z58.5 - - soil <u>Z58.3</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Add subterm and codes	Fever R50.9 - Songo A98.5† N08.0*	Brazil (URC:0042)	Major	January 2006
Revise code	Hallux - malleus (acquired) NEC <u>M20.3</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Add modifier	Hantaan virus disease (<u>Korean hemorrhagic fever</u>) A98.5 † N08.0*	Brazil (URC:0042)	Major	January 2006
Add main term, subterm and codes	<u>Hantavirus disease (with renal manifestations) (Dobrava) (Puumala) (Seoul)</u> A98.5 † N08.0* - with <u>pulmonary manifestations (Andes) (Bayou) (Bermejo) (Black Creek Canal) (Choclo) (JQUITIBA) (Laguna negra) Lechiguanas) (New York) (Oran) (Sin Nombre) B33.4† J17.1*</u>	Brazil (URC:0042)	Major	January 2006
Delete modifiers and revise code Add subterm and code Delete subterm and code Modify subterm Add subterm and code	Hemiplegia G81.9 - congenital (cerebral) (spastic) (spinal) <u>G80.8</u> - - <u>spastic G80.2</u> - infantile (postnatal) G80.2 - spastic G81.1 - - congenital or infantile G80.2 - <u>spinal G81.1</u>	Australia (URC:0045)	Major	January 2006
Revise code Add modifier Add subterm and code	Hydatid - Morgagni's - - female <u>Q50.5</u> - - male (<u>epididymal</u>) Q55.4 - - - <u>testicular Q55.2</u>	UK (URC:0017)	Major	January 2006
Add subterm and code	Hyperactive, hyperactivity - <u>detrusor muscle N32.8</u>	North America (URC:0040)	Major	January 2006

Add subterm and code	Hypoplasia, hypoplastic - lung (lobe) (not associated with short gestation) Q33.6 - - <u>with immaturity, prematurity or low birth weight P28.0</u>	MRG (URC:0118)	Major	January 2006
Modify subterm Add subterm and code	Infection, infected (opportunistic) B99 - protozoal <u>NEC B64</u> - - <u>specified NEC B60.8</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Modify subterm Add subterms and codes Modify subterm Add subterms, codes and cross references	Infection, infected (opportunistic) B99 - with lymphangitis – <i>see</i> <u>Lymphangitis</u> - - <u>antibiotic-resistant bacterial agent (resistant to) U89.9</u> - - - <u>methicillin U80.1</u> - - - <u>multiple antibiotics U88</u> - - - <u>penicillin U80.0</u> - - - <u>penicillin-related antibiotic U80.8</u> - - - <u>specified antibiotic (single) NEC U89.8</u> - - - - <u>multiple antibiotics U88</u> - - - <u>vancomycin U81.0</u> - - - <u>vancomycin-related antibiotic U81.8</u> - - <u>lymphangitis</u> – <i>see</i> <u>Lymphangitis</u> - bacterial NEC A49.9 - - agent NEC, as cause of disease classified elsewhere B96.8 - - - <u>as cause of disease classified elsewhere B96.8</u> - - - <u>resistant to antibiotic – see Resistance (to), antibiotic(s) by bacterial agent</u>	Australia/ NORDIC (URC:0026)	Major	January 2006
Revise code Add subterms and codes	Infestation B88.9 - protozoal NEC <u>B64</u> - - <u>intestinal A07.9</u> - - - <u>specified NEC A07.8</u> - - <u>specified NEC B60.8</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Add subterm and code Add subterm and code	Injury (see also specified injury type) T14.9 - globe (eye) S05.9 - - <u>specified NEC S05.8</u> - vitreous S05.9 - - <u>specified NEC S05.8</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

Add subterms and cross references	Intoxication - maternal medication, via placenta... - <u>meaning</u> - - <u>inebriation – code to F10-F19 with fourth character .0</u> - - <u>poisoning – see Table of drugs and chemicals</u>	MRG (URC:0116)	Major	January 2006
Revise cross reference	Meconium - in liquor – <u>see also</u> Distress, fetal	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Revise code	Meningitis (basal) (cerebral) (spinal) G03.9 - poliovirus A80.9† <u>G02.0*</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Revise code	Meningoencephalitis (<u>see also</u> Encephalitis) G04.9 - parasitic NEC <u>B89†</u> G05.2*	Dutch Committee on Translation of ICD-10 (URC: 0082)	Major	January 2006
Modify subterm Add subterm and code	Monoplegia G83.3 - congenital or infantile (cerebral) (spinal) G80.8 - - <u>spastic G80.1</u>	Australia (URC:0045)	Major	January 2006
Revise code Add modifier Add subterm and code	Morgagni's - cyst, organ, hydatid or appendage - - female <u>Q50.5</u> - - male (<u>epididymal</u>) Q55.4 - - - <u>testicular Q55.2</u>	UK (URC:0017)	Major	January 2006
Modify cross reference Delete subterm and code	Myolipoma (M8860/0) – <u>see also</u> Lipoma – unspecified site D30.0	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Add modifier	Nephropathia epidemica (<u>Puumala hantavirus</u>) A98.5 † N08.0*	Brazil (URC:0042)	Major	January 2006

Revise code	Neuritis M79.2 - syphilitic <u>A52.1</u> † G59.8*	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Revise code Add modifier Add subterm and code	Organ of Morgagni (persistence of) - female <u>Q50.5</u> - male (<u>epididymal</u>) Q55.4 - - <u>testicular Q55.2</u>	UK (URC:0017)	Major	January 2006
Add subterm and code	Overactive - bladder <u>N32.8</u>	North America (URC:0040)	Major	January 2006
Delete modifier Add subterm and code Delete modifier and revise code Add subterm and code Add subterms and codes Modify code Add subterm and code Modify subterm Add subterm and code Modify subterm Modify subterm Add subterm and code Revise code Add subterms	Palsy (see also Paralysis) G83.9 - cerebral (congenital) (infantile) G80.9 - - <u>choreathetoid G80.3</u> - - diplegic (spastic) <u>G80.8</u> - - - <u>spastic G80.1</u> - - dyskinetic G80.3 - - - <u>athetoid G80.3</u> - - - <u>choreathetoid G80.3</u> - - - <u>dystonic G80.3</u> - - <u>dystonic G80.3</u> - - hemiplegic <u>G80.8</u> - - - <u>spastic G80.2</u> - - monoplegic NEC G80.8 - - - <u>spastic G80.1</u> - - not congenital, or infantile acute I64 - - paraplegic NEC G80.8 - - quadriplegic G80.8 - - - <u>spastic G80.0</u> - - spastic <u>G80.1</u> - - - <u>diplegic G80.1</u> - - - <u>hemiplegic G80.2</u>	Australia (URC:0045)	Major	January 2006

and code	<ul style="list-style-type: none"> - - - <u>monoplegic G80.1</u> - - - <u>quadriplegic G80.0</u> - - - <u>specified NEC G80.1</u> - - - <u>tetraplegic G80.0</u> - - tetraplegic G80.8 - - - <u>spastic G80.0</u> 			
Add subterm and code				
Modify subterm, delete code and add cross reference Delete subterm and code Delete code and add cross reference Delete code and add cross reference Delete modifier and code and add cross reference Delete code and add cross reference Delete code and add cross reference	<p>Paralysis, paralytic (complete) (incomplete) (<i>see also</i> Paresis) G83.9</p> <ul style="list-style-type: none"> - congenital (cerebral) (spinal) G80.9 – <u><i>see</i> Palsy, cerebral</u> —spastic G80.0 - infantile (<i>see also</i> Poliomyelitis, paralytic) A80.3 - - spastic G80.0 – <u><i>see</i> Palsy, cerebral, spastic</u> - spastic G83.9 - - cerebral infantile G80.0 – <u><i>see</i> Palsy, cerebral, spastic</u> - - congenital (cerebral) (spinal) G80.2 – <u><i>see</i> Palsy, cerebral, spastic</u> - spinal (cord) G83.8 - - congenital NEC G80.9 – <u><i>see</i> Palsy, cerebral</u> - - spastic NEC G80.0 – <u><i>see</i> Palsy, cerebral, spastic</u> 	Australia (URC:0045)	Major	January 2006
Modify subterm	<p>Paraplegia (lower) G82.2</p> <ul style="list-style-type: none"> - congenital or infantile (cerebral) (spinal) G80.8 	Australia (URC:0045)	Major	January 2006
Add dagger and asterisk code Add dagger and asterisk code	<p>Pinta A67.9</p> <ul style="list-style-type: none"> - cardiovascular lesions A67.2† I98.1* - lesions - - cardiovascular A67.2† I98.1* 	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

<p>Add subterm & code Delete subterms & codes</p>	<p>Pregnancy (single) (uterine) - complicated by (<i>see also</i> Pregnancy, management, affected by) - - malformation - - - <u>placenta, placental (vessel) O43.1</u> — specified NEC O26.8 — placenta, placental (vessel) O43.1</p>	<p>Dutch Committee on Translation of ICD-10 (URC: 0082)</p>	<p>Minor</p>	<p>January 2004</p>
<p>Modify subterm Add subterm and code Add subterms and codes Add subterm and code Add subterm and code</p>	<p>Quadriplegia (chronic) G82.5 - congenital or infantile (cerebral) (spinal) G80.8 - - <u>spastic G80.0</u> - flaccid G82.3 - - <u>congenital (cerebral) G80.8</u> - - - <u>spastic G80.0</u> - spastic G82.4 - - <u>congenital (cerebral) G80.0</u> - <u>spinal G82.4</u></p>	<p>Australia (URC:0045)</p>	<p>Major</p>	<p>January 2006</p>
<p>Add main term, subterms and codes</p>	<p><u>Resistance, resistant (to)</u> - <u>antibiotic(s), by bacterial agent</u> - - <u>methicillin U80.1</u> - - <u>multiple U88</u> - - <u>penicillin U80.0</u> - - <u>penicillin-related U80.8</u> - - <u>specified (single) NEC U89.8</u> - - - <u>multiple U88</u> - - <u>vancomycin U81.0</u> - - <u>vancomycin-related U81.8</u></p>	<p>Australia/NORDIC (URC:0026)</p>	<p>Major</p>	<p>January 2006</p>
<p>Add modifier, revise morphology code and code Delete subterms</p>	<p><u>Reticulosarcoma (diffuse) (M9591/3) C83.3</u> — diffuse C83.0</p>	<p>Australia (URC:0027)</p>	<p>Major</p>	<p>January 2006</p>

and codes	pleomorphic cell type (M9593/3) C83.9			
-----------	--	--	--	--

Add subterms and codes	Sequelae (of) – <i>see also condition</i> - <u>childbirth complication O94</u> - - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u> - complication(s) of	Australia (URC:0112)	Major	January 2006
Add subterms and codes	- - <u>childbirth (delivery), pregnancy or puerperium O94</u> - - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u>			
Add subterms and codes	- <u>delivery complication O94</u> - - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u>			
Revise code	- obstetric cause O94			
Add subterms and codes	- - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u> - <u>pregnancy complication(s) O94</u> - - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u>			
Add subterms and codes	- <u>puerperium complication(s) O94</u> - - <u>resulting in death (one year or more after delivery) O97</u> - - - <u>between 42 days and one year after delivery O96</u>			
Add main term and codes	<u>Sin Nombre virus disease (Hantavirus (cardio)-pulmonary syndrome) B33.4† J17.1*</u>	Brazil (URC:0042)	Major	January 2006
Revise code	Spastic, spasticity – <i>see also Spasm</i> - child (cerebral) (congenital) (paralysis) <u>G80.1</u>	Australia (URC:0045)	Major	January 2006
Add subterm and codes	Syndrome – <i>see also Disease</i> - <u>Hantavirus (cardio)-pulmonary (HPS) (HCPS) B33.4† J17.1*</u>	Brazil (URC:0042)	Major	January 2006
Revise code	Tabacism, tabacosis, tabagism <u>T65.2</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

Delete subterms and codes	Tetraplegia (<i>see also</i> Quadriplegia) G82.5 flaccid G82.3 spastic G82.4	Australia (URC:0045)	Major	January 2006
---------------------------	--	----------------------	-------	--------------

Revise code	Use (of) - harmful F1x.1 - - patent medicines F55 - - - maternal, affecting fetus or newborn <u>P04.1</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004
Delete subterm Revise subterm	Wound, open... T14.1 - multiple T01.9 —sites - - specified <u>sites</u> NEC T01.8	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

Index to External Causes

Instruction	Alphabetic index entries	Source	Major/Minor update	Implementation date
Revise code	Bite, bitten by - arthropod (nonvenomous) NEC W57.- - - venomous (<i>see also</i> Contact, with, by type of arthropod) <u>X25.-</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Major	January 2006
Revise code	Cataclysm, cataclysmic (any injury) NEC X39.- - earthquake <u>X34.-</u>	Dutch Committee on Translation of ICD-10 (URC: 0082)	Major	January 2006
	Pushed, pushing (accidental) (injury in) (overextension) X50.- - by other person(s) (accidental) W51.- - - from - - - transport vehicle NEC (<i>see also</i> Accident, transport) V89.9 - - - - stated as	Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004

Ratified by HoC/WHO at the HoC meeting in Brisbane, October 2002

Add dash to code	- - - - - intentional, homicide (attempt) <u>Y08.-</u>			
------------------	--	--	--	--

Table of Drugs and Chemicals

Instruction	Alphabetic index entries					Source	Major/Minor update	Implementation date
	Brass (fumes)	<u>T56.8</u>	X49.-	X69.-	Y19.-			
Revise Chapter XIX code						Dutch Committee on Translation of ICD-10 (URC: 0082)	Minor	January 2004