
CERVICAL CANCER PROFILE

TOTAL POPULATION, FEMALE (2019): 1 183 000

TOTAL DEATHS. FEMALE (2019): 6 300


Morbidity and Mortality

Crude cervical cancer incidence per 100 000 women (2020):	23.5
Age-standardized cervical cancer incidence	
per 100 000 women (2020):	42.9
Cumulative risk of cervical cancer, ages 0-74 (2020):	3.9%
Cervical cancer deaths (2019):	140
Cervical cancer mortality-to-incidence ratio (2020):	0.7
Population-based cancer registry exists (2021):	Yes



Primary Prevention

HPV vaccination programme coverage among girls (2020)



Among girls turning 15 years old in 2020, **7 in 10** received their first HPV vaccination dose

HPV vaccination programme (2020):

HPV included in national vaccination programme:	Yes
Scale of vaccination programme:	National
Year of introduction:	2019
Primary target cohort:	9-14 years


Related risk factors:

Tobacco use prevalence, women aged 15+ years (2020):	1%
Condom use at last high-risk sex (2013):	57%
HIV incidence per 1000, women aged 15+ years (2020):	1.54

Secondary Prevention

National screening programme	
for cervical cancer exists (2021):	No
Primary screening test used (2021):	-
Target age range of programme (2021):	-
Programme/guidelines exist to strengthen early detection	
of first symptoms at primary health care level (2021):	No
Clearly defined referral system exists from primary care	
to secondary and tertiary care (2021):	Yes

Screening for cervical cancer (2019)



Fewer than 1 in 10 women have been screened for cervical cancer in the last 5 years

Treatment and Supportive Care

National guidelines on cervical cancer management	
exist (2021):	No

Number of radiotherapy units per 10 000 cancer patients (-): ND Number of brachytherapy units per 10 000 cancer patients (-): ND

Cancer diagnosis and treatment services generally available (2021):

No
Yes
Yes
Yes
No

Number of medical staff (per 10 000 cancer patients):

Radiation oncologists (2019):	ND
Medical physicists (2019):	ND
Surgeons (2011):	142
Radiologists (2019):	0
Nuclear medicine physicians (2019):	0

Palliative care for patients with NCDs in the public health system generally available (2021):

In primary health care facilities:	No
In community or home-based care:	No

Reported annual opioid consumption excluding methadone - in oral morphine equivalence per capita (2017):

WHO Cervical Cancer Elimination Strategy Targets for 2030

90% of girls fully vaccinated with the HPV vaccine by the age of 15

70% of women are screened with a high-performance test by 35 years of age and again by 45 years of age

90% of women identified with

ND = data not available

* Programme coverage: % of national target population (among 9-14-year-old girls) † Coverage by age 15: % of population turning 15 that have been vaccinated against HPV at any time between ages 9 to 14

See Explanatory Notes for indicator descriptions.

cervical disease receive treatment