

Botswana

Hypertension profile


Total population (2019): 2 500 000

Total deaths (2019): 20 700


Age-standardized prevalence of hypertension among adults aged 30–79 years (2019)^a

♀ 44% ♂ 40% ♀ 47%


Prevalence of hypertension – global comparison (both sexes)^a




Of the 364 000 adults aged 30–79 years with hypertension:



Trends in uncontrolled hypertension in adults aged 30–79 years^c



Hypertension control rate scenarios



Mortality

	both sexes	males	females	year
Probability of premature mortality from NCDs (%)	27	33	23	2019
Cardiovascular disease deaths	4100	2000	2100	2019
Cardiovascular disease deaths attributable to high systolic blood pressure (%)	63	64	61	2019

Risk factors^e

	both sexes	males	females	year
Mean population salt intake, adults aged 25+ years (g/day)	8	8	7	2019
Current tobacco use, adults aged 15+ years (%)	20	31	9	2019
Obesity, adults aged 18+ years (%)	19	8	29	2016
Total alcohol per capita consumption, adults aged 15+ years (litres)	8	14	3	2019
Physical inactivity, adults aged 18+ years (%)	22	17	26	2016

National response

Targets

National target for blood pressure
National target for salt consumption

Surveillance

Conducted recent, national survey measuring raised blood pressure/hypertension
Conducted recent, national survey on salt/sodium intake
Functioning system for generating reliable cause-specific mortality data on a routine basis

Treatment

Guidelines for management of hypertension

Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake.

See Explanatory notes for description of indicators