# **Switzerland**

## Hypertension profile

Age-standardized prevalence of hypertension among adults aged 30-79 years (2019)<sup>a</sup>

Total population (2019): 8 576 000


Total deaths (2019): 69 100














Of the **1.5 million** adults aged 30–79 years with hypertension:



### Trends in uncontrolled hypertension in adults aged 30-79 years<sup>c</sup>



### Hypertension control rate scenarios



Projected hypertension control rates by scenario:d

Mortality						
	both sexes	males	females	year		
Probability of premature mortality from NCDs (%)	8	10	6	2019		
Cardiovascular disease deaths	21 300	9700	11 600	2019		
Cardiovascular disease deaths attributable to high systolic blood pressure (%)	46	47	46	2019		

	both sexes	males	females	year
Mean population salt intake, adults aged 25+ years (g/day)	8	9	7	2019
Current tobacco use, adults aged 15+ years (%) <sup>f</sup>		28	23	2019
Obesity, adults aged 18+ years (%)	20	22	17	2016
Total alcohol per capita consumption, adults aged 15+ years (litres)		16	5	2019
Physical inactivity, adults aged 18+ years (%)	24	22	26	2016

### **National response**

Risk factors

Targets
National target for blood pressure
National target for salt consumption
Surveillance

Conducted recent, national survey measuring raised blood pressure/hypertension Conducted recent, national survey on salt/sodium intake

Functioning system for generating reliable cause-specific mortality data on a routine basis

# business as usual ---- progress ---- aspirational

Treatment		
22	26	2016
10	ວ	2019

Guidelines for management of hypertension



Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake. f. Data refer to tobacco smoking only, in the absence of sufficient data on all tobacco use.

See Explanatory notes for description of indicators