Czechia

Hypertension profile


Age-standardized prevalence of hypertension among adults aged 30-79 years (2019)^a

Total population (2019): 10 537 000 Total deaths (2019): 113 000

> 42% 49%

Q 34%







Of the **3.3 million** adults aged 30–79 years with hypertension:



Trends in uncontrolled hypertension in adults aged 30-79 years^c



Hypertension control rate scenarios



Projected hypertension control rates by scenario:d business as usual ---- progress ---- aspirational

Mortality				
	both sexes	males	females	year
Probability of premature mortality from NCDs (%)	14	19	10	2019
Cardiovascular disease deaths	47 900	22 600	25 400	2019
Cardiovascular disease deaths attributable to high systolic blood pressure (%)	47	51	43	2019

	both sexes	males	females	year
Mean population salt intake, adults aged 25+ years (g/day)	13	15	11	2019
Current tobacco use, adults aged 15+ years (%) ^f	31	35	27	2019
Obesity, adults aged 18+ years (%)	26	26	25	2016
Total alcohol per capita consumption, adults aged 15+ years (litres)	13	21	6	2019
Physical inactivity, adults aged 18+ years (%)	31	28	34	2016

Risk factors^e

National response			
Targets National target for blood pressure National target for salt consumption	×	Treatment Guidelines for management of hypertension	~
Surveillance Conducted recent, national survey measuring raised blood pressure/hypertension	~	пуренензіон	
Conducted recent, national survey measuring raised blood pressure/hypertension Conducted recent, national survey on salt/sodium intake Functioning system for generating reliable cause-specific mortality data on a routine basis	×		

Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake. f. Data refer to tobacco smoking only, in the absence of sufficient data on all tobacco use.