Germany

Hypertension profile

Total population (2019): 83 148 000

Total deaths (2019): 845 000

Age-standardized prevalence of hypertension among adults aged 30-79 years (2019)^a




ೆ 34%








Of the **19.8 million** adults aged 30–79 years with hypertension:




Trends in uncontrolled hypertension in adults aged 30–79 years^c



Hypertension control rate scenarios

both sexes



Projected hypertension control rates by scenario:^d business as usual •••• progress •••• aspirational •••

Probability of premature mortality from NCDs (%)

Cardiovascular disease deaths

Cardiovascular disease deaths attributable to high systolic blood pressure (%)

both sexes	males	females	year
12	15	9	2019
285 000	158 000	127 000	2019
55	52	57	2019

males

10

Risk factors^e

Mortality

Mean population salt intake, adults aged 25+ years (g/day)	9
Current tobacco use, adults aged 15+ years (%) ^f	23
Obesity, adults aged 18+ years (%)	22
Total alcohol per capita consumption, adults aged 15+ years (litres)	12
Physical inactivity, adults aged 18+ years (%)	42

3	25	20	2019
2	24	20	2016
2	19	5	2019
2	40	44	2016

females

National response

i ai Bets	
National target for blood pressure	
National target for salt consumption	

Surveillance

Targets

Conducted recent, national survey measuring raised blood pressure/hypertension Conducted recent, national survey on salt/sodium intake

Functioning system for generating reliable cause-specific mortality data on a routine basis

TreatmentGuidelines for management of hypertension



vear

2019

Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake. f. Data refer to tobacco smoking only, in the absence of sufficient data on all tobacco use.

See Explanatory notes for description of indicators