Denmark

Hypertension profile

Age-standardized prevalence of hypertension among adults aged 30-79 years (2019)^a

236%

3 43%

Total population (2019): 5 796 000


Total deaths (2019): 54 000

Q 29%







Of the **1.5 million** adults aged 30–79 years with hypertension:



Trends in uncontrolled hypertension in adults aged 30–79 years^c



Hypertension control rate scenarios



Mortauty				
	both sexes	males	females	year
Probability of premature mortality from NCDs (%)	11	13	9	2019
Cardiovascular disease deaths	12 800	6600	6200	2019
Cardiovascular disease deaths attributable to high systolic blood pressure (%)	52	53	50	2019
Risk factors ^e				

	both sexes	males	females	year
Mean population salt intake, adults aged 25+ years (g/day)	8	9	7	2019
Current tobacco use, adults aged 15+ years (%) ^f	18	18	18	2019
Obesity, adults aged 18+ years (%)	20	22	17	2016
Total alcohol per capita consumption, adults aged 15+ years (litres)	9	15	4	2019
Physical inactivity, adults aged 18+ years (%)	29	26	31	2016

National response			
Targets		Treatment	
National target for blood pressure	×	Guidelines for management of	
National target for salt consumption	×	hypertension	~
Surveillance			
Conducted recent, national survey measuring raised blood pressure/hypertension	~		
Conducted recent, national survey on salt/sodium intake	×		
Functioning system for generating reliable cause-specific mortality data on a routine basis	V		

Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake. f. Data refer to tobacco smoking only, in the absence of sufficient data on all tobacco use.