Japan

Hypertension profile

Total population (2019): 125 800 000

Total deaths (2019): 1352 000

Age-standardized prevalence of hypertension among adults aged 30-79 years (2019)^a




් 40%









Of the **33.1 million** adults aged 30–79 years with hypertension:



Trends in uncontrolled hypertension in adults aged 30–79 years^c



Hypertension control rate scenarios



Projected hypertension control rates by scenario:^d business as usual ---- progress ---- aspirational -

Probability of premature mortality from NCDs (%)

Cardiovascular disease deaths

Cardiovascular disease deaths attributable to high systolic blood pressure (%)

both sexes	males	females	year
8	11	6	2019
364 000	165 000	199 000	2019
45	48	42	2019

both sexes males females year Mean population salt intake, adults aged 25+ years (g/day) 10 9 11 2019 Current tobacco use, adults aged 15+ years (%)f 10 2019 21 31 Obesity, adults aged 18+ years (%) 4 5 4 2016 Total alcohol per capita consumption, adults aged 15+ years (litres) 7 11 3 2019 Physical inactivity, adults aged 18+ years (%) 35 34 37 2016

National response

Targets

Mortality

Risk factorse

National target for blood pressure National target for salt consumption

Surveillance

Conducted recent, national survey measuring raised blood pressure/hypertension Conducted recent, national survey on salt/sodium intake

Functioning system for generating reliable cause-specific mortality data on a routine basis

Treatment

Guidelines for management of hypertension

no data

Footnotes: a. SBP ≥140 mmHg or DBP ≥90 mmHg or taking medication for hypertension. b. Control rate: adults aged 30–79 years receiving treatment, with blood pressure SBP <140 mmHg and DBP <90 mmHg. c. SBP ≥140 mmHg or DBP ≥90 mmHg. d. Progress and aspirational scenarios reflect a theoretical scaling up of treatment and control. e. Age-standardized estimates are presented for all indicators except salt intake. f. Data refer to tobacco smoking only, in the absence of sufficient data on all tobacco use.

See Explanatory notes for description of indicators