




		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY†							
	Premature mortality from NCDs	✓	Total NCD deaths	2016
			Risk of premature death between 30-70 years (%)	2016
	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	...
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	14	3	8
	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	16	34	25
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	7	7	7
	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	26	21	23
	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	9	14	11
	Obesity	✓	Obesity, adults aged 18+ (%)	2016	20	36	28
			Obesity, adolescents aged 10-19 (%)	2016	13	14	13
	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	2
	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	9

SELECTED ADULT RISK FACTOR TRENDS



NATIONAL SYSTEMS RESPONSE							
	Drug therapy to prevent heart attacks and strokes	✓	Proportion of population at high risk for CVD or with existing CVD (%)	-	...		
			Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%)	-	...		
			Proportion of primary health care centres reported as offering CVD risk stratification	2017		None	
			Reported having CVD guidelines that are utilized in at least 50% of health facilities	2017		Yes	
	Essential NCD medicines and basic technologies to treat major NCDs	✓	Number of essential NCD medicines reported as "generally available"	2017		9 out of 10	
			Number of essential NCD technologies reported as "generally available"	2017		6 out of 6	

... = no data available
† See Explanatory Notes