


NO DATA AVAILABLE


NO DATA AVAILABLE

		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY [†]							
	Premature mortality	X	Total NCD deaths	2016			
	from NCDs	A	Risk of premature death between 30-70 years (%)	2016			
<u>F</u>	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	
RISK	FACTORS						
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016			
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	33	57	45
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010			
\odot	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	23	8	16
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	25	19	22
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	25	21	23
			Obesity, adults aged 18+ (%)	2016	51	58	55
	Obesity	✓	Obesity, adolescents aged 10-19 (%)	2016	30	29	29
%	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	1
备	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	13
CELEC	TED ADULT BISK FACT	OR TREMPS					















NATIONAL SYSTEMS RESPONSE













Essential NCD medicines and basic technologies to treat major NCDs

Number of essential NCD technologies reported as "generally available"

Number of essential NCD medicines reported as

2017 2017

10 out of 10 6 out of 6