


Burkina Faso


Total population (2016): 18 633 725 ► Population aged 15 years and older (15+): 55% ► Population in urban areas: 31% ► Income group (World Bank): Low income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	4.6		4.9	
Unrecorded	2.5		3.3	
Total**	7.1		8.2	
Total males / females	12.2	2.3	14.2	2.4
WHO African Region	6.3		6.3	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	27.7
Females (15+)	10.5
Both sexes (15+)	22.2

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	29.4	57.5	21.1	59.9
Females	5.2	22.5	3.2	24.3
Both sexes	17.1	46.3	12.3	50.4

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	47.2	69.3	58.5
Former drinkers* (15+)	1.7	7.5	4.7
Abstainers (15+), past 12 months	48.8	76.8	63.1

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	43.8	38.5	66.9	42.3	1322
Road traffic injuries, males / females	75.9	36.4	30.9	22.2	1660
Cancer, males / females	121.8	125.3	7.4	2.4	324

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 3 4 **5** > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	3.4	1.4
Females	0.3	0.0
Both sexes	1.8	0.7
WHO African Region	3.7	1.3

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	No / No / No
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	13 / 13 / 13
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / Yes, No No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.05 / 0.05
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	No