


Bosnia and Herzegovina


Total population (2016): 3 802 134 ► Population aged 15 years and older (15+): 87% ► Population in urban areas: 37% ► Income group (World Bank): Upper-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	4.6		4.4	
Unrecorded	2.5		1.8	
Total**	7.3		6.4	
Total males / females	12.3	2.5	10.9	2.1
WHO European Region	11.2		9.8	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	17.5
Females (15+)	6.1
Both sexes (15+)	13.4

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	33.4	53.7	29.7	63.9
Females	6.9	20.4	5.9	27.5
Both sexes	20.0	41.8	18.2	52.9

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	15.5	39.2	27.4
Former drinkers* (15+)	22.4	26.8	24.6
Abstainers (15+), past 12 months	37.9	66.0	52.1

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	21.7	5.9	68.6	43.0	358
Road traffic injuries, males / females	26.9	5.7	33.9	22.1	175
Cancer, males / females	207.5	115.3	6.9	2.7	380

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 **3** 4 5 > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	9.8	5.5
Females	2.2	1.4
Both sexes	6.0	3.4
WHO European Region	8.8	3.7

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Subnational / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions on on-/off-premise sales of alcoholic beverages (any):	
Hours, days / places, density	No, No / No, No
Specific events / intoxicated persons / petrol stations	Yes / Yes / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.03 / 0.00 / 0.03
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes