
Cameroon


Total population (2016): 23 924 407 > Population aged 15 years and older (15+): 58% > Population in urban areas: 54% > Income group (World Bank): Lower-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	6.1		6.5	
Unrecorded	2.6		2.4	
Total**	8.7		8.9	
Total males / females	14.7	2.9	15.2	2.7
WHO African Region	6.3		6.3	

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine

Spirits Other

	Litres
Males (15+)	28.2
Females (15+)	10.6
Both sexes (15+)	22.6

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	31.5	58.5	23.2	61.7
Females	5.9	23.7	3.7	25.7
Both sexes	18.7	47.4	13.5	51.8

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	44.2	67.0	55.7
Former drinkers* (15+)	2.0	7.9	5.0
Abstainers (15+), past 12 months	46.2	75.0	60.6

 $^{^{\}star}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

(AAI), 2010					
	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	76.4	56.9	67.4	43.7	3639
Road traffic injuries, males / females	63.4	21.9	34.2	25.9	2186
Cancer, males / females	122.9	116.7	6.8	2.3	517

*Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016 LEAST < 1 2 3 4 6 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	11.8	4.0
Females	2.3	0.9
Both sexes	7.0	2.5
WHO African Region	3.7	1.3

 $^{^{\}star}$ 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	21/21/21
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, No / Yes, No No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.08 / 0.08 / 0.08
Legally binding regulations on alcohol advertising / product placement (any)	No / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	No
National monitoring system(s) (any)	No

^{*} Based on alcohol-attributable years of life lost.