


Spain


Total population (2016): 46 064 604 ► Population aged 15 years and older (15+): 85% ► Population in urban areas: 81% ► Income group (World Bank): High income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	9.6		8.5	
Unrecorded	1.2		1.8	
Total**	10.5		10.0	
Total males / females	16.9	4.4	16.4	4.0
WHO European Region	11.2		9.8	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	20.2
Females (15+)	7.0
Both sexes (15+)	14.6

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	41.3	51.1	43.9	62.6
Females	10.7	18.8	11.2	26.4
Both sexes	25.6	37.3	28.0	49.5

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	6.6	20.1	13.6
Former drinkers* (15+)	12.5	22.9	17.9
Abstainers (15+), past 12 months	19.1	43.0	31.4

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	12.8	4.1	73.8	56.3	3 621
Road traffic injuries, males / females	5.6	1.6	39.6	28.3	590
Cancer, males / females	200.7	100.1	8.3	3.9	7 264

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 **2** 3 4 5 > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	2.7	1.2
Females	0.5	0.2
Both sexes	1.5	0.7
WHO European Region	8.8	3.7

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (1999/2009) / Yes
Excise tax on beer / wine / spirits	Yes / No / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions on on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density	Yes, No / Yes, No
Specific events / intoxicated persons / petrol stations	Yes / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.03 / 0.03
Legally binding regulations on alcohol advertising / product placement (any)	Yes / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	Yes / Yes
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes