
United Kingdom of Great Britain and Northern Ireland


Total population (2016): 65 111 143 ➤ Population aged 15 years and older (15+): 82% ➤ Population in urban areas: 84% ➤ Income group (World Bank): High income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961-2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	10.1		9.8	
Unrecorded	1.6		1.1	
Total**	12.3		11.4	
Total males / females	19.6	5.4	18.4	4.8
WHO European Region	11.2		9.8	

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine Spirits Other

	Litres
Males (15+)	21.8
Females (15+)	7.6
Both sexes (15+)	15.6

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	46.4	55.1	49.9	66.4
Females	13.8	22.0	14.5	29.7
Both sexes	29.8	40.6	32.6	52.4

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	5.3	16.4	11.0
Former drinkers* (15+)	10.4	20.6	15.6
Abstainers (15+), past 12 months	15.7	37.0	26.6

 $^{^{\}star}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), $2016\,$

	ASI	OR*	AAF	(%)	AAD** (Number)
Liver cirrhosis, males / females	14.7	7.9	77.0	63.4	5 556
Road traffic injuries, males / females	4.8	1.4	43.8	32.2	735
Cancer, males / females	189.7	143.1	8.0	4.1	10 412

^{*}Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016 LEAST < 1 ② 3 4 5 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	13.0	2.2
Females	4.7	0.7
Both sexes	8.7	1.4
WHO European Region	8.8	3.7

 $^{^{\}star}$ 12-month prevalence estimates (15+); ** including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Subnational / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	sub-national sub-national

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	Subnational
Legally binding regulations on alcohol advertising / product placement (any)	No / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / Yes
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes

^{*} Based on alcohol-attributable years of life lost.