


Haiti


Total population (2016): 10 848 175 ► Population aged 15 years and older (15+): 67% ► Population in urban areas: 60% ► Income group (World Bank): Low income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	5.8		5.8	
Unrecorded	0.6		0.0	
Total**	6.4		5.8	
Total males / females	10.9	2.1	10.0	1.8
WHO Region of the Americas	8.2		8.0	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	26.6	55.5	19.0	59.6
Females	4.9	21.5	3.1	24.0
Both sexes	15.5	44.1	11.1	49.6

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	20.9
Females (15+)	7.7
Both sexes (15+)	16.5

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	25.9	54.2	40.4
Former drinkers* (15+)	26.2	22.9	24.5
Abstainers (15+), past 12 months	52.1	77.1	64.9

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	35.8	18.1	60.8	35.2	785
Road traffic injuries, males / females	25.2	7.9	31.3	21.5	489
Cancer, males / females	150.2	124.8	4.1	1.5	188

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 3 **4** 5 > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	10.3	3.9
Females	3.8	1.8
Both sexes	6.9	2.8
WHO Region of the Americas	8.2	4.1

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	— / —
Excise tax on beer / wine / spirits	— / — / —
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	— / — / —
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	— / — / —
Restrictions on on-/off-premise sales of alcoholic beverages (any):	
Hours, days / places, density	— / —
Specific events / intoxicated persons / petrol stations	— / — / —

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	— / — / —
Legally binding regulations on alcohol advertising / product placement (any)	— / —
Legally binding regulations on alcohol sponsorship / sales promotion (any)	— / —
Legally required health warning labels on alcohol advertisements / containers (any)	— / —
National government support for community action (any)	—
National monitoring system(s) (any)	—