


Liberia


Total population (2016): 4 615 222 ▶ Population aged 15 years and older (15+): 58% ▶ Population in urban areas: 50% ▶ Income group (World Bank): Low income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	3.7		3.5	
Unrecorded	1.6		2.2	
Total**	5.3		5.8	
Total males / females	9.1	1.6	9.9	1.6
WHO African Region	6.3		6.3	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	23.2
Females (15+)	8.7
Both sexes (15+)	18.7

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	21.9	51.3	14.5	54.6
Females	3.5	18.7	2.1	20.5
Both sexes	12.7	41.3	8.4	45.2

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	52.4	74.9	63.7
Former drinkers* (15+)	4.9	6.2	5.6
Abstainers (15+), past 12 months	57.3	81.1	69.2

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	58.5	35.1	60.5	32.5	381
Road traffic injuries, males / females	84.6	40.9	23.3	16.5	345
Cancer, males / females	129.4	113.6	6.6	2.4	80

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 3 4 **5** > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	9.2	3.1
Females	1.3	0.5
Both sexes	5.2	1.8
WHO African Region	3.7	1.3

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions on on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / No, No No / No / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	No / No / No
Legally binding regulations on alcohol advertising / product placement (any)	No / No
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	No
National monitoring system(s) (any)	No