


# Mauritania


Total population (2016): 4 166 463 ► Population aged 15 years and older (15+): 60% ► Population in urban areas: 60% ► Income group (World Bank): Lower-middle income

## ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	0.0		0.0	
Unrecorded	0.1		0.0	
<b>Total**</b>	<b>0.1</b>		<b>0.0</b>	
Total males / females	0.1	0.0	0.1	0.0
WHO African Region	6.3		6.3	

\* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; \*\*adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	4.4
Females (15+)	1.6
Both sexes (15+)	3.8

Prevalence of heavy episodic drinking\* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	0.1	5.3	0.0	5.8
Females	0.0	1.2	0.0	1.3
Both sexes	0.0	4.3	0.0	4.8

\* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	96.9	99.1	98.0
Former drinkers* (15+)	1.6	0.4	1.0
Abstainers (15+), past 12 months	98.6	99.5	99.1

\* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

## HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	53.5	37.6	4.1	1.3	21
Road traffic injuries, males / females	54.5	27.9	0.1	0.1	1
Cancer, males / females	99.5	95.4	0.8	0.1	7

\* Per 100 000 population (15+); \*\*alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score\*, 2016

LEAST < **1** 2 3 4 5 > MOST

\* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016\*

	Alcohol use disorders**	Alcohol dependence
Males	0.7	0.3
Females	0.1	0.0
Both sexes	0.4	0.2
WHO African Region	3.7	1.3

\* 12-month prevalence estimates (15+); \*\*including alcohol dependence and harmful use of alcohol.

## POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Total ban
Excise tax on beer / wine / spirits	Total ban
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	Total ban
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	Total ban
Restrictions on on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Total ban Total ban

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	Total ban
Legally binding regulations on alcohol advertising / product placement (any)	Total ban
Legally binding regulations on alcohol sponsorship / sales promotion (any)	Total ban
Legally required health warning labels on alcohol advertisements / containers (any)	Total ban
National government support for community action (any)	No
National monitoring system(s) (any)	No