
Slovakia


Total population (2016): 5 429 418 ➤ Population aged 15 years and older (15+): 85% ➤ Population in urban areas: 54% ➤ Income group (World Bank): High income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	10.7		10.4	
Unrecorded	1.8		1.6	
Total**	11.9		11.5	
Total males / females	19.3	5.0	18.9	4.6
WHO European Region	11.2		9.8	

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; ** adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine Spirits Other

	Litres
Males (15+)	23.1
Females (15+)	8.0
Both sexes (15+)	16.6

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	55.5	68.1	53.4	76.4
Females	18.3	31.7	17.3	40.9
Both sexes	36.2	52.3	35.8	63.4

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	6.3	19.4	13.1
Former drinkers* (15+)	12.1	22.9	17.7
Abstainers (15+), past 12 months	18.4	42.3	30.8

 $^{^{\}star}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

(1011), 2010					
	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	40.9	12.7	77.5	62.0	1 118
Road traffic injuries, males / females	11.7	3.6	48.1	36.2	164
Cancer, males / females	305.9	155.8	11.8	3.3	1 199

^{*}Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016 LEAST < 1 2 3 4 5 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	22.8	10.2
Females	2.5	1.1
Both sexes	12.2	5.5
WHO European Region	8.8	3.7

 $^{^{\}star}$ 12-month prevalence estimates (15+); ** including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (1993/2016) / Yes
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	No, No / Yes, No Yes / Yes / Yes

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in $\%$	Zero tolerance
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	No / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes

^{*} Based on alcohol-attributable years of life lost.