


Slovenia


Total population (2016): 2 069 362 ► Population aged 15 years and older (15+): 85% ► Population in urban areas: 50% ► Income group (World Bank): High income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	10.5		10.8	
Unrecorded	1.0		1.8	
Total**	11.5		12.6	
Total males / females	18.4	4.7	20.4	5.1
WHO European Region	11.2		9.8	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	24.6
Females (15+)	8.5
Both sexes (15+)	17.7

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	56.3	67.9	56.8	78.1
Females	19.2	32.0	19.7	43.3
Both sexes	37.5	52.6	38.8	65.2

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	5.6	17.4	11.6
Former drinkers* (15+)	11.5	22.6	17.1
Abstainers (15+), past 12 months	17.1	40.0	28.7

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	31.2	8.7	78.1	62.5	391
Road traffic injuries, males / females	9.6	2.8	50.1	33.9	53
Cancer, males / females	253.4	145.0	8.3	3.3	387

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 2 **3** 4 5 > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	23.5	10.5
Females	4.5	2.0
Both sexes	13.9	6.2
WHO European Region	8.8	3.7

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2000/2016) / No
Excise tax on beer / wine / spirits	Yes / No / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions for on-/off-premise sales of alcoholic beverages (any):	
Hours, days / places, density	Yes, No / Yes, No
Specific events / intoxicated persons / petrol stations	Yes / Yes / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	0.05 / 0.00 / 0 / 00
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	No / No
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / No
National government support for community action (any)	Yes
National monitoring system(s) (any)	Yes