
Uzbekistan


Total population (2016): 30 300 446 > Population aged 15 years and older (15+): 72% > Population in urban areas: 38% > Income group (World Bank): Lower-middle income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	20	10*	20	16*
Recorded	1.8		1	.6
Unrecorded	1.4		1.1	
Total**	3.2		2.7	
Total males / females	5.6	0.9	4.8	0.7
WHO European Region	11.2		9.8	

 $^{^{\}star}$ Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; ** adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

Beer

Wine Spirits

Other All

	Litres
Males (15+)	15.8
Females (15+)	5.7
Both sexes (15+)	12.8

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	14.6	48.6	9.1	52.7
Females	2.0	16.9	1.2	19.3
Both sexes	8.1	39.1	5.2	44.2

^{*} Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	38.5	68.3	53.9
Former drinkers* (15+)	31.4	19.7	25.4
Abstainers (15+), past 12 months	69.9	88.0	79.2

 $^{^{\}ast}$ Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASI	OR*	AAF	(%)	AAD** (Number)
Liver cirrhosis, males / females	46.2	31.7	58.6	29.2	3 234
Road traffic injuries, males / females	20.0	4.5	20.7	14.5	557
Cancer, males / females	86.0	74.2	7.8	1.8	657

^{*}Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016 LEAST < 1 2 3 4 5 > MOST

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	9.9	5.3
Females	2.1	1.3
Both sexes	5.9	3.3
WHO European Region	8.8	3.7

 $^{^{\}star}$ 12-month prevalence estimates (15+); ** including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	Yes (2011/—) / Yes
Excise tax on beer / wine / spirits	Yes / No / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	20 / 20 / 20
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	20 / 20 / 20
Restrictions for on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, No / Yes, Yes Yes / Yes / Yes

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	Zero tolerance
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	Yes / Yes
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / Yes
National government support for community action (any)	Yes
National monitoring system(s) (any)	No

^{*} Based on alcohol-attributable years of life lost.