

CONSOLIDATED GUIDELINES ON  
**PERSON-CENTRED  
HIV PATIENT MONITORING  
AND CASE SURVEILLANCE**

**ANNEX 2.3.5**  
ART REGISTER

JUNE 2017

ART register Page 1 District \_\_\_\_\_ Health facility \_\_\_\_\_

Cohort year: _____ Month: _____																					
Registration and personal information						Status at ART start			Fill when applicable					Pregnancy (also record "P" as follow-up status) (For each pregnancy, record EDD, ANC no. and HEI no.)					1st-line regimen	2nd-line regimen	3rd-line regimen
ART start date	Unique ID no.	Patient clinic ID	Name <i>Surname Given name</i>	Sex	Age	Pregnant Breastfeeding	TB+	CD4 (% if U5)	CTX <i>Start month/ year</i>	TB prev. therapy <i>Start, complete month/year</i>	TB Rx <i>Start month/year and TB registration no.</i>	HBsAg test <i>Month/year</i>	HCV Ab test (with reflex HCV RNA if +) <i>Month/ year</i>	Preg 1	Preg 2	Preg 3	Preg 4	Original regimen	Substitutions <i>1st: Reason/Date 2nd: Reason/Date</i>	Switches, substitutions <i>1st: Reason/Date 2nd: Reason/Date</i>	Switches, substitutions <i>1st: Reason/Date 2nd: Reason/Date</i>
1																					
2																					
3																					
4																					
5																					
6																					
7																					
8																					
9																					
10																					
11																					
12																					
13																					
14																					
15																					
16																					
17																					
18																					
19																					
20																					
Sub-totals																					
Adult/adolescent 1st-line regimens 1a = TDF-3TC-EFV 1b = TDF-FTC-EFV 1c = ... 1d = Other	Child <10 years 1st-line regimens 4a = ABC-3TC-EFV 4b = ABC-3TC-LPV/r 4c = AZT-3TC-LPV/r 4d = ... 4e = Other	Adult/adolescent 2nd-line regimens 2a = TDF-3TC-ATV/r 2b = TDF-FTC-ATV/r 2c = TDF-3TC-LPV/r 2d = TDF-FTC-LPV/r 2e = AZT-3TC-ARV/r 2f = AZT-3TC-LPV/r 2g = ... 2h = Other	Child <10 years 2nd-line regimens 5a = AZT-3TC-RAL 5b = ABC-3TC-RAL 5c = AZT-3TC-EFV 5d = AZT-3TC-RAL 5e = TDF-3TC-EFV 5f = TDF-3TC-RAL 5g = ... 5h = Other	Adult/adolescent 3rd-line regimens 3a = DRV/r-DTG-??? 3b = DRV/r-RAL-??? 3c = DTG-??? 3d = RAL-??? 3e = DRV/r-??? 3f = ... 3g = Other	Child <10 years 3rd-line regimens 6a = DRV/r-??? 6b = DRV/r-DTG 6c = DRV/r-RAL 6d = ... 6e = Other	Reasons for substitution 1 Toxicity/side-effects (record code for toxicity) 2 Pregnancy 3 Due to new TB 4 New drug available 5 Drug out of stock 6 Other reason (specify)	Reasons for switch to 2nd- or 3rd-line regimen 7 Clinical treatment failure 8 Immunological failure 9 Virological failure	Treatment-limiting toxicity 1 GI 2 Skin 3 CNS 4 Hepatic dysfunction/jaundice 5 Haematological 6 Fatigue 7 Bone dysfunction 8 Metabolic 9 Headache 10 Kidney dysfunction													

ART register Page 2

Cohort year: _____ Month: _____ Write in year and month																																
Month 0	Month 1	2	3	4	5	6			7	8	9	10	11	12			13	14	15	16	17	18		19	20	21	22	23	24			
						Viral load	CD4 (% if U5)	Follow-up status						Viral load	CD4 (% if U5)	Follow-up status						CD4 (% if U5)	Follow-up status							Viral load	CD4 (% if U5)	Follow-up status
1																																
2																																
3																																
4																																
5																																
6																																
7																																
8																																
9																																
10																																
11																																
12																																
13																																
14																																
15																																
16																																
17																																
18																																
19																																
20																																
Sub-totals																																
<b>Top row:</b> Record follow-up status at end of each month On treatment (record current regimen code) <b>DEAD</b> <b>STOP</b> (add why STOP code), e.g. STOP/1 <b>RESTART</b> (record regimen code), e.g. RESTART/1b <b>LTF</b> Lost to follow up (not seen for at least 3 months from last missed appointment) <b>TI</b> Transfer in (draw a line under list of original cohort patients and enter <b>TI</b> patients [starting ART in same month/year] under line, recording <b>TI</b> in margin and continue follow up) <b>TO</b> Transfer out (record to where), e.g. TO/facility X						<b>Bottom row:</b> Record if currently pregnant or breastfeeding						<b>If follow-up status is "STOP", then add reason(s) (and weeks of interruption if later restarted), e.g. STOP/1/5 weeks</b> 1 Severe illness, hospitalization 2 Drugs out of stock 3 Patient lacks finances 4 Excluded HIV infection in infant 5 Other (specify)						<b>CD4/Viral load:</b> <b>Top row:</b> Record date of test <b>Bottom row:</b> Record test result <b>CD4:</b> cells/mm <sup>3</sup> or % if U5 <b>VL:</b> copies/mL														



ART register Page 4

Cohort year:		Month:		Write in year and month																								
49	50	51	52	53	54	55	56	57	58	59	60 (5 years)			61	62	63	64	65	66	67	68	69	70	71	72 (6 years)			
											Viral load	CD4 (% if U5)	Follow-up status													Viral load	CD4 (% if U5)	Follow-up status
1																												
2																												
3																												
4																												
5																												
6																												
7																												
8																												
9																												
10																												
11																												
12																												
13																												
14																												
15																												
16																												
17																												
18																												
19																												
20																												
Sub-totals																												
<b>Top row:</b> Record follow-up status at end of each month On treatment (record current regimen code) <b>DEAD</b> <b>STOP</b> (add why STOP code) <b>RESTART</b> (record regimen code) <b>LTF</b> Lost to follow up (not seen for at least 3 months from last missed appointment) <b>TI</b> Transfer in (draw a line after original cohort, and enter <b>TI</b> patients [starting ART in same month/year] under line, recording <b>TI</b> in margin and continue follow up) <b>TO</b> Transfer out (record to where)												<b>Bottom row:</b> Record if currently pregnant or breastfeeding			<b>If follow-up status is "STOP", then add reasons (and weeks of interruption if later restarted):</b> 1 Severe illness, hospitalization 2 Drugs out of stock 3 Patient lacks finances 4 Excluded HIV infection in infant 5 Other (specify)			<b>CD4/Viral load:</b> <b>Top row:</b> Record date of test <b>Bottom row:</b> Record test result <b>CD4:</b> cells/mm <sup>3</sup> or % if U5 <b>VL:</b> copies/mL										



ART register Page 5

Cohort year:		Month:		Write in year and month																															
		73	74	75	76	77	78	79	80	81	82	83	84 (7 years)			85	86	87	88	89	90	91	92	93	94	95	96 (8 years)								
													Viral load	CD4 (% if U5)	Follow-up status													Viral load	CD4 (% if U5)	Follow-up status					
1																																			
2																																			
3																																			
4																																			
5																																			
6																																			
7																																			
8																																			
9																																			
10																																			
11																																			
12																																			
13																																			
14																																			
15																																			
16																																			
17																																			
18																																			
19																																			
20																																			
Sub-totals																																			
<b>Top row:</b> Record follow-up status at end of each month On treatment (record current regimen code) <b>DEAD</b> <b>STOP</b> (add why STOP code) <b>RESTART</b> (record regimen code) <b>LTF</b> Lost to follow up (not seen for at least 3 months from last missed appointment) <b>TI</b> Transfer in (draw a line after original cohort, and enter <b>TI</b> patients [starting ART in same month/year] under line, recording <b>TI</b> in margin and continue follow up) <b>TO</b> Transfer out (record to where)													<b>Bottom row:</b> Record if currently pregnant or breastfeeding													<b>If follow-up status is "STOP", then add reasons (and weeks of interruption if later restarted):</b> 1 Severe illness, hospitalization 2 Drugs out of stock 3 Patient lacks finances 4 Excluded HIV infection in infant 5 Other (specify)					<b>CD4/Viral load:</b> <b>Top row:</b> Record date of test <b>Bottom row:</b> Record test result <b>CD4:</b> cells/mm <sup>3</sup> or % if U5 <b>VL:</b> copies/mL				

ART register Page 6

Cohort year:		Month:				Write in year and month																						
97	98	99	100	101	102	103	104	105	106	107	108 (9 years)			109	110	111	112	113	114	115	116	117	118	119	120 (10 years)			
											Viral load	CD4 (% if U5)	Follow-up status													Viral load	CD4 (% if U5)	Follow-up status
1																												
2																												
3																												
4																												
5																												
6																												
7																												
8																												
9																												
10																												
11																												
12																												
13																												
14																												
15																												
16																												
17																												
18																												
19																												
20																												
Sub-totals																												
<b>Top row:</b> Record follow-up status at end of each month On treatment (record current regimen code) <b>DEAD</b> <b>STOP</b> (add why STOP code) <b>RESTART</b> (record regimen code) <b>LTF</b> Lost to follow up (not seen for at least 3 months from last missed appointment) <b>TI</b> Transfer in (draw a line after original cohort, and enter TI patients [starting ART in same month/year] under line, recording TI in margin and continue follow up) <b>TO</b> Transfer out (record to where)												<b>Bottom row:</b> Record if currently pregnant or breastfeeding			<b>If follow-up status is "STOP", then add reasons (and weeks of interruption if later restarted):</b> 1 Severe illness, hospitalization 2 Drugs out of stock 3 Patient lacks finances 4 Excluded HIV infection in infant 5 Other (specify)			<b>CD4/Viral load:</b> <b>Top row:</b> Record date of test <b>Bottom row:</b> Record test result <b>CD4:</b> cells/mm <sup>3</sup> or % if U5 <b>VL:</b> copies/mL										



List of patients who may not or will not start ART soon after enrolment into HIV care\*

	Date enrolled in HIV care	Unique ID no.	Patient clinic no.	Name <i>Surname Given name</i>	Sex	Age	Status at enrolment		Outcome <i>DEAD TO LTF Start ART</i>	Outcome date
							Baseline CD4	TB prev. therapy started <i>month /year</i>	Active TB disease <i>month/ year</i>	
1										
2										
3										
4										
5										
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
16										
17										
18										
19										
20										
21										
22										
23										
24										
25										
26										
27										
28										
29										
30										
31										
32										
33										
34										
35										
36										
37										
38										
39										
40										
Subtotals										

\*Patients who start the process of initiating ART (e.g. adherence preparation) and will start ART soon thereafter should not be entered here unless they decide to delay treatment.