


Federal Democratic Republic of Ethiopia

African Region



I. Epidemiological Profile

Population Distribution (Country reported)

Population	2024	%
High transmission (>1 case per 1000 population)	90.1M	81.9%
Low transmission (0-1 case per 1000 population)	14.8M	13.5%
Malaria free (0 cases)	5.0M	4.6%
Total	110.0M	100%

Parasites and Vectors

Major *Plasmodium* Species

*P. falciparum**: 65.05%

P. vivax: 34.88%

*Includes mixed infections and other species of Plasmodium

Major *Anopheles* Species

An. arabiensis, *An. stephensi*

Reported Cases and Deaths

Total cases (presumed + confirmed)	4.5M
Reported indigenous confirmed cases	4.3M
Confirmed cases from public sector (health facility)	3.6M
Confirmed cases at community level	496.7K
Confirmed cases from private sector	187.2K
Indigenous deaths	1.5K

WHO Estimates

12.4M
Estimated Cases
Confidence Interval: [6.9M,18.6M]
22.4K
Estimated Deaths
Confidence Interval: [8.4K,43.4K]

II. Intervention policies and strategies

Interventions and Policies

Intervention	Policies/Strategies	Policy	Year
Diagnosis	Malaria diagnosis with either microscopy or RDTs are free in the public sector	Yes	2005
Treatment	ACTs for malaria treatment is free in the public sector	Yes	2005
	ACT is delivered at community	Yes	2005
	Pre-referral Rx with rectal artesunate suppositories at community level	Yes	2015
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	Yes*	2018
	The sale of oral artemisinin-based monotherapy drugs	has never been allowed	-
IPT	IPT used to prevent malaria during pregnancy	No	-
	Community based delivery of IPTp (c-IPTp) is used to prevent malaria during pregnancy aligned with WHO recommendation	No	-
	Seasonal malaria chemoprevention (SMC) is used, aligned with WHO recommendation	No	-
	Perennial Malaria Chemoprevention (PMC) is used, aligned with WHO recommendation	No	-
	Intermittent Preventive Treatment in school-aged children (IPTsc) is used, aligned with WHO recommendation	No	-
Surveillance	Malaria is a notifiable disease	Yes	-
	Case investigation and classification is undertaken	Yes*	2019
	Foci investigation and classification is undertaken	Yes	-
	Case reporting from private sector is mandatory	Yes	2016
ITN	ITNs distributed free of charge through through mass campaign to all age groups	Yes	-
	ITNs distributed free of charge through routine channels to all age groups	No	-
	ITNs durability is monitored	Yes	-
IRS	IRS is an intervention at the NMP	Yes	1960

Antimalarial Treatment Policy

Antimalaria treatment policy	Medicine	Year adopted
First-line treatment of unconfirmed malaria	AL+PQ	2014
First-line treatment of <i>P. falciparum</i>	AL+PQ	2014
For treatment failure of <i>P. falciparum</i>	DHA-PPQ	2022
Treatment of severe malaria	AS; AL+PQ	2018
Treatment of <i>P. vivax</i>	CQ+PQ	2018

Type of RDT used (public)

P.f + P.v specific (Combo)

Therapeutic efficacy tests (clinical and parasitological failure, %)

Medicine	Year	Min	Median	Max	Follow-up	No. of studies	Species
AL	2015-2023	0	3.1	8	28 days	8	Pf
CQ	2016-2021	0	0.6	4	28 days	8	Pv
CQ+PQ	2019-2020	0	0	0	28 days	1	Pv
DHA-PPQ	2017-2021	0	0	0	42 days	2	Pv

Resistance status by insecticide class (2020-2024) and use of class for malaria vector control (2024)

Insecticide class	(%) sites ¹	Vectors ²	Used ³
Carbamates	45.5 (15/33)	<i>An. gambiae s.l.</i> , <i>An. stephensi</i> , <i>Other species</i>	No
Neonicotinoids	0 (0/20)	-	Yes
Organophosphates	0 (0/35)	-	Yes
Pyrethroids	83.7 (41/49)	<i>An. gambiae s.l.</i> , <i>An. stephensi</i> , <i>Other species</i>	Yes

¹ Percent of sites for which resistance is confirmed and total number of sites that reported data

² Vectors reported to exhibit resistance to insecticide class

³ Class reported as used for malaria control in 2024

Intervention	Policies/Strategies	Policy	Year
	DDT is used for IRS	No	-
Larval source management	Use of Larval source management	Yes	2012


Yes* = Policy adopted, but not implemented in 2024, or no data exist to support implementation

No = Policy does not exist or policy has been discontinued


Earliest year that policy is adopted was adjusted based on the earliest year that the WHO policy was recommended

III. Data Visualization

Estimated and reported cases




Treatment seeking and reporting completeness




Cases tested and treated




Test positivity




Confirmed malaria cases per 1000 population at risk




Malaria inpatients and deaths




Coverage of ITN and IRS



Sources of financing



Government expenditure by intervention in 2024



Footnotes

Country profiles are generated automatically based on data reported by countries. They are available for all current malaria endemic countries and areas asked to report to the Malaria and Neglected Tropical Diseases Programme annually. Country profiles are based on data validated by the countries as of 15 October 2025.

Survey sources: DHS 2016

Further information on the methods used to estimate malaria cases and deaths can be found in Annex I of the 2025 WMR, and an explanation for the gap between estimated and reported confirmed indigenous cases is provided in [mpac-april2018-erg-report-malaria-burden-session6.pdf](https://cdn.who.int/media/docs/default-source/malaria/mpac-documentation/mpac-april2018-erg-report-malaria-burden-session6.pdf) (https://cdn.who.int/media/docs/default-source/malaria/mpac-documentation/mpac-april2018-erg-report-malaria-burden-session6.pdf?sfvrsn=44e72782_2).

World Malaria Report 2025