

Cook Islands STEPS Survey 2003-2004

Fact Sheet

The STEPS survey of chronic disease risk factors in Cook Islands was carried out from September 2003 to May 2004. Cook Islands carried out Step 1, Step 2 and Step 3. Socio demographic and behavioural information was collected in Step 1. Physical measurements such as height, weight and blood pressure were collected in Step 2. Biochemical measurements were collected to assess blood glucose and cholesterol levels in Step 3. The STEPS survey in Cook Islands was a population-based survey of adults aged 25-64. A multi-stage cluster sampling design was used to produce representative data for that age range in Cook Islands. A total of 2036 adults participated in the Cook Islands STEPS survey. The response rate was 90%.

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females
Step 1 Tobacco Use			
Percentage who currently smoke tobacco	43.9% (40.0 – 47.9)	46.6% (41.4 – 51.8)	41.1% (35.0 – 47.2)
Percentage who currently smoke tobacco daily	33.3% (30.4 – 36.2)	37.5% (33.3 – 41.7)	28.8% (24.9– 32.7)
For those who smoke tobacco daily			
Average age started smoking (years)	20.3 (19.5 – 21.1)	19.7 (18.7 – 20.6)	21.2 (20.0 – 22.4)
Percentage of daily smokers smoking manufactured cigarettes	62.3% (53.7 – 70.9)	55.2% (42.8 – 67.7)	72.1% (61.6 – 82.6)
Mean number of manufactured cigarettes smoked per day (by smokers of manufactured cigarettes)	8.1 (6.6 – 9.5)	8.1 (6.0 – 10.1)	8.0 (6.1 – 10.0)
Step 1 Alcohol Consumption			
Percentage who are lifetime abstainers	26.8% (22.6 – 31.0)	17.9% (14.2 – 21.6)	36.3% (29.6 – 42.9)
Percentage who are past 12 month abstainers	10.3% (8.0 – 12.7)	7.7% (4.2 – 11.1)	13.2% (11.0 – 15.4)
Percentage who currently drink (drank alcohol in the past 30 days)	62.9% (57.1 – 68.6)	74.4% (68.7– 80.2)	50.6% (43.2– 57.9)
Percentage who engage in heavy episodic drinking (men who had 5 or more / women who had 4 or more drinks on any day in the past 30 days)		74.7% (68.9 – 80.5)	51.6% (47.8 – 55.5)
Step 1 Fruit and Vegetable Consumption (in a typical week)			
Mean number of days fruit consumed	3.7 (3.4 – 3.9)	3.4 (3.1 – 3.6)	4.0 (3.7 – 4.3)
Mean number of servings of fruit consumed on average per day	1.5 (1.4 – 1.6)	1.4 (1.2 – 1.6)	1.6 (1.5 – 1.8)
Mean number of days vegetables consumed	4.0 (3.7 – 4.3)	3.7 (3.3 – 4.1)	4.2 (3.8 – 4.7)
Mean number of servings of vegetables consumed on average per day	1.5 (1.4 – 1.7)	1.5 (1.3 – 1.7)	1.6 (1.4 – 1.8)
Percentage who ate less than 5 servings of fruit and/or vegetables on average per day	81.8% (79.4 – 84.2)	83.5% (80.4 – 86.7)	79.9% (76.2 – 83.7)
Step 1 Physical Activity			
Percentage with low levels of activity (defined as < 600 MET-minutes per week)*	75.3% (71.6 – 79.1)	74.4% (68.2– 80.6)	76.3% (72.0 – 80.6)
Percentage with high levels of activity (defined as ≥ 3000 MET-minutes per week)*	11.8% (9.2 – 14.4)	14.1% (9.9 – 18.4)	9.6% (7.2 – 12.1)
Median time spent in physical activity on average per day (minutes) (presented with inter-quartile range)	0.0 (0.0 – 38.6)	0.0 (0.0 – 42.9)	0.0 (0.0 – 34.3)
Percentage not engaging in vigorous activity	92.6% (89.9 – 95.3)	89.0% (84.7 – 93.4)	95.8% (94.5 – 97.2)

^{*} For complete definitions of low and high levels of physical activity, other conditions are specified in the GPAQ Analysis Guide, available at: http://www.who.int/chp/steps/GPAQ/en/index.html



Cook Islands STEPS Survey 2003-2004

Fact Sheet

Results for adults aged 25-64 years (incl. 95% CI)	Both Sexes	Males	Females	
Step 2 Physical Measurements				
Mean body mass index - BMI (kg/m²)	32.8 (32.4 – 33.2)	32.3 (31.9 – 32.8)	33.3 (32.7 – 33.8)	
Percentage who are overweight (BMI ≥ 25 kg/m²)	88.5% (86.9 – 90.1)	89.8% (87.8 – 91.9)	87.1% (84.8 – 89.3)	
Percentage who are obese (BMI ≥ 30 kg/m²)	61.4% (58.7 – 64.2)	57.4% (53.8 – 61.1)	65.7% (62.5 – 68.9)	
Average waist circumference (cm)		103.7 (102.6 – 104.7)	100.0 (98.5 – 101.5)	
Mean systolic blood pressure - SBP (mmHg), including those currently on medication for raised BP	129.1 (125.7 – 132.5)	134.6 (132.1 – 137.2)	123.3 (119.4 – 127.3)	
Mean diastolic blood pressure - DBP (mmHg) , including those currently on medication for raised BP	80.6 (78.9 – 82.3)	83.1 (80.8 – 85.3)	78.0 (76.4 – 79.6)	
Percentage with raised BP (SBP ≥ 140 and/or DBP ≥ 90 mmHg or currently on medication for raised BP)	33.2% (26.5 – 39.9)	40.6% (33.0 – 48.1)	25.5% (16.3 – 34.8)	
Step 3 Biochemical Measurement				
Mean fasting blood glucose, including those currently on medication for raised blood glucose: mmol/L	6.3 (6.1 – 6.4)	6.4 (6.2 – 6.6)	6.1 (5.9 – 6.3)	
Percentage with impaired fasting glycaemia as defined below • capillary whole blood value ≥5.6 mmol/L (100 mg/dl) and <6.1 mmol/L (110 mg/dl)	41.8% (38.1 – 45.4)	47.3% (41.8 – 52.7)	36.0% (33.3 – 38.8)	
Percentage with raised fasting blood glucose as defined below or currently on medication for raised blood glucose • capillary whole blood value ≥ 6.1 mmol/L (110 mg/dl)	23.6% (19.7 – 27.4)	26.1% (20.0 – 32.2)	21.0% (16.4 – 25.5)	
Mean total blood cholesterol, including those currently on medication for raised cholesterol: mmol/L	5.1 (5.0 – 5.2)	5.1 (5.0 – 5.2)	5.0 (4.9 – 5.1)	
Percentage with raised total cholesterol (≥ 5.0 mmol/L or ≥ 190 mg/dl or currently on medication for raised cholesterol)	75.2% (71.9 – 78.5)	77.1% (72.4 – 81.9)	73.2% (68.8 – 77.5)	
Summary of combined risk factors				
 less than 5 servings of fruits & vegetables per day 	verweight (BMI \ge 25 kg/m ²) aised BP (SBP \ge 140 and/or DBP \ge 90 mmHg or urrently on medication for raised BP)			
Percentage with none of the above risk factors	0.3% (0.0 – 0.6)	0.3% (0.0 – 0.8)	0.3% (0.0 – 0.7)	
Percentage with three or more of the above risk factors, aged 25 to 44 years	75.1% (71.2 – 79.0)	81.0% (76.7 – 85.3)	70.0% (66.4 – 73.6)	
Percentage with three or more of the above risk factors, aged 45 to 64 years	80.5% (77.4 – 83.5)	83.8% (80.2 – 87.4)	77.1% (73.6 – 80.5)	
Percentage with three or more of the above risk factors, aged 25 to 64 years	76.6% (73.6 – 79.7)	81.8% (78.5 – 85.1)	71.9% (68.7 – 75.1)	