

[EXT] ECUADOR - comments on the second draft of the WHO Discussion paper - prevention and control of NCDs

Mision Permanente ONU - Ginebra <onuginebra@cancilleria.gob.ec>

Tue 20/09/2022 10:24

To: NCDappendix3 <NCDappendix3@who.int>

Cc: Mision Permanente ONU - Ginebra <onuginebra@cancilleria.gob.ec>

 1 attachments (890 KB)

2022_discussion_paper_01_aug_revisado_dnepc_22082022 (1).pdf;

Dear Sir/Madam:

In relation to the process to update the set of best buys and other recommended interventions for the prevention and control of NCDs (Updated Appendix 3 of the WHO Global NCD Action Plan), please find attached the comments on the second draft of the WHO Discussion paper, submitted by the Ministry of Public Health of Ecuador.

Kind regards

MISION PERMANENTE DEL ECUADOR ANTE LA ONU – GINEBRA

Chemin Camille-Vidart 15 – 1202 Ginebra – Suiza

(0041) 022 732 49 55

onuginebra@cancilleria.gob.ec



Draft Updated Appendix 3 of the WHO Global NCD action plan 2013-2030**MANDATE**

1. Paragraph 3(a) of decision WHA72(11) requested the Director-General to, inter alia:

“to propose updates to the appendices of WHO’s Global Action Plan for the Prevention and Control of NCDs 2013-2020 and WHO’s comprehensive mental health action plan 2013–2020, as appropriate, in consultation with Member States and taking into account the views of other stakeholders, ensuring that the action plans remain based on scientific evidence for the achievement of previous commitments for the prevention and control of noncommunicable diseases, including Sustainable Development Goal target 3.4 (by 2030, reduce by one third premature mortality from noncommunicable diseases through prevention and treatment and promote mental health and well-being) and other related goals and targets”¹

This WHO Discussion paper is prepared pursuant to this decision.

BACKGROUND

2. In line with paragraph 3(a) of decision WHA72(11), the WHO Secretariat is updating the 2017 version of the Appendix 3 of the WHO Global NCD Action Plan 2013-2020, now extended until 2030, considering new scientific evidence as well as new WHO recommendations, since the 2017 update.
3. The WHO Global NCD Action Plan contains five appendices. Appendix 3 consists of a menu of policy options to support the implementation of the 6 objectives of the WHO Global NCD Action Plan. It outlines a list of policy options and cost-effective interventions, Member States should prioritize in order to prevent and control NCDs. The last revision of the Appendix 3 of the WHO Global NCD Action Plan² was endorsed by the World Health Assembly in 2017 (decision WHA70.11). That endorsement by WHA70, in May 2017, enabled the WHO Secretariat to include an updated Appendix 3 in the report of the UN Secretary-General to the United Nations General Assembly on NCDs of December 2017 (see paragraphs 10, 15, Table 3, Table 5, 53, 55, 59 and 60 of A/72/662).

¹ World Health Assembly 74 (2019) Follow-up to the political declaration of the third high-level meeting of the General Assembly on the prevention and control of non-communicable diseases. WHA72(11). [https://apps.who.int/gb/ebwha/pdf_files/WHA72/A72\(11\)-en.pdf](https://apps.who.int/gb/ebwha/pdf_files/WHA72/A72(11)-en.pdf).

² World Health Organization (2017). Tackling NCDs: 'best buys' and other recommended interventions for the prevention and control of noncommunicable diseases. World Health Organization. <https://apps.who.int/iris/handle/10665/259232>.

4. The 2017 update of Appendix 3 contained 88 recommendations including 16 interventions which were classified as “very cost-effective and affordable interventions for all Member States”³ (previously referred to as the “best buys”), as well as other cost-effective interventions (previously referred to as “good buys”)⁴.

5. The main reasons for updating Appendix 3 in 2022 are:

- To take into consideration the emergence of new evidence of cost-effectiveness and/or new WHO recommendations since the adoption of the WHO Global NCD Action Plan 2013-2020,
- To assist countries selecting a combination of these interventions to define locally tailored packages of interventions to accelerate ongoing national NCD responses.

6. This updated version of the Appendix 3, will complement existing global strategies and action plans and a number of new technical products that WHO secretariat is developing along with the NCD Implementation Roadmap for the WHO Global NCD Action Plan ⁵ including the WHO menu of cost-effectiveness intervention for mental health⁶, the recommended interventions to address the health impact of air pollution^{7,8} and the menu of cost-effective interventions for oral health⁹ that are being developed.

PROCESS FOLLOWED BY THE SECRETARIAT TO UPDATE APPENDIX 3

7. The WHO secretariat used the WHO-CHOICE methodology for updating appendix 3 which was endorsed at the World Health Assembly in May 2016¹⁰. This is to retain consistency and familiarity with existing Appendix 3 and previously agreed methods and global strategies. This update is accompanied by a methodological annex providing more information on the methodology used to identify and analyse interventions, and includes assumptions used in the WHO-CHOICE economic modelling and describing updates from the 2017 update. Detailed information on the evidence and assumptions underlying the interventions are provided in separate technical briefs by disease and risk factor area.

8. The Secretariat identified a revised list of 83 interventions (excluding overarching/enabling actions). These include:

- 31 interventions which have been unchanged from the 2017 update and for which key parameters were updated and cost-effectiveness estimates were generated,

³ Generate an extra year of healthy life for a cost that falls below the average annual income or GDP per person

⁴ WHO (2017). Tackling NCDs: 'best buys' and other recommended interventions for the prevention and control of noncommunicable diseases. World Health Organization. <https://apps.who.int/iris/handle/10665/259232>.

⁵ WHO (2022). Draft implementation road map 2023–2030 for the global action plan for the prevention and control of noncommunicable diseases 2013–2030. Annex 1. Document A75/10 Add.8. World Health Organization. https://apps.who.int/gb/ebwha/pdf_files/WHA75/A75_10Add8-en.pdf

⁶ WHO (2021) WHO menu of cost-effective interventions for mental health. Geneva: World Health Organization. <https://www.who.int/publications/i/item/9789240031081>.

⁷ Compendium of WHO and other UN guidance on health and the environment: <https://www.who.int/tools/compendium-on-health-and-environment>

⁸ WHO (2021) WHO global air quality guidelines: particulate matter (PM2.5 and PM10), ozone, nitrogen dioxide, sulfur dioxide and carbon monoxide. Geneva: World Health Organization. <https://www.who.int/publications/i/item/9789240034228>

⁹ https://apps.who.int/gb/ebwha/pdf_files/WHA74/A74_R5-en.pdf

¹⁰ http://apps.who.int/gb/ebwha/pdf_files/WHA69/A69_1Rev1-en.pdf

- 9 interventions from the 2017 update which have been revised to reflect updates in WHO policy or scientific evidence,
- 8 interventions included in the 2017 update that had no analysis carried out at the time and for which cost-effectiveness analysis was performed for the 2022 update,
- 10 new interventions from new WHO guidance and tools, and
- 25 interventions that are considered priorities but have no cost-effectiveness analysis.

For the WHO-CHOICE analysis, interventions were considered for inclusion if they have a WHO recommendation as well as evidence required for the modelling. Interventions included without WHO-CHOICE analysis are drawn from other existing WHO guidance documents and WHO-CHOICE analysis will be considered in future updates. Individual-level interventions that were not been derived from existing WHO disease management guidelines or did not have sufficient data for the WHO-CHOICE modelling have been removed. Their inclusion will be considered in future updates.

9. The list of interventions is not exhaustive but is intended to provide information and guidance on effectiveness and cost-effectiveness of population-based and individual interventions based on current evidence, and to act as the basis for future work to develop and expand the evidence base.

10. It is acknowledged that there are inherent challenges in identifying a core list of interventions whilst being comprehensive enough to meet the needs of Member States. The limitations of considering cost-effectiveness in isolation are recognised. It is recommended that additional criteria reflecting national values are used when determining the list of interventions to be adopted by the Member States¹¹.

11. Following the first web-based and information consultation organized in June, the Secretariat has taken into consideration the feedback received in this second draft of the discussion paper.

KEY CHANGES AND PROPOSED UPDATED APPENDIX 3

12. The key changes compared to the 2017 update include:

- Changes in the definition of some interventions and inclusion of new interventions to align with new WHO policy and guidance. For example, dietary interventions moved from an approach focusing on the health impact and cost of salt intake only to interventions assessing the health outcomes and cost of interventions addressing multiple nutrients such as fibres, vegetables, and salt. Similarly, interventions for chronic respiratory diseases (asthma and chronic obstructive pulmonary disease) now distinguish between acute and long-term patient management to be in line with WHO PEN guidance.
- Changes to the modelling of interventions. This includes changes such as the inclusion of a different population in need or adding new health outcomes due to additional evidence on impact being available. For example, for cardiovascular diseases, the use of new risk charts and threshold leads to different populations in need and drug therapies.
- The inclusion and use of country-level data for 62 low -and middle-income countries compared to regional data for 20 low, middle and high-income countries in the 2017 update.

¹¹ WHO (2021) Principles of health benefit packages. <https://www.who.int/publications/item/9789240020689>

13. Annex 1 outlines the proposed structure and content of the updated Appendix 3 that will be submitted to the 152nd session of WHO Executive Board (EB152) for consideration by Member States at the Seventy-sixth World Health Assembly (WHA76) in 2023. This includes:

- A proposed preamble text to the Appendix 3
- The updated list of interventions for the main Appendix 3, and
- A Methodological Annex

TIMELINE AND NEXT STEPS

14. As per the approach shared with Members States, the timeframe and next steps for finalising the updated Appendix 3 is as follows:

Dates	Activity
June 1 st to 26 June 2022	The Secretariat will publish a first draft WHO Discussion Paper containing a proposed updated Appendix 3 for a web-based consultation; including available results from completed CHOICE analyses
20 June 2022	The Secretariat will convene a first informal consultation with Member States and UN organizations
21 June 2022	The Secretariat will convene a first informal consultation with non-State actors (nongovernmental organizations, private sector entities, philanthropic foundations, and academic institutions).
1 August to 28 August 2022	The Secretariat will publish a second draft WHO Discussion Paper containing a proposed updated Appendix 3 for a web-based consultation; including updated results on CHOICE analyses.
16 September 2022	The Secretariat will convene a second informal consultation with Member States and UN organizations
19 September 2022	The Secretariat will convene a second informal consultation with non-State actors

Annex 1. Proposed updated Appendix 3 of the Global Action Plan for the prevention and control of noncommunicable diseases 2013-2030

What is appendix 3

Provided at the request of Member States, Appendix 3 is a menu of policy options and cost-effective interventions for prevention and control of major noncommunicable diseases. The purpose is to assist Member States in implementing, as appropriate, for national context, (without prejudice to the sovereign rights of nations to determine taxation among other policies), actions to achieve the nine voluntary global targets for NCD prevention and control through the six objectives of the WHO Global NCD Action Plan 2013-2030. The list of interventions is not exhaustive but is intended to provide information and guidance on effectiveness and cost-effectiveness of population-based and individual interventions based on current evidence, and to act as the basis for future work to develop and expand the evidence base.

Why update appendix 3

Appendix 3 is updated to;

- Take into consideration the emergence of new evidence of cost-effectiveness or new WHO recommendations since the adoption of the Global Action Plan for the prevention and control of noncommunicable diseases 2013;
- To accelerate progress towards meeting the nine voluntary global NCD targets and SDG target 3.4, and support the implementation road map 2023-2030 of the Global NCD Action Plan and the Obesity acceleration plan¹², and
- To refine the existing formulation of some interventions based on lessons learnt from the use of the previous two versions.

The area of Appendix 3 where this is most relevant is under Objective 3 (risk factors) and Objective 4 (health systems). All of the “very cost-effective and affordable interventions for all Member States” interventions in the original Appendices were listed under Objectives 3 and 4, and this remains the case in the updated version.

What has changed?

There has been no change to the menu of options listed for Objectives 1 (raising the priority of NCDs), 2 (strengthening leadership & governance), 5 (research) and 6 (monitoring & evaluation) which are process-related recommendations.

¹² WHO (2022) Follow-up to the political declaration of the third high-level meeting of the General Assembly on the prevention and control of non-communicable diseases. Acceleration plan to support Member States in implementing the recommendations for the prevention and management of obesity over the life course. Annex 12. Document A75/10 Add.6. https://apps.who.int/gb/ebwha/pdf_files/WHA75/A75_10Add6-en.pdf

Within Objectives 3 (risk factors) and 4 (health systems), in the 2017 appendix there were 16 interventions considered with a cost-effectiveness ratio $\leq$ \$100 per DALY averted¹³ in low -and middle-income countries (referred to as the “best buys”) and 20 interventions with a cost-effectiveness ratio $>$ \$100/DALY (previously referred to as “good buys”). In addition, 36 interventions without CEA but are part of WHO guidance were also provided. In the 2022 updated Appendix 3, there are now a total of 83 interventions and 21 overarching/enabling actions, representing an expansion from the original list of 88 interventions (including overarching/enabling actions) (Table 1).

In the current update, the cost-effectiveness was examined for 58 interventions. The increase in the number of interventions between 2017 and 2022 updates is due to the availability of new scientific evidence or WHO recommendations as proposed by WHO technical units and/or expert groups since the adoption of the WHO Global NCD Action Plan 2013-2020. In addition, 25 interventions that are part of WHO guidance were also included but without WHO-CHOICE analysis. The absence of cost-effectiveness does not mean that the intervention is not cost-effective, affordable or feasible, but the WHO-CHOICE analysis could not be completed in the 2022 update due to methodological or capacity reasons.

Table 1: Overarching/enabling actions and interventions included in the 2022 update of Appendix 3

Interventions	Overarching/enabling actions	Interventions with WHO-CHOICE analysis	Interventions without WHO-CHOICE analysis
Objective 3			
Tobacco	3	7	2
Harmful use of alcohol	4	5	6
Healthy diet	1	7	4
Physical inactivity	5	2	5
Objective 4			
Cardio-vascular diseases	8	13	5
Diabetes		6	
Chronic respiratory diseases		4	3
Cancer		14	
Total	21	58	25

Table 2 presents the list of interventions for each of the four risk factors and disease areas by income category: low-income countries, lower-middle income countries and upper-middle-income countries. Table 3 ranks all interventions for low and lower-middle income countries by risk factor and disease area, with interventions with an average cost-effectiveness ratio $\leq$ \$100per health life year gained (HLYg) considered

¹³ The international dollar is a hypothetical unit of currency that has the same purchasing power parity that the U.S. dollar had in the United States at a given point in time.

to be the most cost-effective and feasible for implementation in all countries, i.e. the “Best Buys”. Interventions with an average cost-effectiveness ratio > I\$ 100 are listed next and may be considered depending on the country context.

How to use this information

The economic analyses give an assessment of cost-effectiveness, based on the health impact and the economic cost of the intervention. These economic results present a set of parameters for consideration by Member States, however it must be emphasised that such global analyses should be accompanied by analyses further tailored to the local context.

Figure 1 presents the percentage of interventions for which we conducted WHO-CHOICE analysis falling within each band of cost-effectiveness ratio for low-income, lower-middle income and upper-middle income countries. The cost-effectiveness threshold represents the maximum amount one is willing to pay per health outcome. The figure shows that with a threshold of <I\$100 per healthy life year gained 56%, 44% and 33% of the interventions are considered very good value-for-money in low-income, lower-middle income and upper-middle income countries respectively. As this threshold increases, the proportion of interventions considered good value-for-money increases too. If the cost-effectiveness threshold chosen by an upper-middle income country is for example I\$5,000 per healthy life year gained, 86% of interventions in the 2022 update of the Appendix 3 would represent good value for money.


When considering interventions for prevention and management of any disease, including non-communicable diseases, emphasis should be given to both economic and non-economic criteria, as both will affect the implementation and impact of interventions. Non-economic implementation considerations such as acceptability, sustainability, scalability, equity, ethics¹⁴ are essential to consider as part of the prioritisation and implementation of the proposed interventions, based on specific country context.

The WHO secretariat will consider the development of an interactive web-based tool, to help countries see the impact on NCD targets of prioritizing and scaling up the implementation of a set of cost-effective interventions of the updated Appendix 3. Other tools, such as the One Health Tool are available to help individual countries cost specific interventions in their national context¹⁵.

¹⁴ WHO (2021) Principles of health benefit packages. <https://www.who.int/publications/i/item/9789240020689>

¹⁵ <https://www.avenirhealth.org/software-onehealth.php>

Figure 1: Percentage of interventions by cost-effectiveness band for the 3 income groups



Methodological annex

The update of appendix 3 is accompanied by a methodological annex presented at the end of the document. This annex provides more detailed information about the methodology used to identify and analyse interventions, and includes the assumptions used in the WHO-CHOICE economic modelling. The methodological Annex also presents more detailed economic analyses for each intervention, and includes summary tables with costs, health impacts and banded cost-effectiveness ratios for all interventions presented separately for 3 income categories: low-income countries, lower-middle income countries and upper-middle income countries.

Objective 1: To raise the priority accorded to the prevention and control of noncommunicable diseases in global, regional and national agendas and internationally agreed development goals, through strengthened international cooperation and advocacy

Overarching/enabling actions

- Raise public and political awareness, understanding and practice about prevention and control of NCDs
- Integrate NCDs into the social and development agenda and poverty alleviation strategies
- Strengthen international cooperation for resource mobilization, capacity-building, health workforce training and exchange of information on lessons learnt and best practices
- Engage and mobilize civil society and the private sector as appropriate and strengthen international cooperation to support implementation of the action plan at global, regional and national levels
- Implement other policy options in objective 1

Objective 2: To strengthen national capacity, leadership, governance, multisectoral action and partnerships to accelerate country response for the prevention and control of noncommunicable diseases

Overarching/enabling actions

- Prioritize and increase, as needed, budgetary allocations for prevention and control of NCDs without prejudice to the sovereign right of nations to determine taxation and other policies
- Assess national capacity for prevention and control of NCDs
- Develop and implement a national multisectoral policy and plan for the prevention of control of NCDs through multi-stakeholder engagement
- Implement other policy options in objective 2 to strengthen national capacity including human and institutional capacity, leadership, governance, multisectoral action and partnerships for prevention and control of noncommunicable diseases

Objective 3: To reduce modifiable risk factors for noncommunicable diseases and underlying social determinants through creation of health-promoting environments

Tobacco use

Overarching/enabling actions

For the Parties to the WHO Framework Convention on Tobacco Control (WHO FCTC):
- Strengthen the effective implementation of the WHO FCTC and its guidelines, as well as the Protocol to Eliminate Illicit Trade in Tobacco Products, if applicable - Establish and operationalize national mechanisms for coordination of the WHO FCTC implementation as part of national strategy with specific mandate, responsibilities and resources
For the Member States that are not Parties to the WHO FCTC:
Consider implementing the measures set out in the WHO FCTC and its guidelines, as well as the Protocol to Eliminate Illicit Trade in Tobacco Products, if applicable, as the foundational instruments in global tobacco control

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
T1	Increase excise taxes and prices on tobacco products	
T2	Implement plain/standardized packaging and/or large graphic health warnings on all tobacco packages	Requires capacity for implementing and enforcing regulations and legislation
T3	Enact and enforce comprehensive bans on tobacco advertising, promotion and sponsorship	
T4	Eliminate exposure to second-hand tobacco smoke in all indoor workplaces, public places, public transport	
T5	Implement effective mass media campaigns that educate the public about the harms of smoking/tobacco use and second hand smoke 	
T6	Provision of cost-covered effective population-wide support (including brief advice, national toll-free quit line services and mCessation) for tobacco cessation to all tobacco users. 	Requires sufficient, trained providers and a better functioning health system
T7	Provision of cost-covered effective pharmacological interventions to all tobacco users who want to quit. 	

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
T8	Establish a tracking and tracing system to support the elimination of illicit trade in tobacco products 
T9	Ban cross-border tobacco advertising, promotion and sponsorship, including through modern means of communication 

Harmful use of alcohol

Overarching/enabling actions

- Implement the WHO Global strategy to reduce harmful use of alcohol through multisectoral actions in the recommended target areas¹⁶
- Implement the action plan on alcohol to support and complement policy measures and interventions implemented at the national level in the 10 areas recommended of the global strategy¹⁷
- Strengthen leadership and increase commitment and capacity to address the harmful use of alcohol
- Increase awareness and strengthen the knowledge base on the magnitude and nature of problems caused by harmful use of alcohol by awareness programmes, operational research, improved monitoring and surveillance systems

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
A1	Increase excise taxes on alcoholic beverages 	Levying taxes should be combined with other price measures, such as bans on discounts or promotions
A2	Enact and enforce bans or comprehensive restrictions on exposure to alcohol advertising (across multiple types of media) 	Requires capacity and infrastructure for implementing and enforcing regulations and legislation
A3	Enact and enforce restrictions on the physical availability of retailed alcohol (via reduced hours of sale) 	
A4	Enact and enforce drink-driving laws and blood alcohol concentration limits via sobriety checkpoints 	
A5	Provide brief psychosocial intervention for persons with hazardous and harmful alcohol use 	Requires trained providers at all levels of health care

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
A6	Carry out regular reviews of prices in relation to level of inflation and income 
A7	Establish minimum prices for alcohol where applicable 
A8	Enact and enforce an appropriate minimum age for purchase or consumption of alcoholic beverages and reduce density of retail outlets 
A9	Restrict or ban promotions of alcoholic beverages in connection with sponsorships and activities targeting young people
A10	Provide prevention, treatment and care for alcohol use disorders and comorbid conditions in health and social services 
A11	Provide consumer information about, and label, alcoholic beverages to indicate, the harm related to alcohol 

¹⁶ http://www.who.int/substance_abuse/publications/global_strategy_reduce_harmful_use_alcohol/en/

¹⁷ https://apps.who.int/gb/ebwha/pdf_files/EB150/B150_7Add1-en.pdf

Unhealthy diet

Overarching/enabling actions

- Implement the WHO Global Strategy on Diet, Physical Activity and Health¹⁸

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
H1	Reformulation policies for healthier food and beverage products	Requires multisectoral actions with relevant ministries and support by civil society. Regulatory capacity along with multisectoral action is needed.
H2	Front-of-pack labelling as part of comprehensive nutrition labelling policies 	
H3	Public food procurement and service policies for healthy diets 	
H4	Behaviour change communication and mass media campaign for healthy diets 	
H5	Policies to protect children from the harmful impact of food marketing 	
H6	Protection, promotion and support of optimal breastfeeding practices 	
H7	Taxation on sugar-sweetened beverages as part of comprehensive fiscal policies to promote healthy diets 	

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
H7	Subsidies on healthy foods and beverages (e.g. fruits and vegetables) as part of comprehensive fiscal policies for healthy diets 
H8	Menu labelling in food service to promote healthy diets (e.g. reduce total energy intake (kcal) and/or intake of sugars, sodium and unhealthy fats) 
H9	Limiting portion and package size to reduce energy intake and the risk of overweight/obesity
H10	Nutrition education and counselling in different settings (for example, in preschools, schools, workplaces and hospitals) to promote healthy diets 

¹⁸ WHO (2014) Global strategy on diet, physical activity and health. Geneva: World Health Organization.
<https://www.who.int/publications/i/item/9241592222>

Physical inactivity

Overarching/enabling actions

- Global action plan on physical activity 2018–2030: more active people for a healthier world ¹⁹
- ACTIVE: a technical package for increasing physical activity²⁰
- WHO guidelines on physical activity and sedentary behaviour²¹
- Leadership and whole of government commitment to address physical inactivity using a life course approach
- Strong advocacy to increase awareness and knowledge on the cross cutting benefits of increasing physical activity, operational research and knowledge translation and improved monitoring and surveillance systems

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
P1	Provide physical activity assessment, counselling, and behaviour change support as part of routine primary health care services through the use of a brief intervention 	Requires capacity, and sufficiently trained staff in primary care
P2	Implement sustained, population wide, best practice communication campaigns to promote physical activity, with links to community-based programmes and environmental improvements to enable and support behaviour change. 	Requires multisectoral actions with relevant ministries and support by civil society

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
P3	Implement urban and transport planning and urban design, at all levels of government, to provide compact neighbourhoods providing mixed-land use and connected networks for walking and cycling and equitable access to safe, quality public open spaces that enable and promote physical activity and active mobility 
P4	Implement whole-of-school programmes that include quality physical education, and adequate facilities, equipment and programs supporting active travel to/from school and support physical activity for all children of all abilities during and after school 
P5	Improve walking and cycling infrastructure ensuring universal and equitable access to enable and promote safe walking, cycling, other forms of micro mobility (e.g. wheelchairs, scooters and skates) by people of all ages and abilities 
P6	Implement multi-component workplace physical activity programmes 
P7	Provide and promote physical activity through provision of community-based (grass roots) sport and recreation programmes and conduct free mass participation events to encourage engagement by people of all ages and abilities 

¹⁹ <https://apps.who.int/iris/bitstream/handle/10665/272722/9789241514187-eng.pdf>

²⁰ <https://apps.who.int/iris/handle/10665/275415>

²¹ <https://www.who.int/publications/i/item/9789240015128>

Objective 4: To strengthen and orient health systems to address the prevention and control of noncommunicable diseases and the underlying social determinants through people-centred primary health care and universal health coverage

Overarching/enabling actions

- Integrate very cost-effective noncommunicable disease interventions into the basic primary health care package with referral systems to all levels of care to advance the universal health coverage agenda
- Explore viable health financing mechanisms and innovative economic tools supported by evidence
- Scale up early detection and coverage, prioritizing very cost-effective high-impact interventions including cost-effective interventions to address behavioural risk factors
- Train the health workforce and strengthen capacity of health system particularly at primary care level to address the prevention and control of noncommunicable diseases
- Improve the availability of the affordable basic technologies and essential medicines, including generics, required to treat major noncommunicable diseases, in both public and private facilities
- Implement other cost-effective interventions and policy options in objective 4 to strengthen and orient health systems to address noncommunicable diseases and risk factors through people-centered health care and universal health coverage
- Develop and implement a palliative care policy, including access to opioids analgesics for pain relief, together with training for health workers
- Expand the use of digital technologies to increase health service access and efficacy for NCD prevention, and to reduce the costs in health care delivery

Cardiovascular disease

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
CV1	Pharmacological treatment of hypertension in adults using either of the following: thiazide and thiazide-like agents; angiotensin-converting enzyme inhibitors (ACE-Is)/angiotensin-receptor blocker (ARBs); calcium channel blockers (CCBs). 	Feasible to implement in all settings and aligned to latest WHO guidelines (2021) Simple protocols can be followed by non-physician workers depending on the country context
CV2a	Drug therapy (treatment with an antihypertensive and statin) to control CVD risk using a total risk approach and counselling to individuals who have had a heart attack or stroke and to persons with high risk ($\geq 20\%$) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts 	Feasibility and practicality of implementation needs to be assessed and determined. Glucose control not included in this intervention, but in D5 - Control of blood pressure in people with diabetes
CV2b	Drug therapy (treatment with an antihypertensive) to control CVD risk using a total risk approach and counselling to individuals who have had a heart attack or stroke and to persons with high risk ($\geq 10\%$) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts 	Feasibility and practicality of implementation needs to be assessed and determined. Glucose control not included in this intervention, but in D5- Control of blood pressure in people with diabetes.
CV3	Treatment of new cases of acute myocardial infarction with either: acetylsalicylic acid, or acetylsalicylic acid and thrombolysis, or acetylsalicylic acid, thrombolysis and clopidogrel, or primary percutaneous coronary interventions (PCI) initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	Selection of option depends on health system capacity
CV4a	Treatment of acute ischemic stroke with intravenous thrombolytic therapy	Feasibility and practicality of implementation needs to be assessed and determined according to health systems capacity.
CV4b	Treatment of acute ischemic stroke with mechanical thrombectomy within an experienced facility 	Feasibility and practicality of implementation needs to be assessed and determined according to health systems capacity. Requires a surgical facility with appropriately trained workforce.
CV5a	Primary prevention of rheumatic fever and rheumatic heart diseases by increasing appropriate treatment of streptococcal pharyngitis at the primary care level 	

CV5b	Secondary prevention of rheumatic fever and rheumatic heart disease by developing a register of patients who receive regular prophylactic penicillin 	
CV6	Low-dose acetylsalicylic acid within 24 to 48 hours for secondary prevention of ischemic stroke 	
CV7	Comprehensive* care of acute stroke patients in stroke units  *Comprehensive care includes strategies such as staffing by a specialist stroke multidisciplinary team, access to equipment for monitoring and rehabilitation.	Early multidisciplinary approach to be determined and depending on country context. Composition of rehabilitation workforce as an integral part of multidisciplinary team depends on health system capacity.'

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
CV8	Treatment of congestive cardiac failure with angiotensin-converting-enzyme inhibitor, beta-blocker and diuretic 
CV9	Cardiac rehabilitation post myocardial infarction 
CV10	Anticoagulation for medium-and high-risk non-valvular atrial fibrillation and for mitral stenosis with atrial fibrillation 
CV11	Treatment of hypertension using single pill combination anti-hypertensives 
CV12	Secondary prevention of coronary heart disease with a statin, angiotensin-converting-enzyme-inhibitor (ACE-I), beta-blocker and acetylsalicylic acid (low dose) 

Diabetes

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
D1	Foot care to prevent amputation in people with diabetes (including educational programmes, access to appropriate footwear, multidisciplinary clinics) 	
D2	Diabetic retinopathy screening for all diabetes patients and laser photocoagulation for prevention of blindness 	Requires health staff capacity for retinal assessment and photocoagulation
D3	Glycaemic control for people with diabetes, along with standard home glucose monitoring for people treated with insulin to reduce diabetes complications 	
D4	Screening of people with diabetes for proteinuria and treatment with angiotensin-converting enzyme inhibitor for the prevention and delay of renal disease 	
D5	Control of blood pressure in people with diabetes 	
D6	Statin use in people with diabetes > 40years old 	

Chronic respiratory diseases

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
CR1	Acute treatment of asthma exacerbations with inhaled bronchodilators and oral steroids	Requires trained providers at all levels of healthcare 
CR2	Acute treatment of COPD exacerbations with inhaled bronchodilators and oral steroids	
CR3	Long-term management of asthma with inhaled bronchodilator and low-dose beclometasone	
CR4	Long-term management of COPD with inhaled bronchodilator	

Other interventions from WHO guidance (without WHO-CHOICE analysis)

N°	Intervention
CR5	Influenza vaccination for patients with chronic obstructive pulmonary disease 
CR6	Access to improved stoves and cleaner fuels to reduce indoor air pollution 
CR7	Cost-effective interventions to prevent occupational lung diseases, for example, from exposure to silica, asbestos

Cancer

Specific interventions with WHO-CHOICE analysis

N°	Intervention	Non-financial considerations
CA1	Vaccination against human papillomavirus (1-2 doses) of 9–14-year-old girls	
CA2	Cervical cancer: HPV DNA screening, starting at the age of 30 years with regular screening every 5 to 10 years (using a screen-and-treat approach or screen, triage and treat approach)	
CA3	Cervical cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	
CA4	Breast cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	
CA5	Breast Cancer: Screening with mammography (once every 2 years for women aged 50-69 years) linked with timely diagnostic work-up and comprehensive breast cancer treatment in setting where mammographic screening programme is recommended	Requires systems for organised, population-based screening 
CA6	Colorectal cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	
CA7	Basic palliative care for cancer: home-based and hospital care with multi-disciplinary team and access to opiates and essential supportive medicines	Requires access to controlled medicines for pain relief
CA8	Prevention of liver cancer through hepatitis B immunization	
CA9	Oral Cancer: screening in high-risk groups linked with timely diagnostic work-up and comprehensive cancer treatment in setting where significant disease burden and programme is recommended	Requires systems for organised, population-based screening
CA10	Colorectal cancer screening: population-based programme, including through stool-based tests, as appropriate, at age >50 years, linked with timely treatment in settings where screening programme is recommended	Requires systems for organised, population-based screening
CA11	Childhood cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment, focusing on 6 index cancers of WHO Global Initiative for Childhood Cancer	
CA12	Head and neck cancers including oral cancers: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	
CA13	Prostate cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	
CA14	Early detection and comprehensive treatment of cancer for those living with HIV	

Objective 5: To promote and support national capacity for high-quality research and development for the prevention and control of noncommunicable diseases

Overarching/enabling actions

- Develop and implement a prioritized national research agenda for noncommunicable diseases
- Prioritize budgetary allocation for research on noncommunicable disease prevention and control
- Strengthen human resources and institutional capacity for research
- Strengthen research capacity through cooperation with foreign and domestic research institutes
- Implement other policy options in objective 5 to promote and support national capacity for high-quality research, development and innovation

Objective 6: To monitor the trends and determinants of noncommunicable diseases and evaluate progress in their prevention and control

Overarching/enabling actions

- Develop national targets and indicators based on global monitoring framework and linked with a multisectoral policy and plan
- Strengthen human resources and institutional capacity for surveillance and monitoring and evaluation
- Establish and or strengthen a comprehensive noncommunicable disease surveillance system, including reliable registration of deaths by cause, cancer registration, periodic data collection on risk factors and monitoring national response
- Integrate noncommunicable disease surveillance and monitoring into national health information systems
- Implement other policy options in objective 6 to monitor trends and determinants of noncommunicable diseases and evaluate progress in their prevention and control

Methodological annex to Appendix 3: proposed table of contents

The methodological annex will provide further information on the methods used in the development of the 2022 update of the Appendix 3, and highlight the differences compared to the 2017 update. Contents of the Annex will include;

1. Detailed methodological information, including:
 - a. The methodological approach and process used to update appendix 3, including selection of interventions
 - b. Explanation of Generalized Cost Effectiveness Analysis and WHO-CHOICE methodology
 - c. References to the evidence sources and assumptions used in the economic modelling for specific interventions
2. Results of the WHO-CHOICE economic analyses for interventions included in Appendix 3

Detailed methodological information

Identification of interventions

The following criteria used for identifying interventions in 2017 were applied for the 2022 update:

- An intervention must have a demonstrated and quantifiable effect size, from at least one published study in a peer reviewed journal;
- An intervention must have a clear link to one of the global NCD targets.

Additional interventions were considered using the same criteria as above. The intervention list for the 2022 updated Appendix 3 comprised (i) interventions which have been unchanged from the 2017 update, (ii) interventions from the 2017 update which have been re-worded or revised to reflect updates in WHO policy or scientific evidence (iii) interventions included in the 2017 update that had no analysis carried out at the time and for which cost-effectiveness analysis was done in the 2022 update, and (iv) new interventions from new WHO guidance and tools (Table 4).

Table 4: List of interventions for 2022 update and comparison with 2017 update

Interventions	Update of original 2017 interventions (i)	Revised 2017 interventions (ii)	2017 intervention with new CEA (iii)	New interventions with CEA (iv)
Objective 3				
Tobacco	6			1
Harmful use of alcohol	5			
Healthy diet	1	4	2	
Physical inactivity	2			

Objective 4				
Cardio-vascular diseases	7	2	2	2
Diabetes	3		1	2
Chronic respiratory diseases		3		1
Cancer	7		3	4
Total	31	9	8	10

CEA: cost-effectiveness analysis

Choice of economic parameters

Economic parameters are listed below:

1. A menu of interventions for NCDs based on WHO-CHOICE methodology (see below) and presented as a ratio of international dollars (I\$) per healthy life year gained (HLYg)²².
 - a. Cost-effectiveness ratios are presented in bands, ranging from < I\$100/HLYg to >I\$50,000/HLYg²³. The decision to band cost-effectiveness was based on the consensus that the data represent global estimates, therefore banding cost-effectiveness emphasises the relative magnitude of cost-effectiveness rather than a specific amount.
2. Size of health gain: the expected size of population health impact for each intervention was calculated based on total HLY gained per year in a standardized²⁴ population of 1 million people. HLY gained due to an intervention are calculated over a 100-year time frame and evaluated at 95% coverage.
3. Economic cost of implementation: The total cost required per year to implement each intervention was estimated, based on cost in I\$ millions to implement in a standardized population of 1 million people (i.e. I\$ per capita)

While the same methodology has been used for the assessment of the cost-effectiveness of the interventions ensuring comparability of results across areas, different modelling frameworks and assumptions have been used for the modelling of each risk factor and disease. Detailed information on methods, the evidence and assumptions underlying the interventions by disease and risk factor area are provided in separate technical briefs. A comparison across risk factors and diseases will be carried out to check consistency and differences will be explained in a separate document. The evidence used for the modelling of interventions will be periodically revised and updated, and changes in the estimates may occur in the future.

²² The HLY is the equivalent of the Disability Adjusted Life Years (DALYs) used in the 2017 update but framed in a positive way. DALYs are averted while HLY are gained.

²³ Cost-effectiveness bands (in I\$) are: <100, 100-\$500, 500-1,000, 1,000-5,000, 5,000-10,000, 10,000-20,000, 20,000-50,000, >50,000.

²⁴ Standardized over the total population of the analysed countries per income grouping

Country selection

Economic parameters were assessed for 3 country income-groups: low-income countries, lower-middle income countries and upper-middle income countries and this is different compared to the 2017 update where results were presented for 2 country income groups (low -and lower middle-income countries as one group and upper-middle- and high-income countries as the second group). The current update includes selected low -and middle-income countries (LMICs) only since the results from the updated Appendix 3 are most relevant for these countries. Countries were selected so that a significant proportion of the total population and health burden would be represented.

Sixty-two countries were considered in the analysis and listed in Table 5 below²⁵. The list of countries included all low-income countries, the 20 most populous lower-middle-income countries and the 20 most populous upper-middle-income countries. Combined, they represent nearly 80% of the total population and 80% of the global burden of disease. For some type of interventions (e.g. harmful use of alcohol), the analysis was based on a smaller subset of countries.

Table 5: Classification of countries by World Bank income level (2019)

Low-income countries	Lower middle-income countries	Upper middle-income countries ²⁶
Afghanistan	Algeria	Argentina
Burkina Faso	Angola	Brazil
Burundi	Bangladesh	China
Central African Republic	Cameroon	Colombia
Chad	Côte d'Ivoire	Dominican Republic
Democratic Republic of the Congo	Egypt	Ecuador
Eritrea	Ghana	Guatemala
Ethiopia	India	Indonesia
Gambia	Kenya	Iran (Islamic Republic of)
Guinea	Morocco	Iraq
Guinea-Bissau	Myanmar	Jordan
Haiti	Nepal	Kazakhstan
Madagascar	Nigeria	Malaysia
Malawi	Pakistan	Mexico
Mali	Philippines	Peru
Mozambique	Sri Lanka	Russian Federation
Niger	Ukraine	South Africa
Rwanda	United Republic of Tanzania	Thailand
Sierra Leone	Uzbekistan	Turkey
Sudan	Viet Nam	
Tajikistan		
Togo		
Uganda		

²⁵ The 2017 update included 20 countries.

²⁶ One upper-middle income country was excluded during analysis due to lack of data

WHO-CHOICE: A brief methodological overview

General approach

Value for money and efficiency are fundamental considerations guiding investment in health, and WHO-CHOICE provides a way to measure them. This is true in settings where lack of finance is no longer the greatest barrier to achieving better health outcomes; it is also true in less well-resourced settings, where inefficiency is measured in lives lost and human suffering. Cost-effectiveness analysis supports priority setting by defining areas of action where the greatest health gains can be achieved.

Generalized cost-effectiveness analysis (WHO-CHOICE) also allows the definition of an optimal set of interventions, taking into account setting-specific factors such as the burden of disease, health system practice, and economic conditions. Tools to facilitate country-level cost-effectiveness analysis of a wide range of health activities are available. In parallel, WHO-CHOICE publishes and disseminates online a knowledge base of regional-level cost-effectiveness information²⁷.

The use of cost-effectiveness analysis within decision making processes in health is increasingly common globally. However, a series of methodological shortcomings may limit the practical application of cost-effectiveness analysis results. Two examples of this are methodological differences between studies that limit comparability, and use of the current practice as a comparator, which implicitly assumes current resource use is efficient. Generalized Cost-Effectiveness Analysis (GCEA) was developed to overcome such shortcomings of traditional cost-effectiveness analysis.

The GCEA approach enables both existing and new interventions to be evaluated simultaneously. The comparator used in GCEA is a hypothetical “null” scenario, where the impacts of all currently implemented interventions are removed. Uniquely, this method allows existing and new interventions to be analysed at the same time.

Previous cost-effectiveness analyses have been restricted to assessing the efficiency of adding a single new intervention to the existing set or replacing one existing intervention with an alternative. Using WHO-CHOICE, the analyst is no longer constrained by what is already being done, and policymakers can revisit and revise past choices if necessary and feasible. They will have a rational basis for deciding to reallocate resources between interventions to achieve social objectives.

WHO-CHOICE:

- Uses a standardized method for cost-effectiveness analysis that can be applied to all interventions in different settings
- All interventions are evaluated compared to the “null”, a scenario in which we model the absence of health care interventions

²⁷ World Health Organization Choosing Interventions that are Cost-Effective (WHO-CHOICE) programme:
<https://www.who.int/teams/health-systems-governance-and-financing/economic-analysis>

- Impact models are developed using a population-based approach, and healthy life years gained due to an intervention are calculated over a 100-year time frame. Health impacts are not discounted.
- Costs for each intervention are developed using an ingredient based economic costing methodology. Costs are expressed in international dollars (I\$) to ensure comparability across countries and country income groups. Costs are calculated over a 100-year time frame and discounted at 3% per year.
- All interventions are evaluated at 95% coverage.

Table 2: Summary of WHO-CHOICE economic analyses for interventions for NCD prevention and control

Notes: The tables below list all interventions for which WHO-CHOICE economic analysis is available. In this example, separate tables are provided for each of the four main risk factors and four main diseases covered by Objectives 3 and 4 of the WHO Global NCD Action Plan. The intervention name contains more exact detail about the intervention that was modelled -note that these may differ slightly from the wording of the WHO recommended interventions. This Annex is provided for background scientific information only and should not be used as a specific menu for implementation.

		Low-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
Tobacco use										
T1	Increase excise taxes and prices on tobacco products	<100	1,717	<0.01	<100	2,674	0.01	<100	3,093	0.01
T2	Implement plain/standardized packaging and/or large graphic health warnings on all tobacco packages	<100	1,212	<0.01	<100	2,181	0.01	<100	2,319	0.01
T3	Enact and enforce comprehensive bans on tobacco advertising, promotion and sponsorship	<100	933	<0.01	<100	1,490	0.01	<100	1,535	0.01
T4	Eliminate exposure to second-hand tobacco smoke in all indoor workplaces, public places, public transport	<100	1,475	<0.01	<100	2,845	0.01	<100	3,077	0.02
T5	Implement effective mass media campaigns that educate the public about the harms of smoking/tobacco use and second hand smoke	<100	1,345	0.01	<100	2,516	0.03	<100	2,701	0.10
T6	Provision of cost-covered effective population-wide support (including brief advice, national toll-free quit line services and mCessation) for tobacco cessation to all tobacco users.	<100	1,028	0.02	<100	1,724	0.05	<100	1,800	0.09
T7	Provision of cost-covered effective pharmacological interventions to all tobacco users who want to quit.	<100	1,761	0.08	100 – 500	3,583	0.71	100 – 500	3,924	1.32

		Low-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
Harmful use of alcohol										
A1	Increase excise taxes on alcoholic beverages	<100	155	0.01	<100	445	0.02	<100	797	0.04
A2	Enact and enforce bans or comprehensive restrictions on exposure to alcohol advertising (across multiple types of media)	<100	121	<0.01	<100	377	0.01	100 - 500	158	0.03
A3	Enact and enforce restrictions on the physical availability of retailed alcohol (via reduced hours of sale)	<100	132	0.01	<100	383	0.03	100 - 500	190	0.05
A4	Enact and enforce drink-driving laws and blood alcohol concentration limits via sobriety checkpoints	100 - 500	104	0.01	100 - 500	321	0.04	1,000 - 5,000	44	0.11
A5	Provide brief psychosocial intervention for persons with hazardous and harmful alcohol use	<100	244	0.02	100 - 500	647	0.07	100 - 500	691	0.29
Unhealthy diet										
H1	Reformulation policies for healthier food and beverage products	<100	2,449	0.01	<100	4,038	0.02	<100	4,704	0.06
H2	Front-of-pack labelling as part of comprehensive nutrition labelling policies	<100	4,042	<0.01	<100	7,441	0.01	<100	9,069	0.02
H3	Public food procurement and service policies for healthy diets	<100	449	0.01	<100	472	0.02	<100	441	0.04
H4	Behaviour change communication and mass media campaign for healthy diets	<100	2,257	0.01	<100	3,152	0.03	<100	3,506	0.12
H5	Policies to protect children from the harmful impact of food marketing	<100	297	<0.01	<100	446	0.01	<100	610	0.02
H6	Protection, promotion and support of optimal breastfeeding practices	<100	2,052	0.07	<100	3,049	0.11	<100	2,964	0.16

		Low-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
H7	Reduce sugar consumption through effective taxation on sugar-sweetened beverages ²⁸	100 - 500	7	<0.01	100 - 500	12	<0.01	100 - 500	41	<0.01
Physical inactivity										
P1	Brief counselling intervention on physical activity in primary health care	100 - 500	246	0.07	100 - 500	732	0.30	1,000 - 5,000	296	0.90
P2	Physical activity public education and awareness campaign	<100	185	<0.01	<100	617	0.01	100 - 500	162	0.04
Cardiovascular diseases										
CV1	Pharmacological treatment of hypertension in adults using either of the following: thiazide and thiazide-like agents; angiotensin-converting enzyme inhibitors (ACE-Is)/angiotensin-receptor blocker (ARBs); calcium channel blockers (CCBs).	100 - 500	5,281	2.12	500 - 1,000	1,435	0.78	500 - 1,000	3,002	1.97
CV2a	Drug therapy (treatment with an antihypertensive and statin) to control CVD risk using a total risk* approach and counselling to individuals who have had a heart attack or stroke and to persons with high risk (≥ 20%) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts	100 - 500	435	0.21	1,000 - 5,000	274	0.29	500 - 1,000	1,166	0.82
CV2b	Drug therapy (treatment with an antihypertensive) to control CVD risk using a total risk* approach and counselling to individuals who have had a heart attack or stroke and to persons with high risk (≥ 10%) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts	500 - 1,000	2,389	1.24	500 - 1,000	762	0.64	500 - 1,000	2,107	1.79

²⁸ The results on the impact of SSB taxation are preliminary. We are considering additional effects subject to validation.

N°	Intervention	Low-income			Lower-middle-income			Upper-middle-income		
		Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
CV3a	Treatment new cases of acute myocardial infarction with acetylsalicylic acid initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	10,000 - 15,000	7.9	0.08	15,000 - 20,000	2.7	0.04	20,000 - 50,000	3.9	0.08
CV3b	Treatment new cases of acute myocardial infarction with acetylsalicylic acid and thrombolysis initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	5,000 - 10,000	13.0	0.13	15,000 - 20,000	3.7	0.06	15,000 - 20,000	6.1	0.12
CV3c	Treatment of new cases of acute myocardial infarction with acetylsalicylic acid, thrombolysis and clopidogrel, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	5,000 - 10,000	15.4	0.14	15,000 - 20,000	4.2	0.06	15,000 - 20,000	7.2	0.12
CV3d	Treatment of new cases of myocardial infarction with primary percutaneous coronary interventions (PCI), acetylsalicylic acid and clopidogrel, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	>50,000	7.8	1.00	>50,000	2.6	0.24	>50,000	3.9	0.56
CV4a	Treatment of acute ischemic stroke with intravenous thrombolytic therapy	5,000 - 10,000	18.3	0.17	10,000 - 15,000	5.3	0.07	15,000 - 20,000	16.9	0.27
CV4b	Treatment of acute ischemic stroke with mechanical thrombectomy within an experienced facility	20,000 - 50,000	8.0	0.40	20,000 - 50,000	2.9	0.10	>50,000	7.6	0.48
CV5a	Primary prevention of rheumatic fever and rheumatic heart diseases by increasing appropriate treatment of streptococcal pharyngitis at the primary care level	100 - 500	3,430.0	0.44	100 - 500	989.0	0.14	100 - 500	622.4	0.28

		Lower-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
CV5b	Secondary prevention of rheumatic fever and rheumatic heart disease by developing a register of patients who receive regular prophylactic penicillin	<100	722.2	0.03	<100	200.0	0.01	<100	121.9	0.01
CV6	Low-dose acetylsalicylic acid within 24 to 48 hours for secondary prevention of ischemic stroke	20,000 - 50,000	0.8	0.03	10,000 - 15,000	1.2	0.02	>50,000	1.0	0.06
CV7	Comprehensive care of acute stroke patients in stroke units	>50,000	1.6	0.22	>50,000	1.4	0.11	>50,000	1.8	0.39
Diabetes										
D1	Foot care to prevent amputation in people with diabetes (including educational programmes, access to appropriate footwear, multidisciplinary clinics)	100 - 500	80	0.03	1,000 - 5,000	99	0.10	1,000 - 5,000	176	0.20
D2	Diabetic retinopathy screening for all diabetes patients and laser photocoagulation for prevention of blindness	1,000 - 5,000	32	0.04	1,000 - 5,000	48	0.17	1,000 - 5,000	133	0.34
D3	Glycaemic control for people with diabetes, along with standard home glucose monitoring for people treated with insulin to reduce diabetes complications	500 - 1,000	1,091	0.75	1,000 - 5,000	1,996	2.51	1,000 - 5,000	2,784	3.96
D4	Screening of people with diabetes for proteinuria and treatment with angiotensin-converting enzyme inhibitor for the prevention and delay of renal disease	100 - 500	526	0.12	100 - 500	1,012	0.44	500 - 1,000	1,192	0.78
D5	Control of blood pressure in people with diabetes	100 - 500	274	0.04	100 - 500	532	0.17	500 - 1,000	824	0.50
D6	Statin use in people with diabetes > 40years old	<100	515	0.05	100 - 500	1,271	0.22	100 - 500	1,559	0.61

		Low-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
Chronic Respiratory Diseases										
CR1	Acute treatment of asthma exacerbations with inhaled bronchodilators and oral steroids	<100	1,220	0.05	100 - 500	1,051	0.13	500 - 1,000	828	0.43
CR2	Acute treatment of COPD exacerbations with inhaled bronchodilators and oral steroids	<100	183	<0.01	100 - 500	378	0.04	100 - 500	389	0.16
CR3	Long-term management of asthma with inhaled bronchodilator and low-dose beclometasone	<100	1,204	0.10	100 - 500	1,123	0.16	100 - 500	733	0.32
CR4	Long-term management of COPD with inhaled bronchodilator	<100	920	0.03	<100	1,811	0.16	100 - 500	1,723	0.39
Cancer										
CA1	Vaccination against human papillomavirus (1-2 doses) of 9–14 year old girls	<100	2,170	0.04	<100	1,878	0.05	<100	2,524	0.05
CA2	Cervical cancer: HPV DNA screening, starting at the age of 30 years with regular screening every 5 to 10 years (using a screen-and-treat approach or screen, triage and treat approach)	<100	1,012	0.02	<100	940	0.05	<100	1,686	0.14
CA3	Cervical cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	621	0.02	<100	605	0.02	<100	578	0.03
CA4	Breast cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	929	0.05	<100	1,405	0.11	100 - 500	2,721	0.30
CA5	Breast Cancer: Screening with mammography (once every 2 years for women aged 50-69 years) linked with timely diagnostic work-up and comprehensive breast cancer treatment in setting where	100 - 500	990	0.17	100 - 500	1,492	0.39	100 - 500	2,900	0.94

N°	Intervention	Low-income			Lower-middle-income			Upper-middle-income		
		Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
	mammographic screening programme is recommended									
CA6	Colorectal cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	356	0.02	<100	643	0.04	100 - 500	2,877	0.34
CA7	Basic palliative care for cancer: home-based and hospital care with multi-disciplinary team and access to opiates and essential supportive medicines ²⁹			<0.01			<0.01			0.02
CA8	Prevention of liver cancer through hepatitis B immunization	<100	213	0.01	100 - 500	175	0.02	<100	684	0.01
CA9	Oral Cancer: screening in high-risk groups linked with timely diagnostic work-up and comprehensive cancer treatment in setting where significant disease burden and programme is recommended	1,000 - 5,000	43	0.12	100 - 500	1,371	0.38	1,000 - 5,000	227	0.90
CA10	Colorectal cancer screening: population-based programme, including through stool-based tests, as appropriate, at age >50 years, linked with timely treatment in settings where screening programme is recommended	100 - 500	564	0.11	100 - 500	1,036	0.29	100 - 500	4,613	1.13
CA11	Childhood cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment, focusing on 6 index cancers of WHO Global Initiative for Childhood Cancer	<100	262	0.02	<100	1,148	0.10	<100	354	0.03
CA12	Head and neck cancers including oral cancers: early diagnosis programs	500 - 1,000	18	0.01	500 - 1,000	102	0.08	1,000 - 5,000	47	0.05

²⁹ Impact model not available for palliative care

		Low-income			Lower-middle-income			Upper-middle-income		
N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
	linked with timely diagnostic work-up and comprehensive cancer treatment									
CA13	Prostate cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	100 - 500	224	0.03	100 - 500	272	0.05	100 - 500	1,040	0.19
CA14	Early detection and comprehensive treatment of cancer for those living with HIV	<100	247	0.01	<100	84	<0.01	100 - 500	34	<0.01

Table 3: Ranking of interventions by average cost-effectiveness ratio for NCD risk factors and non-communicable diseases based on analysis in low and lower-middle income countries

N°	Intervention	Banded cost-effectiveness ratio	Health impact per year (HLY gained per 1 million)	Economic cost per year (\$ in millions per 1 million)
Tobacco use				
T1	Increase excise taxes and prices on tobacco products	<100	2,400	0.01
T2	Implement plain/standardized packaging and/or large graphic health warnings on all tobacco packages	<100	1,904	0.01
T3	Enact and enforce comprehensive bans on tobacco advertising, promotion and sponsorship	<100	1,331	0.01
T4	Eliminate exposure to second-hand tobacco smoke in all indoor workplaces, public places, public transport	<100	2,453	0.01
T5	Implement effective mass media campaigns that educate the public about the harms of smoking/tobacco use and second hand smoke	<100	2,182	0.02
T6	Provision of cost-covered effective population-wide support (including brief advice, national toll-free quit line services and mCessation) for tobacco cessation to all tobacco users.	<100	1,525	0.04
T7	Provision of cost-covered effective pharmacological interventions to all tobacco users who want to quit.	100 - 500	3,062	0.53
Harmful use of alcohol				
A1	Increase excise taxes on alcoholic beverages	<100	362	0.02
A2	Enact and enforce bans or comprehensive restrictions on exposure to alcohol advertising (across multiple types of media)	<100	304	0.01
A3	Enact and enforce restrictions on the physical availability of retailed alcohol (via reduced hours of sale)	<100	311	0.02
A4	Enact and enforce drink-driving laws and blood alcohol concentration limits via sobriety checkpoints	100 - 500	259	0.03
A5	Provide brief psychosocial intervention for persons with hazardous and harmful alcohol use	100 – 500	532	0.05

Unhealthy diet				
H1	Reformulation policies for healthier food and beverage products	<100	3,584	0.02
H2	Front-of-pack labelling as part of comprehensive nutrition labelling policies	<100	6,470	0.01
H3	Public food procurement and service policies for healthy diets	<100	465	0.02
H4	Behaviour change communication and mass media campaign for healthy diets	<100	2,897	0.03
H5	Policies to protect children from the harmful impact of food marketing	<100	403	0.01
H6	Protection, promotion and support of optimal breastfeeding practices	<100	2,764	0.10
H7	Reduce sugar consumption through effective taxation on sugar-sweetened beverages ³⁰	100 - 500	11	<0.01
Physical inactivity				
P2	Implement sustained, population wide, best practice communication campaigns to promote physical activity, with links to community-based programmes and environmental improvements to enable and support behaviour change.	<100	451	0.01
P1	Provide physical activity assessment, counselling, and behaviour change support as part of routine primary health care services through the use of a brief intervention	500 - 1,000	573	0.38
Cardiovascular diseases				
CV5b	Secondary prevention of rheumatic fever and rheumatic heart disease by developing a register of patients who receive regular prophylactic penicillin	<100	349	0.01
CV1	Pharmacological treatment of hypertension in adults using either of the following: thiazide and thiazide-like agents; angiotensin-converting enzyme inhibitors (ACE-Is)/angiotensin-receptor blocker (ARBs); calcium channel blockers (CCBs).	100 - 500	2,535	1.16
CV5a	Primary prevention of rheumatic fever and rheumatic heart diseases by increasing appropriate treatment of streptococcal pharyngitis at the primary care level	100 - 500	1,687	0.23
CV2a	Drug therapy (treatment with an antihypertensive and statin) to control CVD risk using a total risk* approach and counselling to individuals who have had a heart attack or stroke and to persons with high risk (≥ 20%) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts	500 - 1,000	320	0.27
CV2b	Drug therapy (treatment with an antihypertensive) to control CVD risk using a total risk* approach and counselling to individuals who have had a heart attack or stroke and to	500 - 1,000	1,227	0.81

³⁰ The results on the impact of SSB taxation are preliminary. We are considering additional effects subject to validation

	persons with high risk ($\geq 10\%$) of a fatal and non-fatal cardiovascular event in the next 10 years using the updated WHO CVD risk charts			
CV3a	Treatment new cases of acute myocardial infarction with acetylsalicylic acid, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	10,000 - 15,000	4	0.05
CV3b	Treatment of new cases of acute myocardial infarction with acetylsalicylic acid and thrombolysis, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	10,000 - 15,000	6	0.08
CV3c	Treatment of new cases of acute myocardial infarction with acetylsalicylic acid, thrombolysis and clopidogrel, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	10,000 - 15,000	7	0.09
CV4a	Treatment of acute ischemic stroke with intravenous thrombolytic therapy	10,000 - 15,000	9	0.10
CV6	Low-dose acetylsalicylic acid within 24 to 48 hours for secondary prevention of ischemic stroke	15,000 - 20,000	1	0.02
CV4b	Treatment of acute ischemic stroke with mechanical thrombectomy within an experienced facility	20,000 - 50,000	4	0.19
CV3d	Treatment of new cases of myocardial infarction with primary percutaneous coronary interventions (PCI), acetylsalicylic acid and clopidogrel, initially treated in a hospital setting with follow up carried out through primary health care facilities at a 95% coverage rate	>50,000	4	0.46
CV7	Comprehensive care of acute stroke patients in stroke units	>50,000	1	0.14
Diabetes				
D4	Screening of people with diabetes for proteinuria and treatment with angiotensin-converting enzyme inhibitor for the prevention and delay of renal disease	100 - 500	873	0.35
D5	Control of blood pressure in people with diabetes	100 - 500	459	0.14
D6	Statin use in people with diabetes > 40years old	100 - 500	1,055	0.17
D1	Foot care to prevent amputation in people with diabetes (including educational programmes, access to appropriate footwear, multidisciplinary clinics)	500 - 1,000	93	0.08
D2	Diabetic retinopathy screening for all diabetes patients and laser photocoagulation for prevention of blindness	1,000 - 5,000	44	0.13
D3	Glycaemic control for people with diabetes, along with standard home glucose monitoring for people treated with insulin to reduce diabetes complications	1,000 - 5,000	1,737	2.01
Chronic respiratory diseases				

CR1	Acute treatment of asthma exacerbations with inhaled bronchodilators and oral steroids	<100	1,100	0.11
CR2	Acute treatment of COPD exacerbations with inhaled bronchodilators and oral steroids	<100	322	0.03
CR4	Long-term management of COPD with inhaled bronchodilator	<100	1,556	0.12
CR3	Long-term management of asthma with inhaled bronchodilator and low-dose beclometasone	100 - 500	1,146	0.14
Cancer				
CA1	Vaccination against human papillomavirus (1-2 doses) of 9–14 year old girls	<100	1,962	0.05
CA2	Cervical cancer: HPV DNA screening, starting at the age of 30 years with regular screening every 5 to 10 years (using a screen-and-treat approach or screen, triage and treat approach)	<100	961	0.04
CA3	Cervical cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	610	0.02
CA4	Breast cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	1,269	0.09
CA6	Colorectal cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	<100	561	0.04
CA8	Prevention of liver cancer through hepatitis B immunization	<100	186	0.02
CA11	Childhood cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment, focusing on 6 index cancers of WHO Global Initiative for Childhood Cancer	<100	895	0.08
CA14	Early detection and comprehensive treatment of cancer for those living with HIV	<100	131	<0.01
CA5	Breast Cancer: Screening with mammography (once every 2 years for women aged 50-69 years) linked with timely diagnostic work-up and comprehensive breast cancer treatment in setting where mammographic screening programme is recommended	100 - 500	1,349	0.33
CA9	Oral Cancer: screening in high-risk groups linked with timely diagnostic work-up and comprehensive cancer treatment in setting where significant disease burden and programme is recommended	100 - 500	991	0.31
CA13	Prostate cancer: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	100 - 500	258	0.04

CA10	Colorectal cancer screening: population-based programme, including through stool-based tests, as appropriate, at age >50 years, linked with timely treatment in settings where screening programme is recommended	100 - 500	901	0.24
CA12	Head and neck cancers including oral cancers: early diagnosis programs linked with timely diagnostic work-up and comprehensive cancer treatment	500 - 1,000	78	0.06
CA7	Basic palliative care for cancer: home-based and hospital care with multi-disciplinary team and access to opiates and essential supportive medicines ³¹			<0.01

³¹ Impact model not available for palliative care

WHO Discussion Paper (version dated 1 August 2022)

List of comments by page number

Page: 10

Number: 1

Trabajar más, en medios de comunicación, E INCLUSO EN REDES SOCIALES desalentando mal consumo de tabaco y alcohol también.

Number: 2

Trabajar con adolescentes fomentando la prevención por pares, evitando que adopten estereotipos sobre el consumo de cigarrillos y alcohol.

Number: 3

Considerar el porcentaje de recaídas luego de la intervención medicamentosa. Se podría dar en casos específicos que tengan dependencia de otras drogas que se consuman de modo similar al Tabaco.

Number: 4

Controlar tabaco que llega por contrabando.

Number: 5

Es difícil controlar la publicidad de otros países, hay sitios de frontera donde medios de países vecinos llegan más fácilmente que los nacionales. Considerar si Colombia, Perú y Brasil se unirán a estas estrategias. Se podrá celebrar convenios con Facebook, Instagram etc.? con el fin de limitar publicidad en estas redes sociales?

Page: 11

Number: 1

Es un excelente medio de limitar la venta. tiene la debilidad de no afectar el alcohol artesanal y el ingreso de licores de contrabando.

Number: 2

La debilidad en este punto es la publicidad por redes sociales hacer convenio con Facebook, Instagram etc.

Number: 3

Podría fomentar la venta oculta y contrabando. En Ecuador se evidencia que tiendas de barrio venden licores, e incluso la gente consume licor en ellas o sus alrededores, la Policía no puede actuar sino dando una advertencia. Quizá sería conveniente reformar ciertas leyes para poder emitir multas u otro tipo de sanciones que desanimen el consumo de licor en la vía pública, canchas etc.

Number: 4

Se cumple parcialmente. La Policía y las Agencias de Tránsito tienen limitadas capacidades. La gente reacciona "defendiendo" al infractor de la "agresión" de la autoridad que exige el cumplimiento de la ley.

Number: 5

La ley no permite el ingreso forzoso de una persona con consumo peligroso en un CETAD. No existen CETADs suficientes.

Number: 6

Implicaría focalizar precios

Number: 7

El control en el consumo de alcohol artesanal sería limitado. Una forma de controlar este consumo, sería que se adquiriera el alcohol artesanal directamente al productor en finca, para usarlo en incrementar octanaje de combustibles

Number: 8

Coordinar con Municipios, que son quienes dan los permisos de funcionamiento de los comercios, mediante las LUAEs

Number: 9

La ley no permite el ingreso forzoso de una persona con consumo peligroso en un CETAD. No existen CETADs suficientes.

Number: 10

el aviso de que el alcohol es peligroso para la salud debe ser mas grande, quizá usar imágenes que hagan pensar dos veces en consumirlo - Vehículos accidentados, personas enfermas con cirrosis etc.

Page: 12

Number: 1

No aplicable para comidas tradicionales y comidas artesanales por ejemplo.

Number: 2

Establecer convenios con asociaciones de restaurantes, y patios de comidas populares para disminuir el uso de sal en las preparaciones, y no poner el salero en la mesa a menos que el usuario lo pida específicamente. Limitar los permisos de funcionamiento para locales de comida Rápida??, Endurecer los controles en las fabricas de alimentos procesados?

Number: 3

Limitar la publicidad de Comida chatarra y comidas rápidas

Number: 4

Trabajar desde la academia para que inculquen en sus estudiantes una cultura de alimentación saludable.

Number: 5

Fortalecer la estrategia de fomento lactancia materna

Number: 6

Pocas bebidas comerciales usan azúcar, casi todas usan edulcorantes, sin embargo existen otras industrias que continúan usando jarabe de maíz y otros con alto contenido de glucosa. Controlar la venta de dulces artesanales a base de panela como melcochas etc.

Number: 7

Implementar como una estrategia de fomento a la agricultura orgánica

Number: 8

Trabajar con escolares y adolescentes para eliminar costumbres establecidas Controlar que en los bares escolares no se expendan comida chatarra. No se podría confiar esta responsabilidad a las autoridades de los planteles, por la facilidad de arreglos privados entre interesados.

Number: 9

Estrategia a mediano y largo plazo.

Page: 13

Number: 1

Se requiere personal de salud motivado a prescribir la actividad física. Cambio de mentalidad entre el personal mas antiguo.

Generar diferenciación entre "hacer deporte" y "hacer ejercicio" o "actividad física". Hay costumbres arraigadas (consumo de bebidas alcohólicas luego de "hacer deporte")

Number: 2

Estímulos a los programas que fomenten la actividad física; no necesariamente la practica de deportes

Number: 3

Coordinar con programa de Municipio saludable

Number: 4

Transportarse de casa a escuela en Biciclobuses escolares como existe en Holanda? para ello debe fortalecerse programas de seguridad, y estricto cumplimiento a la zonificación escolar.

Number: 5

En coordinación con los programas de Municipio saludable. Así también debe aplicarse estrictamente la zonificación escolar

Number: 6

En la practica las "pausas activas" no han funcionado, el personal las cumple parcialmente, o no lo hace, lo ven como "Una obligación más". Habría que generar estrategias para motivarlos a hacer actividad física. No necesariamente de índole deportiva, o competitiva.

Number: 7

Actividades de caminata y marcha rápida; antes que actividades competitivas de maratones o carreras 5k 10k etc.

Page: 15

Number: 1

En Ecuador ya se ha implementado la estrategia HEARTS

Number: 2

Existe la GPC de Hipertensión Arterial. Se esta trabajando en adoptar la Guía de tratamiento farmacológico de la OPS.

Number: 3

Existe la GPC de Hipertensión Arterial. Se esta trabajando en Adoptar la Guía de tratamiento farmacológico de la OPS.

Number: 4

Existe la Estrategia Actúa con Velocidad (ACV)

Solo en el Hospital de Especialidades Portoviejo existe el Centro de manejo especializado de enfermedades Cardiovasculares

Number: 5

Los protocolos de manejo varían en las diferentes unidades de salud.

Page: 16

Number: 1

Hasta la fecha, no existe dicho registro

Number: 2

Existe la estrategia Actúa con Velocidad (ACV)

Number: 3

Solo en el Hospital de Especialidades Portoviejo existe el Centro de manejo especializado de enfermedades Cardiovasculares

Number: 4

Los protocolos de manejo vaian en las diferentes unidades de salud. Solo en el Hospital de Especialidades Portoviejo existe el Centro de manejo especializado de enfermedades Cardiovasculares

Number: 5

Los protocolos de manejo vaian en las diferentes unidades de salud.

Number: 6

Los protocolos de manejo vaian en las diferentes unidades de salud. Solo en el Hospital de Especialidades Portoviejo existe el Centro de manejo especializado de enfermedades Cardiovasculares

Number: 7

La combinación de antihipertensivos en una sola píldora se encuentra en proceso de ser incluida en el cuadro de medicamentos básicos

Number: 8

Los protocolos de manejo varían en las diferentes unidades de salud. Solo en el Hospital de Especialidades Portoviejo existe el Centro de manejo especializado de enfermedades Cardiovasculares

Page: 17

Number: 1

En varias unidades del país existen clínicas de Diabetes y Clínicas de Heridas que previenen y tratan pie diabético

Number: 2

Existe la GPC de Diabetes Mellitus

Number: 3

Existe la GPC de Diabetes Mellitus

Number: 4

Existe la GPC de Diabetes Mellitus

Number: 5

Existe la GPC de Diabetes Mellitus y se ha implementado la estrategia HEARTS

Number: 6

Existe la GPC de Diabetes Mellitus

Page: 18

Number: 1

En las unidades de salud se manejan protocolos basados en las guías Global Initiative for Chronic Obstructive Lung Disease (GOLD)

Number: 2

Se cumple este procedimiento

Number: 3

Mas del 97% de las cocinas domesticas y comerciales en Ecuador usan GLP.

Page: 19

Number: 1

Estas vacunas son parte de la Estrategia Nacional de Inmunización

Number: 2

Existe la Estrategia Nacional para la Atencion Integral del Cancer, la cual actualmente esta vigente y se está trabajando en su actualizacion.

Number: 3

Se estan implementando planes piloto para la realizacion de pruebas moleculares para detección de VPH

Number: 4

Existe la Estrategia Nacional para la Atencion Integral del Cancer, la cual actualmente esta vigente y se está trabajando en su actualizacion.

Number: 5

Existe la Estrategia Nacional para la Atencion Integral del Cancer, la cual actualmente esta vigente y se está trabajando en su actualizacion.

Number: 6

La vacuna para Hepatitis B es parte de la ENI, Vacunación Obligatoria dentro de las primeras 24 horas de nacido

Number: 7

Se está trabajando en el Protocolo de Primera Linea para el Manejo de Leucemia linfoblástica Aguda, que es el cáncer mas frecuente en niños

Number 8

Cada unidad de salud que tiene ORL o cirugía de cabeza y cuello, maneja protocolos basados en Guías Internacionales

Number 9

Cada Unidad de salud que cuenta con servicio de Urología, maneja sus protocolos de detección y tratamiento basados en guías internacionales