

Global School-based Student Health Survey



Bolivia

2018 Fact Sheet

The 2018 Bolivia GSHS was a school-based survey of students in 2nd Secondary - 6th Secondary, which are typically attended by students aged 13-17. A two-stage cluster sample design was used to produce data representative of all students in 2nd Secondary - 6th Secondary in Bolivia. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate.

The Bolivia GSHS measured alcohol use; drug use; hygiene; mental health; physical activity; protective factors; sexual behaviors; and violence and unintentional injury. Students self-reported their responses to each question on a computer scanable answer sheet.

The school response rate was 94%, the student response rate was 84%, and the overall response rate was 79%. A total of 7,931 students participated in the Bolivia GSHS. Prevalence estimates (percentages) and 95% confidence intervals are presented below.

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Alcohol Use									
Percentage of students who currently drank alcohol (at least one drink of alcohol on at least one day during the 30 days before the survey)	20.0 (18.0-22.1)	20.4 (17.6-23.4)	19.4 (16.5-22.6)	29.3 (25.6-33.2)	30.0 (26.5-33.7)	28.1 (23.7-32.8)	24.0 (21.8-26.4)	24.6 (22.4-27.0)	23.1 (20.0-26.5)
Percentage of students who ever drank so much alcohol that they were really drunk one or more times during their life	15.3 (13.6-17.1)	15.6 (13.6-17.9)	14.6 (12.1-17.4)	29.2 (25.9-32.7)	32.4 (29.2-35.8)	25.6 (21.4-30.4)	21.3 (19.2-23.5)	23.1 (20.9-25.4)	19.3 (16.4-22.4)
Percentage of students who drank alcohol before age 14 years for the first time, among students who ever had a drink of alcohol other than a few sips	65.8 (62.7-68.9)	71.0 (66.6-75.0)	60.2 (56.6-63.6)	32.6 (29.7-35.7)	39.1 (35.2-43.3)	24.3 (20.5-28.6)	48.5 (45.1-51.8)	54.0 (50.3-57.7)	41.9 (38.2-45.7)
Drug Use									
Percentage of students who ever used marijuana one or more times during their life	6.2 (5.2-7.5)	7.5 (5.9-9.5)	4.8 (3.5-6.4)	10.9 (8.8-13.4)	13.5 (10.6-17.1)	8.1 (6.1-10.7)	8.3 (7.0-9.7)	10.2 (8.3-12.4)	6.2 (5.0-7.6)

Global School-based Student Health Survey



Bolivia

2018 Fact Sheet

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Drug Use									
Percentage of students who used drugs before age 14 years for the first time, among students who ever used drugs	73.1 (65.1-79.8)	74.1 (65.4-81.2)	70.4 (60.0-79.0)	45.9 (39.2-52.7)	45.3 (38.9-51.8)	44.2 (32.4-56.6)	58.6 (51.7-65.2)	58.8 (52.1-65.2)	56.5 (46.2-66.2)
Hygiene									
Percentage of students who usually cleaned or brushed their teeth one or more times per day during the 30 days before the survey	88.9 (87.3-90.3)	86.5 (84.3-88.3)	91.4 (89.8-92.8)	89.4 (88.0-90.6)	88.0 (85.8-89.9)	91.2 (89.9-92.4)	89.1 (88.0-90.1)	87.1 (85.5-88.7)	91.3 (90.5-92.1)
Percentage of students who never or rarely washed their hands after using the toilet or latrine during the 30 days before the survey	10.9 (9.0-13.0)	11.9 (9.7-14.4)	9.7 (7.7-12.0)	10.9 (9.3-12.6)	12.7 (10.4-15.4)	8.6 (7.1-10.4)	10.9 (9.5-12.4)	12.2 (10.4-14.3)	9.2 (7.9-10.8)
Mental Health									
Percentage of students who seriously considered attempting suicide during the 12 months before the survey	22.1 (20.2-24.1)	16.1 (14.3-18.0)	27.6 (24.8-30.6)	22.2 (20.0-24.5)	15.8 (13.6-18.4)	28.8 (26.2-31.5)	22.1 (20.5-23.9)	16.0 (14.6-17.4)	28.1 (26.0-30.3)
Percentage of students who attempted suicide one or more times during the 12 months before the survey	22.0 (19.9-24.1)	17.8 (15.6-20.2)	25.6 (23.0-28.5)	20.8 (18.7-23.1)	15.8 (13.8-18.1)	25.6 (22.7-28.7)	21.5 (19.9-23.1)	16.9 (15.4-18.5)	25.6 (23.5-27.8)
Percentage of students who did not have any close friends	10.0 (8.7-11.5)	10.7 (9.2-12.6)	9.2 (7.5-11.2)	11.0 (9.7-12.3)	10.6 (9.2-12.3)	11.1 (9.2-13.3)	10.4 (9.3-11.7)	10.7 (9.5-12.0)	10.0 (8.5-11.7)

Global School-based Student Health Survey



Bolivia

2018 Fact Sheet

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Physical Activity									
Percentage of students were physically active at least 60 minutes per day on all 7 days during the 7 days before the survey	11.8 (10.4-13.3)	14.6 (12.5-16.9)	9.1 (7.6-10.8)	11.4 (10.1-12.8)	15.0 (12.7-17.6)	7.8 (6.2-9.8)	11.6 (10.6-12.7)	14.8 (13.3-16.4)	8.6 (7.3-10.0)
Percentage of students who attended physical education classes on three or more days each week during this school year	41.0 (38.3-43.8)	41.3 (38.1-44.5)	40.6 (37.3-44.1)	38.2 (35.8-40.6)	39.0 (36.2-42.0)	37.6 (34.1-41.2)	39.8 (37.5-42.1)	40.3 (37.9-42.7)	39.3 (36.5-42.2)
Percentage of students who spent three or more hours per day sitting and watching television, playing computer games, or talking with friends, when not in school or doing homework during a typical or usual day	29.8 (27.0-32.7)	30.1 (27.2-33.2)	29.4 (26.1-32.9)	33.6 (30.7-36.5)	34.1 (30.7-37.7)	32.9 (29.1-37.0)	31.4 (29.1-33.8)	31.9 (29.6-34.2)	30.9 (27.9-34.0)
Protective Factors									
Percentage of students who missed classes or school without permission on one or more days during the 30 days before the survey	35.1 (33.2-37.2)	35.3 (33.0-37.6)	34.8 (31.9-37.8)	43.6 (41.3-45.8)	44.8 (41.7-47.9)	41.7 (38.1-45.3)	38.8 (37.1-40.5)	39.5 (37.4-41.7)	37.7 (35.2-40.2)
Percentage of students who reported that their parents or guardians most of the time or always understood their problems and worries during the 30 days before the survey	31.8 (29.9-33.7)	32.8 (30.1-35.6)	30.9 (28.1-33.9)	28.9 (26.6-31.3)	30.3 (27.2-33.6)	27.5 (24.8-30.4)	30.5 (28.9-32.2)	31.7 (29.2-34.2)	29.5 (27.3-31.7)

Global School-based Student Health Survey



Bolivia

2018 Fact Sheet

	Students Aged 13-15 Years			Students Aged 16-17 Years			Students Aged 13-17 Years		
	Total	Males	Females	Total	Males	Females	Total	Males	Females
Protective Factors									
Percentage of students who reported that their parents or guardians most of the time or always really knew what they were doing with their free time during the 30 days before the survey	38.6 (36.0-41.2)	37.0 (33.3-40.8)	40.1 (37.4-42.9)	36.9 (33.9-40.0)	36.6 (32.6-40.7)	37.6 (34.7-40.5)	37.8 (35.4-40.3)	36.8 (33.3-40.4)	39.0 (37.0-41.1)
Sexual Behaviors									
Percentage of students who ever had sexual intercourse	20.6 (18.4-23.0)	25.0 (21.3-29.2)	16.3 (14.2-18.6)	39.9 (37.1-42.9)	47.6 (44.0-51.2)	32.0 (29.0-35.1)	29.1 (26.7-31.5)	35.2 (31.8-38.7)	23.0 (20.9-25.3)
Percentage of students who had sexual intercourse before age 14 years for the first time, among students who ever had sexual intercourse	55.8 (49.2-62.2)	59.6 (51.5-67.1)	50.0 (40.0-60.0)	21.7 (18.7-25.0)	26.9 (22.8-31.5)	12.4 (9.2-16.4)	34.3 (30.3-38.5)	38.9 (34.1-43.9)	26.6 (21.4-32.5)
Percentage of students who used a condom during last sexual intercourse, among students who ever had sexual intercourse	62.8 (58.1-67.2)	65.3 (59.9-70.4)	57.8 (50.6-64.7)	62.8 (58.7-66.8)	63.6 (58.5-68.5)	61.0 (54.9-66.7)	62.8 (59.4-66.1)	64.2 (60.5-67.8)	59.9 (54.6-64.9)
Violence and Unintentional Injury									
Percentage of students who were in a physical fight one or more times during the 12 months before the survey	30.3 (28.1-32.6)	40.3 (37.6-43.0)	20.4 (18.5-22.4)	30.8 (27.6-34.1)	40.6 (37.0-44.2)	20.1 (17.3-23.2)	30.5 (28.0-33.1)	40.4 (37.8-43.1)	20.3 (18.4-22.3)
Percentage of students who were seriously injured one or more times during the 12 months before the survey	51.5 (49.6-53.5)	55.6 (53.1-58.1)	47.3 (44.2-50.4)	51.9 (49.3-54.6)	56.0 (52.8-59.2)	47.6 (44.3-51.0)	51.7 (49.8-53.6)	55.8 (53.5-58.0)	47.4 (44.9-50.0)

For additional information, please contact:
Yecid Humacayo Morales, Jefe de la Unidad de Alimentación y Nutrición
Ministerio de Salud Bolivia, La Paz, Bolivia,
yhumacayo@minsalud.gob.bo

For general information about the GSHS, please visit:
<http://www.who.int/ncds/surveillance/gshs/>
or
<http://www.cdc.gov/gshs>