

Mexico - Tijuana (Ages 13-15)

Global Youth Tobacco Survey (GYTS)

FACT SHEET



The Mexico - Tijuana GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Mexico - Tijuana could include in a comprehensive tobacco control program.

The Mexico - Tijuana GYTS was a school-based survey of students in grades 1, 2 and 3 conducted in 2003.

A two-stage cluster sample design was used to produce representative data for Mexico - Tijuana. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 100.0%, the class response rate was 98.4%, the student response rate was 83.0%, and the overall response rate was 81.7%. A total of 1,447 students aged 13-15 participated in the Mexico - Tijuana GYTS.

Prevalence

37.4% of students had ever smoked cigarettes (Boy = 40.5%, Girl = 34.6%)
 17.2% currently use any tobacco product (Boy = 18.0%, Girl = 16.3%)
 12.4% currently smoke cigarettes (Boy = 11.9%, Girl = 12.9%)
 7.4% currently use other tobacco products (Boy = 8.8%, Girl = 6.0%)
 23.3% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

19.7% think boys and 34.1% think girls who smoke have more friends
 10.5% think boys and 24.2% think girls who smoke look more attractive

Access and Availability - Current Smokers

19.7% usually smoke at home
 24.9% buy cigarettes in a store
 68.2% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

36.8% live in homes where others smoke in their presence
 51.1% are around others who smoke in places outside their home
 90.8% think smoking should be banned from public places
 81.5% think smoke from others is harmful to them
 41.4% have one or more parents who smoke
 55.1% have most or all friends who smoke

Cessation - Current Smokers

29.7% want to stop smoking
 54.6% tried to stop smoking during the past year
 53.3% have ever received help to stop smoking

Media and Advertising

87.6% saw anti-smoking media messages, in the past 30 days
 88.6% saw pro-cigarette ads on billboards, in the past 30 days
 74.5% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
 10.8% have an object with a cigarette brand logo
 9.6% were offered free cigarettes by a tobacco company representative

School

55.2% had been taught in class, during the past year, about the dangers of smoking
 24.4% had discussed in class, during the past year, reasons why people their age smoke
 47.7% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 17.2% of students currently use any form of tobacco; 12.4% currently smoke cigarettes; 7.4% currently use some other form of tobacco.
- SHS exposure is high – over one-third of the students live in homes where others smoke, and half the students are exposed to smoke around others outside of the home; 2 in 5 students have at least one parent who smokes.
- Eight in 10 students think smoke from others is harmful to them.
- Nine in 10 students think smoking in public places should be banned.
- Three in 10 current smokers want to stop smoking.
- One in 10 students has an object with a cigarette brand logo on it.
- Over 4 in 5 students saw anti-smoking media messages in the past 30 days; almost 9 in 10 students saw pro-cigarette ads on billboards and three-quarters saw pro-tobacco ads in newspapers or magazines in the past 30 days.