



	1 343 500 LIVES CAN BE SAVED BY 2025 BY IMPLEMENTING ALL OF THE WHO "BEST BUYS"						
		NATIONAL TARGET SET		DATA YEAR	MALES	FEMALES	TOTAL
MORTALITY*							
	Premature mortality	✓	Total NCD deaths	2016	3 313 000	2 682 000	5 995 000
	from NCDs	<b>V</b>	Risk of premature death between 30-70 years (%)	2016	27	20	23
P.	Suicide mortality	-	Suicide mortality rate (per 100 000 population)	2016	-	-	16
RISK FACTORS							
	Harmful use of alcohol	✓	Total alcohol per capita consumption, adults aged 15+ (litres of pure alcohol)	2016	9	2	6
K	Physical inactivity	✓	Physical inactivity, adults aged 18+ (%)	2016	24	43	33
	Salt/Sodium intake	✓	Mean population salt intake, adults aged 20+ (g/day)	2010	10	9	9
8	Tobacco use	✓	Current tobacco smoking, adults aged 15+ (%)	2016	20	2	11
	Raised blood pressure	✓	Raised blood pressure, adults aged 18+ (%)	2015	24	23	24
À	Diabetes	✓	Raised blood glucose, adults aged 18+ (%)	2014	8	8	8
À	Obesity	✓	Obesity, adults aged 18+ (%)	2016	3	5	4
			Obesity, adolescents aged 10-19 (%)	2016	2	1	2
<b>%</b>	Ambient air pollution	-	Exceedance of WHO guidelines level for annual PM2.5 concentration (by a multiple of)	2016	-	-	7
à	Household air pollution	-	Population with primary reliance on polluting fuels and technologies (%)	2016	-	-	59

# SELECTED ADULT RISK FACTOR TRENDS




# **OBESITY**






### **RAISED BLOOD PRESSURE**



# NATIONAL SYSTEMS RESPO







Proportion of high risk persons receiving any drug therapy and counselling to prevent heart attacks and strokes (%) Proportion of primary health care centres reported as offering CVD risk stratification

Reported having CVD guidelines that are utilized in at least 50% of health facilities Number of essential NCD medicines reported as

"generally available" Number of essential NCD technologies reported as

% of the population

2017 25% to 50% 2017 Yes



â