

Dichloromethane

Dichloromethane, or methylene chloride, is widely used as a solvent for many purposes, including coffee decaffeination and paint stripping. Exposure from drinking-water is likely to be insignificant compared with that from other sources.

Guideline value	0.02 mg/l (20 µg/l)
Occurrence	Has been found in surface water samples at concentrations ranging from 0.1 to 743 µg/l; levels usually higher in groundwater because volatilization is restricted, with concentrations as high as 3600 µg/l reported; mean concentrations in drinking-water less than 1 µg/l
TDI	6 µg/kg body weight, derived from a NOAEL of 6 mg/kg body weight per day for hepatotoxic effects in a 2-year drinking-water study in rats, using an uncertainty factor of 1000 (100 for interspecies and intraspecies variation and 10 for concern about carcinogenic potential)
Limit of detection	0.3 µg/l by purge-and-trap GC with MS detection (note that dichloromethane vapour readily penetrates tubing during the procedure)
Treatment performance	20 µg/l should be achievable using air stripping
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Assessment date	1993
Principal reference	WHO (2003) <i>Dichloromethane in drinking-water</i>

Dichloromethane is of low acute toxicity. An inhalation study in mice provided conclusive evidence of carcinogenicity, whereas drinking-water studies in rats and mice provided only suggestive evidence. IARC has placed dichloromethane in Group 2B (possible human carcinogen); however, the balance of evidence suggests that it is not a genotoxic carcinogen and that genotoxic metabolites are not formed in relevant amounts in vivo.

1,2-Dichloropropane

1,2-Dichloropropane (CAS No. 78-87-5), or 1,2-DCP, is used as an insecticide fumigant on grain and soil and to control peach tree borers. It is also used as an intermediate in the production of tetrachloroethene and other chlorinated products and as a solvent. 1,2-DCP is relatively resistant to hydrolysis, is poorly adsorbed onto soil and can migrate into groundwater.

Provisional guideline value	0.04 mg/l (40 µg/l)
	The guideline value is provisional owing to limitations of the toxicological database.

12. CHEMICAL FACT SHEETS

Occurrence	Detected in groundwater and drinking-water, usually at concentrations below 20 µg/l, although levels as high as 440 µg/l have been measured in well water
TDI	14 µg/kg body weight based on a LOAEL of 71.4 mg/kg body weight per day (100 mg/kg body weight per day adjusted for daily dosing) for changes in haematological parameters in a 13-week study in male rats, with an uncertainty factor of 5000 (100 for interspecies and intraspecies variation, 10 for use of a LOAEL and 5 to reflect limitations of the database, including the limited data on in vivo genotoxicity and use of a subchronic study)
Limit of detection	0.02 µg/l by purge-and-trap GC with an electrolytic conductivity detector or GC-MS
Treatment performance	1 µg/l should be achievable using GAC
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Assessment date	1998
Principal reference	WHO (2003) <i>1,2-Dichloropropane (1,2-DCP) in drinking-water</i>

1,2-DCP was evaluated by IARC in 1986 and 1987. The substance was classified in Group 3 (not classifiable as to its carcinogenicity to humans) on the basis of limited evidence for its carcinogenicity in experimental animals and insufficient data with which to evaluate its carcinogenicity in humans. Results from in vitro assays for mutagenicity were mixed. The in vivo studies, which were limited in number and design, were negative. In accordance with the IARC evaluation, the evidence from the long-term carcinogenicity studies in mice and rats was considered limited, and it was concluded that the use of a threshold approach for the toxicological evaluation of 1,2-DCP was appropriate.

1,3-Dichloropropane

1,3-Dichloropropane (CAS No. 142-28-9) has several industrial uses and may be found as a contaminant of soil fumigants containing 1,3-dichloropropene. It is rarely found in water.

Reason for not establishing a guideline value	Available data inadequate to permit derivation of health-based guideline value
Assessment date	1993
Principal reference	WHO (2003) <i>1,3-Dichloropropane in drinking-water</i>

1,3-Dichloropropane is of low acute toxicity. There is some indication that it may be genotoxic in bacterial systems. No short-term, long-term, reproductive or developmental toxicity data pertinent to exposure via drinking-water could be located in the literature. The available data are considered insufficient to permit recommendation of a guideline value.