

1,3-Dichloropropene

1,3-Dichloropropene (CAS Nos. 542-75-6 isomer mixture; 10061-01-5 *cis* isomer; 10061-02-6 *trans* isomer) is a soil fumigant, the commercial product being a mixture of *cis* and *trans* isomers. It is used to control a wide variety of soil pests, particularly nematodes in sandy soils. Notwithstanding its high vapour pressure, it is soluble in water at the gram per litre level and can be considered a potential water contaminant.

Guideline value	0.02 mg/l (20 µg/l)
Occurrence	Has been found in surface water and groundwater at concentrations of a few micrograms per litre
Basis of guideline value derivation	Calculated by applying the linearized multistage model to the observation of lung and bladder tumours in female mice in a 2-year gavage study
Limit of detection	0.34 and 0.20 µg/l by purge-and-trap packed column GC using an electrolytic conductivity detector or microcoulometric detector for the <i>cis</i> and <i>trans</i> isomers, respectively
Treatment performance	No information found on removal from water
Assessment date	1993
Principal reference	WHO (2003) <i>1,3-Dichloropropene in drinking-water</i>

1,3 Dichloropropene is a direct-acting mutagen that has been shown to produce forestomach tumours following long term oral gavage exposure in rats and mice. Tumours have also been found in the bladder and lungs of female mice and the liver of male rats. Long term inhalation studies in the rat have proved negative, whereas some benign lung tumours have been reported in inhalation studies in mice. IARC has classified 1,3 dichloropropene in Group 2B (possible human carcinogen).

Dichlorprop

The half-lives for degradation of chlorophenoxy herbicides, including dichlorprop (CAS No. 120-36-5), or 2,4-DP, in the environment are in the order of several days. Chlorophenoxy herbicides are not often found in food.

Guideline value	0.1 mg/l (100 µg/l)
Occurrence	Chlorophenoxy herbicides not frequently found in drinking-water; when detected, concentrations usually no greater than a few micrograms per litre
TDI	36.4 µg/kg body weight, based on a NOAEL of 3.64 mg/kg body weight per day for renal toxicity in a 2-year dietary study in rats, applying an uncertainty factor of 100 (for intraspecies and interspecies variation)
Limit of detection	1 µg/l to 1 mg/l for various methods commonly used for the determination of chlorophenoxy herbicides in water, including solvent extraction, separation by GC, gas-liquid chromatography, thin-layer chromatography or HPLC, with ECD or UV detection