

It should be noted that the lowest reported odour threshold for cyanide in drinking-water is 0.17 mg/l, which is below the short-term health-based value. It is therefore possible that a small number of individuals will detect cyanide by odour at concentrations below the health-based value.

The health-based value relates to total cyanide concentration at the tap, including cyanide from cyanogen chloride in drinking-water as a by-product of disinfection with chlorine. Cyanogen chloride rapidly breaks down to cyanide in the distribution system or when ingested. As the low levels of cyanide normally found in drinking-water are mostly a consequence of the presence of cyanogen chloride, it is not considered necessary to develop a guideline value for long-term exposure to cyanide.

### **Cyanogen chloride**

Cyanogen chloride may be formed as a by-product of chloramination or chlorination of water. It is also formed by the chlorination of cyanide ion present in raw water.

Reason for not establishing a guideline value	Occurs in drinking-water at concentrations well below those of health concern
Assessment date	2009
Principal references	IPCS (2004) <i>Hydrogen cyanide and cyanides</i> WHO (2009) <i>Cyanogen chloride in drinking-water</i>

Cyanogen chloride is rapidly metabolized to cyanide in the body. There are few data available on the oral toxicity of cyanogen chloride.

As cyanogen chloride is unlikely to be found in drinking-water at concentrations that are of health concern, it is considered unnecessary to develop a formal guideline value for cyanogen chloride. Instead, for guidance purposes, a health-based value is derived based on cyanide.

Using a NOAEL for cyanide of 4.5 mg/kg body weight per day for minor changes in the testis in a subchronic study in which rats were exposed through their drinking-water and an uncertainty factor of 100, a TDI for cyanide of 0.045 mg/kg body weight (corresponding to a cyanogen chloride dose of 0.11 mg/kg body weight) can be derived. In view of the minor nature of the changes observed and the NOAEL in a previous chronic study, it is not considered necessary to include an additional uncertainty factor to allow for the length of the study. Further, it appears that a dose that may be toxic in acute poisoning would certainly be tolerated under chronic conditions, owing to efficient detoxification. Assuming a 60 kg adult drinking 2 litres of water per day and allowing 20% of the TDI to come from water because of the potential for exposure to cyanogenic glycosides in food, the health-based value for long-term exposure is 0.3 mg/l for cyanide or 0.6 mg/l for cyanogen chloride (rounded values).

Although low concentrations of cyanide in raw waters will be converted to cyanogen chloride by chlorination, cyanogen chloride may also be formed during the production of chloramines in situ as a residual disinfectant to maintain the hygienic condition of the distribution system. It is important that treatment be optimized to minimize the formation of cyanogen chloride while maintaining adequate chloramine residuals where chloramination is practised.