

Provisional guideline value	0.05 mg/l (50 µg/l)
	The guideline value is designated as provisional on the basis of technical achievability.
Occurrence	Found in groundwater and surface water distribution systems at concentrations up to about 100 µg/l, with mean concentrations below 20 µg/l
Basis of guideline value derivation	Linear multistage model applied to combined data for carcinomas and adenomas in male mice exposed to doses up to 429 mg/kg body weight per day for up to 2 years
Limit of detection	< 0.1–0.4 µg/l by GC with ECD; practical quantification limit 1 µg/l
Treatment performance	Concentrations may be reduced by installing or optimizing coagulation to remove precursors or by controlling the pH during chlorination.
Additional comments	The concentration associated with a 10^{-5} upper-bound excess lifetime cancer risk is 40 µg/l. In some circumstances, however, it may not be possible to adequately disinfect potable water and maintain DCA levels below 40 µg/l, so the provisional guideline value of 50 µg/l is retained.
Assessment date	2004
Principal reference	WHO (2005) <i>Dichloroacetic acid in drinking-water</i>

IARC reclassified DCA as Group 2B (possibly carcinogenic to humans) in 2002, based on the absence of data on human carcinogenicity and sufficient evidence of its carcinogenicity in experimental animals. This classification was based primarily on findings of liver tumours in rats and mice. Genotoxicity data are considered to be inconclusive, particularly at lower doses. Glycogen deposition, peroxisome proliferation, changes in signal transduction pathways and DNA hypomethylation have all been observed following DCA exposure and have been hypothesized to be involved in its carcinogenicity. However, the available data are not sufficient to establish a cancer mode of action with reasonable certainty, especially at the very low exposure levels expected to apply to humans ingesting chlorinated drinking-water. Recent data suggest that there may be more than one mechanism leading to tumours, as altered hepatic foci from treated mice were found to have three different types of cellular characteristics.

Dichlorobenzenes (1,2-dichlorobenzene, 1,3-dichlorobenzene, 1,4-dichlorobenzene)

The dichlorobenzenes (DCBs) are widely used in industry and in domestic products such as odour-masking agents, chemical dyestuffs and pesticides. Sources of human exposure are predominantly air and food.

Guideline values	<i>1,2-Dichlorobenzene</i> : 1 mg/l (1000 µg/l)
	<i>1,4-Dichlorobenzene</i> : 0.3 mg/l (300 µg/l)
Occurrence	Have been found in raw water sources at levels as high as 10 µg/l and in drinking-water at concentrations up to 3 µg/l; much higher concentrations (up to 7 mg/l) present in contaminated groundwater

TDIs	<p><i>1,2-Dichlorobenzene</i>: 429 µg/kg body weight, based on a NOAEL of 60 mg/kg body weight per day for tubular degeneration of the kidney identified in a 2-year mouse gavage study, adjusting for daily dosing and using an uncertainty factor of 100 (for interspecies and intraspecies variation)</p> <p><i>1,4-Dichlorobenzene</i>: 107 µg/kg body weight, based on a LOAEL of 150 mg/kg body weight per day for kidney effects identified in a 2-year rat study, adjusting for daily dosing and using an uncertainty factor of 1000 (100 for interspecies and intraspecies variation and 10 for the use of a LOAEL instead of a NOAEL and the carcinogenicity end-point)</p>
Limit of detection	0.01–0.25 µg/l by gas–liquid chromatography with ECD; 3.5 µg/l by GC using a photoionization detector
Treatment performance	0.01 mg/l should be achievable using air stripping
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Additional comments	Guideline values for both 1,2- and 1,4-DCB far exceed their lowest reported taste thresholds in water of 1 and 6 µg/l, respectively.
Assessment date	1993
Principal reference	WHO (2003) <i>Dichlorobenzenes in drinking-water</i>
Reason for not establishing a guideline value	Available data inadequate to permit derivation of health-based guideline value for <i>1,3-dichlorobenzene</i>
Assessment date	1993
Principal reference	WHO (2003) <i>Dichlorobenzenes in drinking-water</i>

1,2-Dichlorobenzene

1,2-DCB is of low acute toxicity by the oral route of exposure. Oral exposure to high doses of 1,2-DCB affects mainly the liver and kidneys. The balance of evidence suggests that 1,2-DCB is not genotoxic, and there is no evidence for its carcinogenicity in rodents.

1,3-Dichlorobenzene

There are insufficient toxicological data on this compound to permit a guideline value to be proposed, but it should be noted that it is rarely found in drinking-water.

1,4-Dichlorobenzene

1,4-DCB is of low acute toxicity, but there is evidence that it increases the incidence of renal tumours in rats and of hepatocellular adenomas and carcinomas in mice after long-term exposure. IARC has placed 1,4-DCB in Group 2B (possibly carcinogenic to humans). 1,4-DCB is not considered to be genotoxic, and the relevance for humans of the tumours observed in experimental animals is doubtful.