

Mecoprop

The half-lives for degradation of chlorophenoxy herbicides, including mecoprop (CAS No. 93-65-2; 7085-19-0 racemic mixture), also known as 2(2-methyl-chlorophenoxy) propionic acid or MCPA, in the environment are in the order of several days. Chlorophenoxy herbicides are not often found in food.

Guideline value	0.01 mg/l (10 µg/l)
Occurrence	Chlorophenoxy herbicides not frequently found in drinking-water; when detected, concentrations usually no greater than a few micrograms per litre
TDI	3.33 µg/kg body weight, based on a NOAEL of 1 mg/kg body weight for effects on kidney weight in 1- and 2-year studies in rats, with an uncertainty factor of 300 (100 for interspecies and intraspecies variation and 3 for limitations in the database)
Limit of detection	0.01 µg/l by GC-MS; 0.01–0.02 µg/l by GC with ECD
Treatment performance	0.1 µg/l should be achievable using GAC or ozonation
Guideline value derivation	<ul style="list-style-type: none"> • allocation to water 10% of TDI • weight 60 kg adult • consumption 2 litres/day
Assessment date	1993
Principal reference	WHO (2003) <i>Chlorophenoxy herbicides (excluding 2,4-D and MCPA) in drinking-water</i>

Chlorophenoxy herbicides, as a group, have been classified in Group 2B (possibly carcinogenic to humans) by IARC. However, the available data from studies in exposed populations and experimental animals do not permit assessment of the carcinogenic potential to humans of any specific chlorophenoxy herbicide. Therefore, drinking-water guidelines for these compounds are based on a threshold approach for other toxic effects. Effects of dietary administration of mecoprop in short-term and long-term studies include decreased relative kidney weight (rats and dogs), increased relative liver weight (rats), effects on blood parameters (rats and dogs) and depressed body weight gain (dogs).

Mercury

Mercury is used in the electrolytic production of chlorine, in electrical appliances, in dental amalgams and as a raw material for various mercury compounds. Methylation of inorganic mercury has been shown to occur in fresh water and in seawater, although almost all mercury in uncontaminated drinking-water is thought to be in the form of Hg²⁺. Thus, it is unlikely that there is any direct risk of the intake of organic mercury compounds, especially of alkylmercurials, as a result of the ingestion of drinking-water. However, there is a possibility that methylmercury will be converted into inorganic mercury. Food is the main source of mercury in non-occupationally

exposed populations; the mean dietary intake of mercury in various countries ranges from 2 to 20 µg/day per person.

Guideline value	0.006 mg/l (6 µg/l) for inorganic mercury
Occurrence	Mercury is present in the inorganic form in surface water and groundwater at concentrations usually below 0.5 µg/l, although local mineral deposits may produce higher levels in groundwater
TDI	2 µg/kg body weight for inorganic mercury based on a NOAEL of 0.23 mg/kg body weight per day for kidney effects in a 26-week study in rats and applying an uncertainty factor of 100 (for interspecies and intraspecies variation) after adjusting for daily dosing
Limit of detection	0.05 µg/l by cold vapour AAS; 0.6 µg/l by ICP; 5 µg/l by flame AAS
Treatment performance	It should be possible to achieve a concentration below 1 µg/l by treatment of raw waters that are not grossly contaminated with mercury using methods that include coagulation/sedimentation/filtration, PAC and ion exchange.
Guideline value derivation	
• allocation to water	10% of TDI
• weight	60 kg adult
• consumption	2 litres/day
Additional comments	A similar TDI may be obtained by applying an uncertainty factor of 1000 (an additional uncertainty factor of 10 for adjustment from a LOAEL to a NOAEL) to the LOAEL for renal effects of 1.9 mg/kg body weight per day in a 2-year NTP study in rats. The current guideline value applies to inorganic mercury, which is the form found in drinking-water, whereas the previous guideline value applied to total (inorganic and organic) mercury.
Assessment date	2004
Principal references	IPCS (2003) <i>Elemental mercury and inorganic mercury compounds</i> WHO (2005) <i>Mercury in drinking-water</i>

The toxic effects of inorganic mercury compounds are seen mainly in the kidney in both humans and laboratory animals following short-term and long-term exposure. In rats, effects include increased absolute and relative kidney weights, tubular necrosis, proteinuria and hypoalbuminaemia. In humans, acute oral poisoning results primarily in haemorrhagic gastritis and colitis; the ultimate damage is to the kidney. The overall weight of evidence is that mercury(II) chloride has the potential to increase the incidence of some benign tumours at sites where tissue damage is apparent and that it possesses weak genotoxic activity but does not cause point mutations.

Methoxychlor

Methoxychlor (CAS No. 72-43-5) is an insecticide used on vegetables, fruit, trees, fodder and farm animals. It is poorly soluble in water and highly immobile in most agricultural soils. Under normal conditions of use, methoxychlor does not seem to be of environmental concern. Daily intake from food and air is expected to be below 1