

TOBACCO CONTROL FACT SHEET

Kazakhstan

Health impact of tobacco control policies
in line with the WHO Framework Convention
on Tobacco Control (FCTC)



Based on the current level of adult smoking in Kazakhstan (1), premature deaths attributable to smoking are projected to be more than 1.4 million of the 2.8 million smokers alive today (Table 1) and may increase in the absence of stronger policies.

TABLE 1.

Initial smoking prevalence and projected premature deaths

Smoking prevalence (%)		Smokers (n)	Projected premature deaths of current smokers (n)			
Male	Female	Total	Male ^a	Female ^a	Total ^a	Total ^b
42.4	4.5	2 845 500	1 272 000	150 750	1 422 750	924 788

^a Premature deaths are based on relative risks from large-scale studies of high-income countries.

^b Premature deaths are based on relative risks from large-scale studies of low- and middle-income countries.

Source: WHO (1).

Key findings

Within 15 years, the effects of individual tobacco control policies when fully implemented in line with the WHO FCTC (2) are projected to reduce smoking prevalence by:

- 26.5% by increasing excise cigarette taxes from the current level of 28.57% to 75% and prevent much smoking among young people;
- 6.3% with stronger enforcement of comprehensive smoke-free laws;
- 6.6% by banning most forms of direct and indirect advertising to create a comprehensive ban on advertising, promotion and sponsorship with enforcement;
- 9% by requiring that strong graphic health warnings be added to tobacco products;
- 3.3% by increasing from moderate provision to a well publicized and comprehensive tobacco-cessation policy; and
- 6.3% by increasing awareness of the harms of tobacco use through a high-level media campaign.

With this stronger set of policies and consistent with the WHO FCTC (2), smoking prevalence can be reduced by 36% within five years, 47% within 15 years and 56% within 40 years. More than 800 000 deaths could be averted in the long term (Table 2). The SimSmoke tobacco control model (3) incorporates synergies in implementing multiple policies (such as strong media campaigns with smoke-free laws and tobacco-cessation policies).

TABLE 2.

Effect of tobacco control policies (individual and combined) on initial smoking prevalence and smoking-attributable deaths

Tobacco control policy	Relative change in smoking prevalence (%)			Reduction in smoking-attributable deaths in 40 years (n)			
	5 years	15 years	40 years	Male ^a	Female ^a	Total ^a	Total ^b
Protect through smoke-free laws	–5.5	–6.3	–6.8	87 087	10 321	97 408	63 315
Offer tobacco-cessation services	–1.9	–3.3	–4.8	60 812	7 207	68 019	44 212
Mass media campaigns	–5.5	–6.3	–6.6	83 952	9 950	93 902	61 036
Warnings on cigarette packages	–6.0	–9.0	–12.0	152 640	18 090	170 730	110 975
Enforce marketing restrictions	–5.5	–6.6	–7.2	90 948	10 779	101 727	66 122
Raise cigarette taxes	–17.7	–26.5	–35.4	449 752	53 302	503 054	326 985
Combined policies	–35.9	–47.0	–56.2	715 406	84 786	800 192	520 125

^a Smoking-attributable deaths are based on relative risks from large-scale studies of high-income countries.

^b Smoking-attributable deaths are based on relative risks from large-scale studies of low- and middle-income countries.

→ Monitor tobacco use

The prevalence of current adult smokers (15 and above) in Kazakhstan in 2014 was 22.4% (men: 42.4%; women: 4.5%) (1).

→ Protect people from tobacco smoke

Health-care, government and education facilities (including universities), indoor offices and workplaces, and cafes, pubs and bars are completely smoke-free in Kazakhstan (Table 3). Smoking violations incur fines for the establishment and patron, but no funds are dedicated to enforcement and no system is in place for citizen complaints and further investigations (4).

TABLE 3.

Complete smoke-free indoor public places

Health-care facilities	Education facilities (except universities)	Universities	Government facilities	Indoor offices and workplaces	Restaurants	Cafes, pubs and bars	Public transport	All other indoor public places
✓	✓	✓	✓	✓	–	✓	–	–

Source: WHO (4).

✓ = completely smoke-free. – = not completely smoke-free.

→ Offer help to quit tobacco use

Smoking-cessation services are available in some health clinics and primary care facilities and in offices of health professionals, with costs fully covered by the national health service or national health insurance. Cessation support can also be accessed at hospitals, where the cost is partially covered, and in the community (without cost coverage). Nicotine replacement therapy can be purchased over the counter in a pharmacy without a prescription, but is not cost-covered. A toll-free quit line is available (4).

→ Warn about the dangers of tobacco

Health warnings are legally mandated to cover 40% of the front and rear of the principal display area, with 12 such warnings approved by law. The position of health warnings on packages rotates and the messages are written in the principal language(s) of the country. The law sets the font size/style and colour for package warnings and mandates that the warnings include a photograph or graphic. It also mandates that warnings appear on each package and any outside packaging and labelling used in retail sale (4).

→ Enforce bans on tobacco advertising, promotion and sponsorship

Through a law adopted in 2003 and amended in 2007 (5), Kazakhstan has bans in place on all forms of direct and some forms of indirect advertising (Table 4). The law requires fines for violations of these bans (4).

TABLE 4.

Bans on direct and indirect advertising

Direct advertising		Indirect advertising	
National television and radio	✓	Free distribution in mail or through other means	✓
International television and radio	✓	Promotional discounts	—
Local magazines and newspapers	✓	Non-tobacco products identified with tobacco brand names	—
International magazines and newspapers	✓	Appearance of tobacco brands in television and/or films (product placement)	✓
Billboards and outdoor advertising	✓	Appearance of tobacco products in television and/or films	—
Advertising at point of sale	✓	Sponsored events	—
Advertising on the Internet	✓	Tobacco products display at point of sale	—

Source: WHO (4).

✓ = banned. — = not banned.

Kazakhstan does not have:

- bans on tobacco companies/tobacco industry publicizing their activities;
- bans on entities other than tobacco companies/tobacco industry publicizing activities of the tobacco companies;
- bans on tobacco companies funding or making contributions (including in-kind contributions) to smoking-prevention media campaigns, including those directed at young people; and
- a requirement to present prescribed anti-tobacco advertisements before, during or after the broadcasting or showing of any visual entertainment (4).

→ Raise taxes on tobacco

A pack of cigarettes in Kazakhstan costs 210 KZT¹ (US\$ 1.15), of which 39.29% is tax (10.71% is value-added tax and 28.57% excise taxes) (4).

¹ The currency code is according to International Organization for Standardization, ISO 4217 currency names and code elements.

About the SimSmoke model

The abridged version of the SimSmoke tobacco control model, developed by David Levy of Georgetown University, United States of America, projects the reduction in smoking prevalence and smoking-attributable deaths as a result of implementing tobacco control policies (individually and in combination) (3). Specifically, the model projects the effects from:

- protecting from second-hand smoke through stronger smoke-free laws
- offering greater access to smoking-cessation services
- placing warnings on tobacco packages and other media/educational programmes
- enforcing bans on advertising, promotion and sponsorship
- raising cigarette prices through higher cigarette taxes (6).

Data on smoking prevalence among adults for the SimSmoke model were taken from the most recent nationally representative survey covering a wide age range; data on tobacco control policies were taken from the 2015 WHO report on the global tobacco epidemic (4).

Funding

This fact sheet was made possible by funding from the Government of the Russian Federation.

References

1. Global Adult Tobacco Survey. Fact sheet. Kazakhstan 2014. Geneva: World Health Organization; 2014 (<http://www.who.int/tobacco/surveillance/survey/gats/kazfactsheet.pdf>, accessed 27 December 2016).
2. WHO Framework Convention on Tobacco Control [website]. Geneva: World Health Organization; 2016 (<http://www.who.int/fctc/en/>, accessed 27 December 2016).
3. Levy DT, Fouad H, Levy J, Dragomir A, El Awa F. Application of the abridged SimSmoke model to four eastern Mediterranean countries. *Tob Control* 2016; 25(4):413–21. doi:10.1136/tobaccocontrol-2015-052334.
4. WHO report on the global tobacco epidemic, 2015: raising taxes on tobacco. Geneva: World Health Organization; 2015 (http://www.who.int/tobacco/global_report/2015/en/, accessed 27 December 2016).
5. Tobacco control database for the WHO European Region [online database]. Copenhagen: WHO Regional Office for Europe; 2016 (<http://data.euro.who.int/tobacco/>, accessed 27 December 2016).
6. MPOWER. In: Tobacco free initiative (TFI) [website]. Geneva: World Health Organization; 2016 (<http://www.who.int/tobacco/mpower/en/>, accessed 27 December 2016).

Acknowledgements

Data analysis: David Levy and Jeffrey Levy, Georgetown University, Washington (DC), United States of America

Text: Kristina Mauer-Stender, Nataliia Toropova, Elizaveta Lebedeva, WHO Regional Office for Europe

Text Editing: Alex Mathieson, Edinburgh, United Kingdom

Graphic design: Carli Hyland, Hill+Knowlton Strategies, Copenhagen, Denmark